

DAFTAR PUSTAKA

- AJCC. (2018). *AJCC Cancer Staging Manual 8th Edition* (M. B. Amin, S. B. Edge, F. L. Greene, D. R. Byrd, R. K. Brookland, M. K. Washington, J. E. Gershenwald, C. C. Compton, K. R. Hess, D. C. Sullivan, J. M. Jessup, J. D. Brierley, L. E. Gaspar, R. L. Schilsky, C. M. Balch, D. P. Winchester, E. A. Asare, M. Madera, D. M. Gress, & L. R. Meyer, Eds.). Springer International Publishing. <https://doi.org/10.1007/978-3-319-40618-3>
- Almomani R, Khanfar M, Bodoor K, et al. Evaluation of patched-1 protein expression level in low risk and high risk basal cell carcinoma subtypes. *Asian Pac J Cancer Prev*. 2019;20(9):2851–2857. doi:10.31557/APJCP.2019.20.9.2851
- American Cancer Society. Cancer facts and figures 2000. www.cancer.org.2001 (Accessed on March 08, 2005).
- American Cancer Society. Key Statistics for Basal and Squamous Cell Skin Cancers. <https://www.cancer.org/cancer/basal-and-squamous-cell-skin-cancer/about/key-statistics.html> (Accessed on October 22, 2021).
- Ammasaignandan V, Shariff VNSA, Ramesh A. Basal cell carcinoma- a prospective clinico epidemiological and pathological study. *Int J Res Med Sci*. 2017;5(6):2712.
- Aszterbaum, M., Rothman, A., Fisher, M., Xie, J., Bonifas, J.M., Zhang, X., Epstein Jr, E.H., Johnson, R.L. and Scott, M.P., 1998. Identification of mutations in the human PATCHED gene in sporadic basal cell carcinomas and in patients with the basal cell nevus syndrome. *Journal of investigative dermatology*, 110(6), pp.885-888.
- Athar M, Li C, Kim AL, Spiegelman VS, Bickers DR. Sonic hedgehog signaling in Basal cell nevus syndrome. *Cancer Res*. 2014; 74:4967–4975.
- Athar M, Tang X, Lee JL, Kopelovich L, Kim AL. Hedgehog signaling in skin development and cancer. *Exp Dermatol*. 2006; 15:667–677.
- Atwood SX, Chang AL, Oro AE. Hedgehog pathway inhibition and the race against tumor evolution. *J Cell Biol*. 2012; 199:193–197.
- Atwood, S.X., Sarin, K.Y., Whitson, R.J., Li, J.R., Kim, G., Rezaee, M., Ally, M.S., Kim, J., Yao, C., Chang, A.L.S. and Oro, A.E., 2015. Smoothed variants explain the majority of drug resistance in basal cell carcinoma. *Cancer cell*, 27(3), pp.342-353.
- Berezikov E. Evolution of microRNA diversity and regulation in animals. *Nat Rev Genet*. 2011; 12:846–860.
- Bertoli G, Cava C, Castiglioni I. MicroRNAs as Biomarkers for Diagnosis, Prognosis and Theranostics in Prostate Cancer. *Int J Mol Sci*. 2016; 17:421.
- Boldt K, van Reeuwijk J, Lu Q, et al. An organelle-specific protein landscape identifies novel diseases and molecular mechanisms. *Nat Commun*. 2016; 7:11491.
- Bonilla X, Parmentier L, King B, et al. Genomic analysis identifies new drivers and progression pathways in skin basal cell carcinoma. *Nat Genet*. 2016; 48:398–406.
- Bossi, P., Perrone, F., Cortellazzi, B., Licitra, L., Dal Col, V., Laurini, E., Fermeiglia, M. and Pricl, S., 2013. Abstract PR05: Resistance to Hedgehog inhibitor through Smoothed receptor mutation in basal cell carcinoma. *Molecular Cancer Therapeutics*, 12(11_Supplement), pp.PR05-PR05.
- Bresler, SCA-Ohoo, Padwa, BL., Granter, SR. Nevroid Basal Cell Carcinoma Syndrome (Gorlin Syndrome). *Head Neck Pathol*. 2016; 10:119–124.
- Briscoe, J. and Thérond, P.P. (2013) ‘The mechanisms of hedgehog signalling and its roles in development and disease’, *Nature Reviews Molecular Cell Biology*, 14(7), pp. 416–429. doi:10.1038/nrm3598.
- Campione, E., Di Prete, M., Lozzi, F., Lanna, C., Spallone, G., Mazzeo, M., Cosio, T.,

- Rapanotti, C., Dika, E., Gaziano, R. and Orlandi, A., 2020. High-risk recurrence basal cell carcinoma: focus on hedgehog pathway inhibitors and review of the literature. *Chemotherapy*, 65(1-2), pp.2-10.
- Chaudhary SC, Tang X, Arumugam A, et al. Shh and p50/Bcl3 signaling crosstalk drives pathogenesis of BCCs in gorlin syndrome. *Oncotarget*. 2015; 6:36789–36814.
- Christenson LJ, Borrowman TA, Vachon CM, et al. Incidence of basal cell and squamous cell carcinomas in a population younger than 40 years. *JAMA* 2005; 294:681.
- Chuang TY, Popescu A, Su WP, Chute CG. Basal cell carcinoma. A population-based incidence study in Rochester, Minnesota. *J Am Acad Dermatol* 1990; 22:413.
- Ciążyńska M, Narbutt J, Woźniacka A, Lesiak A. Trends in basal cell carcinoma incidence rates: a 16-year retrospective study of a population in central Poland. *Postep dermatologii i Alergol*. 2018;35(1):47–52.
- Colavito SA, Zou MR, Yan Q, Nguyen DX, Stern DF. Significance of glioma-associated oncogene homolog 1 (GLI1) expression in claudin-low breast cancer and crosstalk with the nuclear factor kappa-light-chain-enhancer of activated B cells (NFkappaB) pathway. *Breast Cancer Res*. 2014; 16:444.
- Couzens AL, Knight JD, Kean MJ, et al. Protein interaction network of the mammalian Hippo pathway reveals mechanisms of kinase-phosphatase interactions. *Sci Signal*. 2013; 6:rs15.
- Crowson, A.N., 2006. Basal cell carcinoma: biology, morphology and clinical implications. *Modern pathology*, 19, pp.S127-S147.
- Croyle MJ, Lehman JM, O'Connor AK, et al. Role of epidermal primary cilia in the homeostasis of skin and hair follicles. *Development*. 2011; 138:1675–1685.
- Dahlén A, Mertens F, Mandahl N, Panagopulos. 2005. GLI1 (glioma-associated oncogene homolog-1). *Atas Genet Cytogenet Oncol Haematol*; 9(2):136-8
- Dalmay T. MicroRNAs and cancer. *J Intern Med*. 2008; 263:366–375. [PubMed: 18298485]
77. Sand M, Gambichler T, Skrygan M, et al. Expression levels of the microRNA processing enzymes Drosha and Dicer in epithelial skin cancer. *Cancer Invest*. 2010; 28:649–653.
- Daya-Grosjean, L. and Sarasin, A., 2000. UV-specific mutations of the human patched gene in basal cell carcinomas from normal individuals and xeroderma pigmentosum patients. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 450(1-2), pp.193-199.
- De Craene B, Denecker G, Vermassen P, et al. Epidermal Snail expression drives skin cancer initiation and progression through enhanced cytoprotection, epidermal stem/progenitor cell expansion and enhanced metastatic potential. *Cell Death Differ*. 2014; 21:310–320.
- Demers AA, Nugent Z, Mihalcioiu C, et al. Trends of nonmelanoma skin cancer from 1960 through 2000 in a Canadian population. *J Am Acad Dermatol* 2005; 53:320.
- Deng, L.J., Jia, M., Luo, S.Y., Li, F.Z. and Fang, S., 2022. Expression of hedgehog signaling pathway proteins in basal cell carcinoma: clinicopathologic study. *Clinical, Cosmetic and Investigational Dermatology*, pp.2353-2361.
- Dennler S, Andre J, Verrecchia F, Mauviel A. Cloning of the human GLI2 Promoter: transcriptional activation by transforming growth factor-beta via SMAD3/beta-catenin cooperation. *J Biol Chem*. 2009; 284:31523–31531.
- Di Magno L, Coni S, Di Marcotullio L, Canettieri G. Digging a hole under Hedgehog: downstream inhibition as an emerging anticancer strategy. *Biochim Biophys Acta*. 2015; 1856:62–72.
- DiDonato JA, Mercurio M, Karin M. NF-kappaB and the link between inflammation and cancer. *Immunol Rev*. 2012; 246:379–400.
- Dong J, Feldmann G, Huang J, et al. Elucidation of a universal size-control mechanism in *Drosophila* and mammals. *Cell*. 2007; 130:1120–1133.

- Dorn KV, Hughes CE, Rohatgi R. A Smoothened-Evc2 complex transduces the Hedgehog signal at primary cilia. *Dev Cell*. 2012; 23:823–835.
- Drummond IA. Cilia functions in development. *Curr Opin Cell Biol*. 2012; 24:24–30. [PubMed: 2226236] 39. Brooks ER, Wallingford JB. Multiciliated cells. *Curr Biol*. 2014; 24:R973–R982.
- Elder, D.E., Massi, D., Scolyer, R.A. and Willemze, R., 2018. WHO classification of skin tumours. International Agency for Research on cancer 2018.
- Epstein, E.H., 2008. Basal cell carcinomas: attack of the hedgehog. *Nature Reviews Cancer*, 8(10), pp.743-754.
- Ezratty EJ, Stokes N, Chai S, Shah AS, Williams SE, Fuchs E. A role for the primary cilium in Notch signaling and epidermal differentiation during skin development. *Cell*. 2011; 145:1129–1141s.
- Fernandez-Flores, A. and de la Torre, F.R., 2020. Considerations on Lateral and Deep Surgical Margins in Resected Skin Tumors. *Actas dermo-sifiliograficas*, 111(6), pp.533-535.
- Garcia-Gonzalo FR, Phua SC, Roberson EC, et al. Phosphoinositides regulate ciliary protein trafficking to modulate hedgehog signaling. *Dev Cell*. 2015; 34:400–409.
- Gerdes JM, Liu Y, Zaghloul NA, et al. Disruption of the basal body compromises proteasomal function and perturbs intracellular Wnt response. *Nat Genet*. 2007; 39:1350–1360
- Goetz SC, Anderson KV. The primary cilium: a signalling centre during vertebrate development. *Nat Rev Genet*. 2010; 11:331–344.
- Goldenberg G, Karagiannis T, Palmer JB, et al. Incidence and prevalence of basal cell carcinoma (BCC) and locally advanced BCC (LABCC) in a large commercially insured population in the United States: a retrospective cohort study. *J Am Acad Dermatol*. 2016; 75:957–966. e952.
- Gorlin, R.J. (1995) 'Nevoid basal cell carcinoma syndrome', *Dermatologic Clinics*, 13(1), pp. 113–125. doi:10.1016/s0733-8635(18)30114-1.
- Green A, Battistutta D, Hart V, et al. Skin cancer in a subtropical Australian population: incidence and lack of association with occupation. The Nambour Study Group. *Am J Epidemiol* 1996; 144:1034.
- Habbig S, Bartram MP, Muller RU, et al. NPHP4, a cilia-associated protein, negatively regulates the Hippo pathway. *J Cell Biol*. 2011; 193:633–642.
- Habbig S, Bartram MP, Sagmuller JG, et al. The ciliopathy disease protein NPHP9 promotes nuclear delivery and activation of the oncogenic transcriptional regulator TAZ. *Hum Mol Genet*. 2012; 21:5528–5538
- Hahn H, Wicking C, Zaphiropoulos PG, et al. Mutations of the human homolog of Drosophila patched in the nevoid basal cell carcinoma syndrome. *Cell* 1996; 85:841.
- Hajizadeh, E., Abdollahi, M., Baghestani, A. R., & Haghighat, S. (2018). Prognostic cut point for breast cancer age of diagnosis. *International Journal of Cancer Management*, 11(5). <https://doi.org/10.5812/ijcm.9291>
- Hanahan, D. and Weinberg, R.A., 2000. The hallmarks of cancer. *cell*, 100(1), pp.57-70.
- Hanahan, D., 2022. Hallmarks of cancer: new dimensions. *Cancer discovery*, 12(1), pp.31-46.
- Hannuksela-Svahn A, Pukkala E, Karvonen J. Basal cell skin carcinoma and other nonmelanoma skin cancers in Finland from 1956 through 1995. *Arch Dermatol* 1999; 135:781.
- Hatta, N., Hirano, T., Kimura, T., Hashimoto, K., Mehregan, D.R., Ansai, S., Takehara, K. and Takata, M., 2005. Molecular diagnosis of basal cell carcinoma and other basaloid cell neoplasms of the skin by the quantification of Gli1 transcript levels. *Journal of cutaneous pathology*, 32(2), pp.131-136.
- Heffelfinger C, Ouyang Z, Engberg A, et al. Correlation of global microRNA expression with basal cell carcinoma subtype. G3 (Bethesda). 2012; 2:279–286.

- Hilgendorf KI, Johnson CT, Jackson PK. The primary cilium as a cellular receiver: organizing ciliary GPCR signaling. *Curr Opin Cell Biol.* 2016; 39:84–92.
- Inamura K, Ishikawa Y. MicroRNA in lung cancer: novel biomarkers and potential tools for treatment. *J Clin Med.* 2016;5.
- Itesako T, Seki N, Yoshino H, et al. The microRNA expression signature of bladder cancer by deep sequencing: the functional significance of the miR-195/497 cluster. *PloS ONE.* 2014; 9:e84311.
- Jaju PD, Nguyen CB, Mah AM, et al. Mutations in the Kinetochore Gene KNSTRN in Basal Cell Carcinoma. *J Invest Dermatol.* 2015; 135:3197–3200.
- Janjua, et al. Research Article: Basal Cell Carcinoma of the Head and Neck Region: An Analysis of 171 Cases. *Journal of Skin Cancer.* 2012. Didownload dari : [Basal Cell Carcinoma of the Head and Neck Region: An Analysis of 171 Cases - PMC \(nih.gov\)](#)
- Javelaud, D., Alexaki, V.I., Dennler, S., Mohammad, K.S., Guise, T.A. and Mauviel, A., 2011. TGF- β /SMAD/GLI2 signaling axis in cancer progression and metastasis. *Cancer research*, 71(17), pp.5606-5610.
- Jemal AM, Gracia M. Ward E, Thun MJ. Global cancer incidence (surveillance, epidemiology and End result databas) in Devita VT, Lawrence TS, Rosenberg SA, editors. Devita, Hellman & Rosenberg”scancer. Principles and practice of Oncologi.8th ed.:Lippincott Williams&Wilks ;2008 p.25-76
- Jeng, K.S., Sheen, I.S., Leu, C.M., Tseng, P.H. and Chang, C.F., 2020. The role of smoothened in cancer. *International Journal of Molecular Sciences*, 21(18), p.6863.
- Jiang K, Liu Y, Fan J, et al. PI(4)P promotes phosphorylation and conformational change of smoothened through interaction with its C-terminal tail. *PLoS Biol.* 2016; 14:e1002375.
- Johnson MR, Hageboutros A, Wang K, et al. Life-threatening toxicity in a dihydropyrimidine dehydrogenase-deficient patient after treatment with topical 5-fluorouracil. *Clin Cancer Res* 2012; 5:2006.
- Jung B, Padula D, Burtscher I, et al. Pitchfork and Gprasp2 target smoothened to the primary cilium for hedgehog pathway activation. *PLoS ONE.* 2016; 11:e0149477.
- Kasper M, Schnidar H, Neill GW, et al. Selective modulation of Hedgehog/GLI target gene expression by epidermal growth factor signaling in human keratinocytes. *Mol Cell Biol.* 2012; 26:6283–6298.
- Kauvar, A.N. et al. (2015) ‘Consensus for nonmelanoma skin cancer treatment’, *Dermatologic Surgery*, 41(5), pp. 550–571. doi:10.1097/dss.0000000000000296.
- Kevinsyah, A.P., 2014. Profil Histopatologi Karsinoma Sel Basal Di Instalasi Patologi Anatomi RSUP Dr Sardjito Yogyakarta Tahun 2012-2013 (Doctoral dissertation, Universitas Gadjah Mada).
- Kim HS, Kim YS, Lee C, Shin MS, Kim JW, Jang BG. Expression profile of sonic hedgehog signaling-related molecules in basal cell carcinoma. *PLoS One.* 2019;14(11):e0225511. doi:10.1371/journal.pone.0225511
- Kim, D.P., Kus, K.J. and Ruiz, E., 2019. Basal cell carcinoma review. *Hematology/Oncology Clinics*, 33(1), pp.13-24.
- Kim, H.S., Kim, Y.S., Lee, C., Shin, M.S., Kim, J.W. and Jang, B.G., 2019. Expression profile of sonic hedgehog signaling-related molecules in basal cell carcinoma. *PLoS One*, 14(11), p.e0225511.
- KM, O.A.H. and Hu, Z., 1997. Basal cell carcinomas in mice overexpressing Sanie Hedgehog. *Science*, 276, pp.817-21.
- Kong JH, Yang L, Dessaud E, et al. Notch activity modulates the responsiveness of neural progenitors to sonic hedgehog signaling. *Dev Cell.* 2015; 33:373–387.
- Kovalenko A, Kim JC, Kang TB, et al. Caspase-8 deficiency in epidermal keratinocytes triggers an inflammatory skin disease. *J Exp Med.* 2009; 206:2161–2177.

- Kumar V, Abbas A, Aster J. Robbins basic pathology. 10th ed. Philadelphia: Elsevier; 2018.
- Kurek D, Garinis GA, van Doorninck JH, van der Wees J, Grosveld FG. Transcriptome and phenotypic analysis reveals Gata3-dependent signaling pathways in murine hair follicles. *Development*. 2007; 134:261–272.
- Kuzhandaivel A, Schultz SW, Alkhori L, Alenius M. Cilia-mediated hedgehog signaling in *Drosophila*. *Cell Rep*. 2014; 7:672–680.
- Kyrgidis, A., Vahtsevanos, K., Tzellos, T.G., Xirou, P., Kitikidou, K., Antoniadis, K., Zouboulis, C.C. and Triaridis, S., 2010. Clinical, histological and demographic predictors for recurrence and second primary tumours of head and neck basal cell carcinoma. A 1062 patient-cohort study from a tertiary cancer referral hospital. *European Journal of Dermatology*, 20(3), pp.276-282.
- Lacour, J.P., 2002. Carcinogenesis of basal cell carcinomas: genetics and molecular mechanisms. *British Journal of Dermatology*, 146(s61), pp.17-19.
- Lara, F., Santamaría, J.R. and Garbers, L.E.F.D.M., 2017. Recurrence rate of basal cell carcinoma with positive histopathological margins and related risk factors. *Anais Brasileiros de Dermatologia*, 92, pp.58-62.
- Lehman JM, Laag E, Michaud EJ, Yoder BK. An essential role for dermal primary cilia in hair follicle morphogenesis. *J Invest Dermatol*. 2009; 129:438–448.
- Lemeshow, S., Hosmer Jr, D.W., Klar, J. and Lwanga, S.K., 1990. Part 1: statistical methods for sample size determination. *Adequacy of Sample Size in Health Studies*, 247.
- Lewin J, Carucci J. Advances in the management of basal cell carcinoma. *F1000Prime Rep*. 2015;7(April 2016).
- Li, C., Chi, S. and Xie, J., 2011. Hedgehog signaling in skin cancers. *Cellular signalling*, 23(8), pp.1235-1243.
- Lin, N.U., Vanderplas, A., Hughes, M.E., Theriault, R.L., Edge, S.B., Wong, Y.N., Blayney, D.W., Niland, J.C., Winer, E.P. and Weeks, J.C., 2012. Clinicopathologic features, patterns of recurrence, and survival among women with triple-negative breast cancer in the National Comprehensive Cancer Network. *Cancer*, 118(22), pp.5463-5472.
- Liu Y, Snedecor ER, Choi YJ, et al. Gorab is required for dermal condensate cells to respond to hedgehog signals during hair follicle morphogenesis. *J Invest Dermatol*. 2016; 136:378–386.
- Loho, L.L. and Durry, M.F., 2013. Basalioma. *Jurnal Biomedik: JBM*, 5(3).
- Lubeek SF, van Vugt LJ, Aben KK, van de Kerkhof PC, Gerritsen M-JP. The epidemiology and clinicopathological features of basal cell carcinoma in patients 80 years and older: A systematic review. *JAMA dermatol*. 2017;153(1):71-8
- M, Schnidar H, Eichberger T, Neill GW, Philpott MP, et al. 2004. Activation of the BCL2 promoter in response to Hedgehog/GLI signal transduction is predominantly mediated by GLI2. *Cancer Res.*; 64: 7724- 31
- Malicki J, Avidor-Reiss T. From the cytoplasm into the cilium: bon voyage. *Organogenesis*. 2014; 10:138–157.
- Maloverjan A, Piirsoo M, Michelson P, Kogerman P, Osterlund T. Identification of a novel serine/ threonine kinase ULK3 as a positive regulator of Hedgehog pathway. *Exp Cell Res*. 2010; 316:627–637.
- Marcellina P, Mappias A, Anwar AI, Ganda IJ, Hatta M, Masadah R. Expression of Patched-1 Protein in Aggressive and Nonaggressive Basal Cell Carcinoma. *Am J Clin Exp Med*, 2016. 4(5):122
- Marcet B, Chevalier B, Coraux C, Kodjabachian L, Barbry P. MicroRNA-based silencing of Delta/ Notch signaling promotes multiple cilia formation. *Cell Cycle*. 2011; 10:2858–2864.
- Maria Kristanti Sambuaga; Aida S.D. Suriadiredja. 2016. Association of the expression of

GLI1 transcription factor as the component of sonic hedgehog pathway with the aggressive characteristics of the growth pattern of basal cell carcinomas histological subtypes. Universitas Indonesia Library. Di download dari : <https://lontar.ui.ac.id/detail?id=20423493&lokasi=lokal>

Marks R, Staples M, Giles GG. Trends in non-melanocytic skin cancer treated in Australia: the second national survey. *Int J Cancer* 1993; 53:585.

Mawardi P, Kalim H, Kalim KH, Fitri LE, Mintaroem K, Mudigdo A, et al. Mid-face location of primary basal cell carcinoma related to cancer aggressivity. *Asian Pac JTrop Dis*. 2016;6(8):650-3

May-Simera HL, Kelley MW. Cilia, Wnt signaling, and the cytoskeleton. *Cilia*. 2012; 1:7.

Megan wetzel, John strickley. Tye Haeberle, timothy s. Brown. Depth of Invasion of Aggressive and Nonaggressive Basal Cell Carcinoma. *J Clin Aesthet Dermatol*. 2019;12(3):12–14.

Didownload:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6440705/pdf/jcad_12_3_12.pdf

Meng X, Poon R, Zhang X, et al. Suppressor of fused negatively regulates beta-catenin signaling. *J Biol Chem*. 2001; 276:40113–40119.

Messina J, Epstein EH. Basal cell carcinoma. In: Elder D, Massi D, Scolyer R, Willemze R, editors. *WHO classification of skin tumours*. 4th ed. Lyon: The International Agency for Research on Cancer (IARC); 2018. p. 26-34.

Milenkovic L, Scott MP, Rohatgi R. Lateral transport of Smoothed from the plasma membrane to the membrane of the cilium. *J Cell Biol*. 2009; 187:365–374.

Montagna E, Lopes OS. Molecular basis of basal cell carcinoma. Vol. 92, *Anais Brasileiros de Dermatologia*. 2017.p.517-20. 11.

Muzic JG, Schmitt AR, Wright AC, et al. Incidence and Trends of Basal Cell Carcinoma and Cutaneous Squamous Cell Carcinoma: A Population-Based Study in Olmsted County, Minnesota, 2000 to 2010. *Mayo Clin Proc* 2017; 92:890.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology: Basal cell skin cancer. Version 2.2022. https://www.nccn.org/professionals/physician_gls/pdf/nmsc.pdf (Accessed on August 13, 2021).

National Comprehensive Cancer Network (NCCN) Guidelines in Oncology: Basal Cell Skin Cancer. Version 2.2022. Available at: <https://www.nccn.org>.

Noubissi FK, Kim T, Kawahara TN, et al. Role of CRD-BP in the growth of human basal cell carcinoma cells. *J Invest Dermatol*. 2014; 134:1718–1724.

Oninla OA, Oninla SO, Ajani AA, Report C. Gender Dermatoses: The Role of Sex Hormones in Skin Diseases. 2017;2(2):1–4.

Otsuka, A., Levesque, M.P., Dummer, R. and Kabashima, K., 2015. Hedgehog signaling in basal cell carcinoma. *Journal of dermatological science*, 78(2), pp.95-100.

Pal K, Hwang SH, Somatilaka B, et al. Smoothed determines beta-arrestin-mediated removal of the G protein-coupled receptor Gpr161 from the primary cilium. *J Cell Biol*. 2016; 212:861–875.

Paroo Z, Ye X, Chen S, Liu Q. Phosphorylation of the human microRNA-generating complex mediates MAPK/Erk signaling. *Cell*. 2009; 139:112–122.

Pellegrini C, Maturo MG, Di Nardo L, et al. Understanding the Molecular Genetics of Basal Cell Carcinoma. *Int J Mol Sci* 2017; 18.

Peris K, Fargnoli MC, Garbe C, et al. Diagnosis and treatment of basal cell carcinoma: European consensus-based interdisciplinary guidelines. *Eur J Cancer* 2019; 118:10.

Piperno G, Mead K, LeDizet M, Moscatelli A. Mutations in the “dynein regulatory complex” alter the ATP-insensitive binding sites for inner arm dyneins in *Chlamydomonas*

- axonemes. *J Cell Biol.* 1994; 125:1109–1117.
- Priel, S., Cortelazzi, B., Dal Col, V., Marson, D., Laurini, E., Fermeglia, M., Licitra, L., Pilotti, S., Bossi, P. and Perrone, F., 2015. Smoothed (SMO) receptor mutations dictate resistance to vismodegib in basal cell carcinoma. *Molecular oncology*, 9(2), pp.389-397.
- Pusapati GV, Hughes CE, Dorn KV, et al. EFCAB7 and IQCE regulate hedgehog signaling by tethering the EVC-EVC2 complex to the base of primary cilia. *Dev Cell.* 2014; 28:483–496.
- Quazi, S. J., Aslam, N., Saleem, H., Rahman, J., & Khan, S., 2020. Surgical margin of excision in basal cell carcinoma: a systematic review of literature. *Cureus*, 12(7).
- Ramelyte, E., Nägeli, M.C., Hunger, R., Merat, R., Gaide, O., Navarini, A.A., Cozzio, A., Wagner, N.B., Maul, L.V. and Dummer, R., 2023. Swiss Recommendations for Cutaneous Basal Cell Carcinoma. *Dermatology*, 239(1), pp.122-131.
- Reifenberger J, Wolter M, Knobbe CB, et al. Somatic mutations in the PTCH, SMOH, SUFUH and TP53 genes in sporadic basal cell carcinomas. *Br J Dermatol* 2005; 152:43.
- Reifenberger, J. et al. (1998) ‘Mutational analysis of the SHH and PTCH genes in sporadic basal cell carcinomas’, *Journal of Dermatological Science*, 16. doi:10.1016/s0923-1811(98)83850-4.
- Reifenberger, J., Wolter, M., Knobbe, C.B., Köhler, B., Schönicke, A., Scharwächter, C., Kumar, K., Blaschke, B., Ruzicka, T. and Reifenberger, G., 2005. Somatic mutations in the PTCH, SMOH, SUFUH and TP53 genes in sporadic basal cell carcinomas. *British Journal of Dermatology*, 152(1), pp.43-51.
- Reizner GT, Chuang TY, Elpern DJ, et al. Basal cell carcinoma in Kauai, Hawaii: the highest documented incidence in the United States. *J Am Acad Dermatol* 1993; 29:184.
- Richtig, G., Aigelsreiter, A.M., Asslaber, M., Weiland, T., Pichler, M., Eberhard, K., Sygulla, S., Schauer, S., Hoefler, G. and Aigelsreiter, A., 2019. Hedgehog pathway proteins SMO and GLI expression as prognostic markers in head and neck squamous cell carcinoma. *Histopathology*, 75(1), pp.118-127.
- Riesye Arisanty, Muhammad Habiburrahman, Maria A.P Maharani. Clinicopathologic and Histomorphological Aspect of Basal Cell Carcinoma in Dr. Cipto Mangunkusumo Hospital: A Retrospective Analysis of Twenty Years Experience. *eJ KI Vol. 9, No. 2, Agustus 2021.* Didownload dari : <https://ejki.fk.ui.ac.id/index.php/journal/article/view/34/20>
- Rifani, L., Arisanty, R. and Handayani, S.I., 2019. Jalur Pensinyalan Hedgehog pada Karsinoma Sel Basal. *Pratista Patologi*, 6(2), pp.68-80.
- Riobo NA, Lu K, Ai X, Haines GM, Emerson CP Jr. Phosphoinositide 3-kinase and akt are essential for sonic hedgehog signaling. *Proc Natl Acad Sci USA.* 2006; 103:4505–4510.
- Rogers HW, Weinstock MA, Harris AR, et al. Incidence estimate of nonmelanoma skin cancer in the United States, 2006. *Arch Dermatol* 2010; 146:283.
- Rossi, M., Magnoni, L., Miracco, C., Mori, E., Tosi, P., Pirtoli, L., Tini, P., Oliveri, G., Cosci, E. and Bakker, A., 2011. β -catenin and Gli1 are prognostic markers in glioblastoma. *Cancer biology & therapy*, 11(8), pp.753-761.
- Rubin, A.I., Chen, E.H. and Ratner, D. (2005) ‘Basal-cell carcinoma’, *New England Journal of Medicine*, 353(21), pp. 2262–2269. doi:10.1056/nejmra044151.
- Sand M, Bechara FG, Sand D, et al. Long-noncoding RNAs in basal cell carcinoma. *Tumour Biol.* 2016; 37:10595–10608
- Sand M, Skrygan M, Georgas D, et al. Expression levels of the microRNA maturing microprocessor complex component DGCR8 and the RNA-induced silencing complex (RISC) components argonaute-1, argonaute-2, PACT, TARBP1, and TARBP2 in epithelial skin cancer. *Mol Carcinog.* 2012; 51:916–922.
- Sand M, Skrygan M, Sand D, et al. Expression of microRNAs in basal cell carcinoma. *Br J*

- Saydam O, Shen Y, Wurdinger T, et al. Downregulated microRNA-200a in meningiomas promotes tumor growth by reducing E-cadherin and activating the Wnt/ β -catenin signaling pathway. *Mol Cell Biol*. 2009; 29:5923–5940
- Scotto J, Fears TR, Fraumeni JF Jr, et al. Incidence of nonmelanoma skin cancer in the United States in collaboration with Fred Hutchinson Cancer Research Center. NIH publication No. 83-2433, U.S. Dept. of Health and Human Services, Public Health Service, National Institutes of Health, National Cancer Institute, Bethesda, MD 1983:xv. p.113.
- Sekulic, A. et al. (2012) ‘Efficacy and safety of Vismodegib in advanced basal-cell carcinoma’, *New England Journal of Medicine*, 366(23), pp. 2171–2179. doi:10.1056/nejmoa1113713.
- Shi S-R, Jiang G, Taylor CR. 2000. Antigen retrieval techniques: immunohistochemistry and molecular morphology. Eaton Publishing Company; p.23-6
- Skoda AM, Simovic D, Karin V, Kardum V, Vranic S, Serman L. The role of the Hedgehog signaling pathway in cancer : A comprehensive review. *Bosn J Basic Med Sci*. 2018; 18: 8-20.
- Sonkoly E, Loven J, Xu N, et al. MicroRNA-203 functions as a tumor suppressor in basal cell carcinoma. *Oncogenesis*. 2012; 1:e3.
- Stacey SN, Helgason H, Gudjonsson SA, et al. New basal cell carcinoma susceptibility loci. *Nat Commun*. 2015; 6:6825
- Stephens PM, Martin B, Ghafari G, Luong J, Nahar VK, Pham L, et al. Skin Cancer Knowledge, Attitudes, and Practices among Chinese Population: A Narrative Review. *Dermatol Res Pract*. 2018;2018:1-9
- Subramaniam P, Olsen CM, Thompson BS, Whiteman DC, Neale RE. Anatomical distributions of basal cell carcinoma and squamous cell carcinoma in a population-based study in Queensland, Australia. *JAMA Dermatology*. 2017;153:175–82
- Swanson NA, Grekin RC, Baker SR. Mohs surgery: Techniques, indications, and applications in head and neck surgery. *Head & neck surgery*. 1983;6(2):683-92
- Swartling FJ. Myc proteins in brain tumor development and maintenance. *Ups J Med Sci*. 2012; 117:122–131. [PubMed: 22348395] 33. Meyer N, Penn LZ. Reflecting on 25 years with MYC. *Nat Rev Cancer*. 2008; 8:976–990.
- Tan ST, Ghaznawie M, Reginata G. Deteksi dini karsinoma sel basal. *Indones J Cancer*. 2016;10(2):61-6.
- Tan ST, Reginata G. Diagnosis dan Tata Laksana Karsinoma Sel Basal. *CDK-235*, 2015. 42(12): 897-900. Verkouteren JAC, Ramdas KHR, Wakkee M, Nijsten T. Epidemiology of basal cell carcinoma: scholarly review. *Br J Dermatol* 2017; 177:359.
- Teng, Y., Yu, Y., Li, S., Huang, Y., Xu, D., Tao, X. and Fan, Y., 2021. Ultraviolet radiation and basal cell carcinoma: an environmental perspective. *Frontiers in Public Health*, 9, p.666528.
- Thomson J, Hogan S, Leonardi-Bee J, et al. Interventions for basal cell carcinoma: abridged Cochrane systematic review and GRADE assessments. *Br J Dermatol* 2021; 185:499.
- Tiyawatanaroj, A., Sudtikoonaseth, P. and Chayangsu, O., 2022. Basal cell carcinoma trends in Thailand: A 10-year retrospective study of demographic, clinical and histopathological features. *Dermatology Reports*, 14(1).
- Toha, S.S., Rahman, A., Mochtar, M., Julianto, I., Dharmawan, N., Mawardi, P., Wasita, B. and Setyawan, N.A., 2019. Kejadian Karsinoma Sel Basal di RSUD Dr. Moewardi Surakarta Berdasarkan Subtipe Histopatologi menurut Jenis Kelamin, Usia, Lokasi Anatomi, dan Diameter Tumor. *Cermin Dunia Kedokteran*, 46(4), pp.256-260.
- Tuson M, He M, Anderson KV. Protein kinase A acts at the basal body of the primary cilium to prevent Gli2 activation and ventralization of the mouse neural tube. *Development*.

- Valcourt, D.M., Dang, M.N., Wang, J. and Day, E.S., 2020. Nanoparticles for manipulation of the developmental Wnt, Hedgehog, and Notch signaling pathways in cancer. *Annals of biomedical engineering*, 48, pp.1864-1884.
- Varjosalo M, Bjorklund M, Cheng F, et al. Application of active and kinase-deficient kinome collection for identification of kinases regulating hedgehog signaling. *Cell*. 2008; 133:537–548.
- Vornicescu C, Şenilă SC, Bejinariu NI, et al. Predictive factors for the recurrence of surgically excised basal cell carcinomas: a retrospective clinical and immunopathological pilot study. *Exp Ther Med*. 2021;22(5):1–10. doi:10.3892/etm.2021.10771
- Walling HW, Fosko SW, Geraminejad PA, Whitaker DC, Arpey CJ. Aggressive basal cell carcinoma: Presentation, pathogenesis, and management. *Cancer Metastasis Rev*. 2004;23(3-4):389-402.
- Wallingford JB, Mitchell B. Strange as it may seem: the many links between Wnt signaling, planar cell polarity, and cilia. *Genes Dev*. 2011; 25:201–213
- Wang H, Xu Y, Shi J, et al. Photodynamic therapy in the treatment of basal cell carcinoma: a systematic review and meta-analysis. *Photodermatol Photoimmunol Photomed* 2015; 31:44.
- Wang, G.Y. et al. (2011) ‘Basal cell carcinomas arise from hair follicle stem cells in *ptch1*+/- mice’, *Cancer Cell*, 19(1), pp. 114–124. doi:10.1016/j.ccr.2010.11.007.
- Welsch MJ, Troiani BM, Hale L, DelTondo J, Helm KF, Clarke LE. Basal cell carcinoma characteristics as predictors of depth of invasion. *J Am Acad Dermatol*. 2012;67(1):47-53
- Wolf, D.J. (1987) ‘Surgical margins for basal cell carcinoma’, *Archives of Dermatology*, 123(3), p. 340. doi:10.1001/archderm.1987.01660270078019.
- Wong, S.Y. and Dlugosz, A.A. (2014) ‘Basal cell carcinoma, hedgehog signaling, and targeted therapeutics: The long and winding road’, *Journal of Investigative Dermatology*, 134. doi:10.1038/skinbio.2014.4.
- Wrone DA, Swetter SM, Egbert BM, et al. Increased proportion of aggressive-growth basal cell carcinoma in the Veterans Affairs population of Palo Alto, California. *J Am Acad Dermatol* 1996; 35:907.
- Wu F, Zhang Y, Sun B, McMahon AP, Wang Y. Hedgehog signaling: from basic biology to cancer therapy. *Cell Chem Biol*. 2017; 24: 252-80. 8.
- Wu S, Han J, Li WQ, et al. Basal-cell carcinoma incidence and associated risk factors in U.S. women and men. *Am J Epidemiol* 2013; 178:890.
- Xie, J., Murone, M., Luoh, S.M., Ryan, A., Gu, Q., Zhang, C., Bonifas, J.M., Lam, C.W., Hynes, M., Goddard, A. and Rosenthal, A., 1998. Activating Smoothed mutations in sporadic basal-cell carcinoma. *Nature*, 391(6662), pp.90-92.
- Xu, M., Li, X., Liu, T., Leng, A. and Zhang, G., 2012. Prognostic value of hedgehog signaling pathway in patients with colon cancer. *Medical Oncology*, 29, pp.1010-1016.
- Yahya YF, Pohan SS, Soetjipto S, Sudiana IK. Ekspresi β Catenin dan $\beta 4$ Integrin pada Karsinoma Sel Basal Agresif dan Non Agresif. *Jurnal Biosains Pascasarjana*, 2012. 14(1):1-11.
- Yang N, Li L, Eguether T, Sundberg JP, Pazour GJ, Chen J. Intraflagellar transport 27 is essential for hedgehog signaling but dispensable for ciliogenesis during hair follicle morphogenesis. *Development*. 2016; 142:2194–2202.
- Yang, Z., Cui, Y., Ni, W., Kim, S. and Xuan, Y., 2017. Gli1, a potential regulator of esophageal cancer stem cell, is identified as an independent adverse prognostic factor in esophageal squamous cell carcinoma. *Journal of cancer research and clinical oncology*, 143, pp.243-254.
- Yoo YA, Kang MH, Lee HJ, et al. Sonic hedgehog pathway promotes metastasis and

- lymphangiogenesis via activation of Akt, EMT, and MMP-9 pathway in gastric cancer. *Cancer Res.* 2011; 71:7061–7070.
- Youssef, K.K. et al. (2010) ‘Identification of the cell lineage at the origin of basal cell carcinoma’, *Nature Cell Biology*, 12(3), pp. 299–305. doi:10.1038/ncb2031.
- Zedan, W., Robinson, P.A., Markham, A.F. and High, A.S., 2001. Expression of the Sonic Hedgehog receptor ‘PATCHED’ in basal cell carcinomas and odontogenic keratocysts. *The Journal of Pathology: A Journal of the Pathological Society of Great Britain and Ireland*, 194(4), pp.473–477.
- Zhang H, Pasolli HA, Fuchs E. Yes-associated protein (YAP) transcriptional coactivator functions in balancing growth and differentiation in skin. *Proc Natl Acad Sci USA*. 2011; 108:2270–2275.
- Zhang L, Nadeem L, Connor K, Xu G. Mechanisms and Therapeutic Targets of microRNA-associated Chemoresistance in Epithelial Ovarian Cancer. *Curr Cancer Drug Targets*. 2016; 16:429–441.
- Zhao B, Wei X, Li W, et al. Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth control. *Genes Dev*. 2007; 21:2747–2761.
- Zhu H, Lo HW. 2010. The human glioma-associated oncogene homolog 1 (GLI1) family of transcription factors in gene regulation and diseases. *Current Genomics*.; 11(4): 238–45
- Zhu, W., You, Z., Li, T., Yu, C., Tao, G., Hu, M. and Chen, X., 2011. Correlation of hedgehog signal activation with chemoradiotherapy sensitivity and survival in esophageal squamous cell carcinomas. *Japanese journal of clinical oncology*, 41(3), pp.386–393.
- Zindy F, Kawauchi D, Lee Y, et al. Role of the miR-17 approximately 92 cluster family in cerebellar and medulloblastoma development. *Biol Open*. 2014; 3:597–605.