

DAFTAR PUSTAKA

- Ahmad, F.I., Wagiran, A., Abd Samad, A., Rahmat, Z. and Sarmidi, M.R. (2016). Improvement of efficient in vitro regeneration potential of mature callus induced from Malaysian upland rice seed (*Oryza sativa* cv. Panderas). *Saudi Journal of Biological Sciences*, 23(1): 69-77.
- Al-Ashoush, A.M., Shibli, R.A., Tahtamouni, R.W., Al-Qudah, T.S. and Abu-Irarmaileh, B.B. (2020). Enhancement of Pentacyclic Triterpenoids (Betulinic and Oleanolic acids) production from callus cultures of *Lantana camara* L. *Advances in Horticultural Science*, 34(3): 255-263.
- Ali, A., Nongpiur, R. C., & Chrungoo, N. K. (2023). An efficient *Agrobacterium*-mediated transformation and regeneration protocol for aromatic black rice, *Oryza sativa* L.(cv. Chakhao Poireiton and Chakhao Amubi). *Plant Cell, Tissue and Organ Culture (PCTOC)*, 152(3), 619-633.
- Arifa, A.H., Syamsir, E. & Budijanto, S. (2021). Karakterisasi Fisikokimia Beras Hitam (*Oryza sativa* L.) dari Jawa Barat, Indonesia. *Agritech*, 41(1): 15-24.
- Asano, T., Kusano, H., Okuda, T., Kubo, N., Shimada, H., & Kadowaki, K. I. (2002). Rpp16 and Rpp17, from a common origin, have different protein characteristics but both genes are predominantly expressed in rice phloem tissues. *Plant and cell physiology*, 43(6), 668-674.
- Asbel, L. E., & Levison, M. E. (2000). Cephalosporins, carbapenems, and monobactams. *Infectious Disease Clinics*, 14(2), 435-447.
- Azevedo, F., Pereira, H., & Johansson, B. (2017). Colony PCR. In *PCR* (pp. 129-139). Springer, New York, NY.
- Babaoglu, M., Davey, M. R., & Power, J. B. (2000). Genetic engineering of grain legumes: key transformation events. *AgBiotechNet*, 2(ABN 050).
- Backer, C.A. & R.C. B. van den Brink. (1968). *Flora of Java Vol III*. WoltersNoordhoff N. Groningen. Netherlands.
- Bergkessel, M., & Guthrie, C. (2013). Colony PCR. In *Methods in enzymology* (Vol. 529, pp. 299-309). Academic Press.
- Bhojwani, S. S., & Dantu, P. K. (2013). Plant tissue culture: an introductory text (Vol. 318). India: Springer.
- Breed, R. S., Murray, E. G. D., & Smith, N. R. (1957). Bergey's manual of determinative bacteriology. *Bergey's Manual of Determinative Bacteriology*, (7th Edition). The Williams & Wilkins Company, Baltimore.

- Callcott, T. E., Santhakumar, B. A., Luo, J., & Blanchard, L. C. (2018). Therapeutic potential of rice-derived polyphenols on obesity-related oxidative stress and inflammation. *Journal of Applied Biomedicine*, 16(4): 255-262.
- Cappucino, J.G., dan Sherman, N. (1998). *Microbiology: Laboratory Manuals 5th edition*. Benjamin/Cummings Science Publishing, California.
- Chanchula, N., Jaruwattanaphan, T. and Jala, A. (2014). Differential effects of sucrose and plant growth regulator on shoot multiplication and bulbil formation in *Oxalis Versicolour* in vitro. *ITJEMAST*, 5(4): 227-234.
- Chen, P.N., Kuo, W.H., Chiang, C.L., Chiou, H.L., Hsieh, Y.S. & Chu, S.C. (2006). Black rice anthocyanins inhibit cancer cells invasion via repressions of MMPs and u-PA expression. *Chemico-biological interactions*, 163(3): 218-229.
- Clough, S. J., & Bent, A. F. (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *The plant journal*, 16(6), 735-743.
- Cordeiro, D., Alves, A., Ferraz, R., Casimiro, B., Canhoto, J., & Correia, S. (2023). An Efficient *Agrobacterium*-Mediated Genetic Transformation Method for *Solanum betaceum* Cav. Embryogenic Callus. *Plants*, 12(5), 1202.
- Craggs, T. D. (2009). Green fluorescent protein: structure, folding and chromophore maturation. *Chemical Society Reviews*, 38(10), 2865-2875.
- Dindar, B., & Seven, O. (2009). Photodegradation of 2, 4-Dichlorophenoxy acetic acid on TiO₂, ZnO and Fe₂O₃ by sunlight. *Asian Journal of Chemistry*, 21(3), 2270.
- Doi, K., Izawa, T., Fuse, T., Yamanouchi, U., Kubo, T., Shimatani, Z., Yano, M., & Yoshimura, A. (2004). *Ehd1*, a B-type response regulator in rice, confers short-day promotion of flowering and controls FT-like gene expression independently of *Hd1*. *Genes & development*, 18(8), 926-936.
- Dreger, M., Mól, R., Deja, A., Raj, E., Mańkowska, G. and Wielgus, K. (2019). Improved plant regeneration in callus cultures of *Sorghum bicolor* (L.) Moench. *In Vitro Cellular & Developmental Biology-Plant*, 55(2): 190-198.
- Duyi, S., Baran, A., & Chandra, D. S. (2017). Pigmented rice a potential source of bioactive compounds: a review. *International Journal of Food Science & Technology*, 52(5): 1073-1081.
- Dwiyani, R. (2015). *Kultur Jaringan Tanaman*. Pelawa Sari Percetakan & Penerbit. Denpasar.
- Eibl, R., Eibl, D., Pörtner, R., Catapano, G. & Czermak, P. (2008). *Cell and tissue reaction engineering*. Springer Science & Business Media.

- Fatima, Z., Mujib, A., Fatima, S., Arshi, A. and Umar, S. (2009). Callus induction, biomass growth, and plant regeneration in *Digitalis lanata* Ehrh.: influence of plant growth regulators and carbohydrates. *Turkish Journal of Botany*, 33(6): 393-405.
- George, E.F., Hall, M.A., De Klerk, G. (2008). *Plant Propagation by Tissue Culture 3rd Edition Vol. I*. Springer, Netherlands.
- Grewal, D., Gill, R., & Gosal, S. S. (2006). Influence of antibiotic cefotaxime on somatic embryogenesis and plant regeneration in indica rice. *Biotechnology Journal: Healthcare Nutrition Technology*, 1(10), 1158-1162.
- Grossmann, K. (2003). Mediation of herbicide effects by hormone interactions. *J Plant Growth Regul*, 22(1): 109-122.
- Heikrujam, J., Kishor, R., & Behari Mazumder, P. (2020). The Chemistry Behind Plant DNA Isolation Protocols. IntechOpen. doi: 10.5772/intechopen.92206
- Hendawey, M.H. and Abo El Fadl, R.E. (2014). Biochemical studies on the production of active constituents in *Stevia rebaudiana* L. callus. *Glob J Biotechnol Biochem*, 9: 76-93.
- Hooykaas, P.J.J. (1989) Transformation of plant cells via *Agrobacterium*. *Plant Molecular Biology*, 13, 327-336.
- Hoque, K.M.A., Azdi, Z.A. and Prodhan, S.H. (2013). Development of callus initiation and regeneration system of different indigenous indica rice varieties. *J. Biol*, 1: 46-51.
- Hwang, H. H., Yu, M., & Lai, E. M. (2017). *Agrobacterium*-mediated plant transformation: biology and applications. *The Arabidopsis Book*, 15.
- Ikeuchi, M., Sugimoto, K. & Iwase, A. (2013). Plant callus: mechanisms of induction and repression. *The plant cell*, 25(9): 3159-3173.
- Islam, M. M., Ahmed, M., & Mahaldar, D. (2005). In vitro callus induction and plant regeneration in seed explants of rice (*Oryza sativa* L.). *Research Journal of Agriculture and Biological Sciences*, 1(1), 72-75.
- Izawa, T., Oikawa, T., Sugiyama, N., Tanisaka, T., Yano, M., & Shimamoto, K. (2002). Phytochrome mediates the external light signal to repress FT orthologs in photoperiodic flowering of rice. *Genes & development*, 16(15), 2006-2020.
- Jan, R., Khan, M. A., Asaf, S., Lee, I. J., & Kim, K. M. (2020). Modulation of sugar and nitrogen in callus induction media alter PAL pathway, SA and biomass accumulation in rice callus. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 143(3), 517-530.

- Jeng, T. L., Lai, C. C., Ho, P. T., Shih, Y. J., & Sung, J. M. (2012). Agronomic, molecular and antioxidative characterization of red- and purple-pericarp rice (*Oryza sativa* L.) mutants in Taiwan. *Journal of Cereal Science*, 56(2): 425-431.
- Jiang, F., Feng, Z., Liu, H., & Zhu, J. (2015). Involvement of plant stem cells or stem cell-like cells in dedifferentiation. *Frontiers in Plant Science*, 6: 1028.
- Jiménez, V. M., Guevara, E., Herrera, J., & Bangerth, F. (2001). Endogenous hormone levels in habituated nucellar Citrus callus during the initial stages of regeneration. *Plant Cell Reports*, 20, 92-100.
- Jo, E.A., Tewari, R.K., Hahn, E.J., Paek, K.Y. (2008). In vitro sucrose concentration affects growth and acclimatization of *Alocasia amazonica* plantlets. *Plant Cell, Tissue Organ Cult*, 96(3):307.
- Khan, T., Abbasi, B. H., Zeb, A., & Ali, G. S. (2018). Carbohydrate-induced biomass accumulation and elicitation of secondary metabolites in callus cultures of *Fagonia indica*. *Industrial Crops and Products*, 126, 168-176.
- Khatoon, H., Anokhe, A., dan Kalia, V. (2022). Catalase Test: A Biochemical Protocol for Bacterial identification. *AgriCos e-Newsletter*, 03(01): 53-55.
- Kojima, S., Takahashi, Y., Kobayashi, Y., Monna, L., Sasaki, T., Araki, T., & Yano, M. (2002). *Hd3a*, a rice ortholog of the *Arabidopsis* FT gene, promotes transition to flowering downstream of Hd1 under short-day conditions. *Plant and cell physiology*, 43(10), 1096-1105.
- Komiya, R., Ikegami, A., Tamaki, S., Yokoi, S., & Shimamoto, K. (2008). *Hd3a* and *RFT1* are essential for flowering in rice. *Development*, 135(4): 767-774.
- Krikorian, A. D., & Smith, D. L. (1992). Somatic embryogenesis in carrot (*Daucus carota*). In *Plant Tissue Culture Manual: Supplement 1* (pp. 19-50). Boston, MA: Springer US.
- Kristantini, T., Basunanda, P., Murti, R.H., & Widyayanti, S. (2012). Morphological of genetic relationships among black rice landraces from Yogyakarta and surrounding areas. *ARPJ Agric Biol Sci*, 7:12–18
- Kristantini, S.W., Sutarno, S. & Sudarmaji, S. (2015). Keragaman genetik lima kultvar lokal padi beras hitam asal Yogyakarta berdasarkan sifat morfologi. *Prosiding Seminar Nasional Sumber Daya*, 90: 90-100.
- Kurepa, J., Shull, T. E., & Smalle, J. A. (2019). Antagonistic activity of auxin and cytokinin in shoot and root organs. *Plant Direct*, 3(2), e00121.
- Kurniasih, N.S., Susandarini, R., Susanto, F.A., Nuringtyas, T.R., Jenkins, G. & Purwestri, Y.A. (2019). Characterization of Indonesian pigmented rice

(*Oryza sativa*) based on morphology and Single Nucleotide Polymorphisms. *Biodiversitas Journal of Biological Diversity*, 20(4): 1208-1214.

Kushwaha, U. K. S. (2016). *Black Rice: Research, History and Development*. Springer International Publishing, Switzerland.

Lee, K., Jeon, H., & Kim, M. (2002). Optimization of a mature embryo-based in vitro culture system for high-frequency somatic embryogenic callus induction and plant regeneration from japonica rice cultivars. *Plant Cell, Tissue and Organ Culture*, 71, 237-244.

Legha, M. R., Prasad, K. V., Singh, S. K., Kaur, C., Arora, A., & Kumar, S. (2012). Induction of carotenoid pigments in callus cultures of *Calendula officinalis* L. in response to nitrogen and sucrose levels. *In Vitro Cellular & Developmental Biology-Plant*, 48(1), 99-106.

Li, M., Li, H., Hu, X., Pan, X., & Wu, G. (2011). Genetic transformation and overexpression of a rice *Hd3a* induces early flowering in *Saussurea involucreata* Kar. et Kir. ex Maxim. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 106(3), 363-371.

Li, F., Cui, X., Feng, Z., Du, X., & Zhu, J. (2012). The effect of 2, 4-D and kinetin on dedifferentiation of petiole cells in *Arabidopsis thaliana*. *Biologia plantarum*, 56(1), 121-125.

Low, L., Yang, S., Kok, D. A. , Ong- Abdullah, J., Tan, N., & Lai, K. (2018). Transgenic Plants: Gene Constructs, Vector and Transformation Method. In (Ed.), *New Visions in Plant Science*. IntechOpen. Pp 41-61.

Maneewan, K., Bunnag, S., Theerakulpisut, P., Kosittrakun, M., & Suwanagul, A. (2005). Transformation of rice (*Oryza sativa* L.) cv. Chainat 1 using chitinase gene. *Transformation*, 27(6), 1152.

Manimaran, P., Kumar, G. R., Reddy, M. R., Jain, S., Rao, T. B., Mangrauthia, S. K., Sundaram, R.M., Ravichandran, S., & Balachandran, S. M. (2013). Infection of early and young callus tissues of indica rice BPT 5204 enhances regeneration and transformation efficiency. *Rice Science*, 20(6), 415-426.

Mau, J.L., Lee, C.C., Chen, Y.P. & Lin, S.D. (2017). Physicochemical, antioxidant and sensory characteristics of chiffon cake prepared with black rice as replacement for wheat flour. *LWT*, 75: 434-439.

Mehrotra, S. & Goyal, V. (2016). *Agrobacterium-Mediated Alien Gene Transfer Biofabricates Designer Plants*. In Watson, R. R., Preedy, V.R. (Eds.), *Genetically Modified Organisms in Food* (pp 63-73). Academic Press.

Morton, E. R., & Fuqua, C. (2012). Laboratory maintenance of *Agrobacterium*. *Current protocols in microbiology*, 24(1), 3D-1.

- Mostafiz, S.B., & Wagiran, A. (2018). Efficient callus induction and regeneration in selected indica rice. *Agronomy*, 8(5), 77.
- Nauerby, B., Billing, K., & Wyndaele, R. (1997). Influence of the antibiotic timentin on plant regeneration compared to carbenicillin and cefotaxime in concentrations suitable for elimination of *Agrobacterium tumefaciens*. *Plant Science*, 123(1-2), 169-177.
- Navarro, C., Abelenda, J. A., Cruz-Oró, E., Cuéllar, C. A., Tamaki, S., Silva, J., Shimamoto, K., & Prat, S. (2011). Control of flowering and storage organ formation in potato by FLOWERING LOCUS T. *Nature*, 478(7367), 119-122.
- Newell, C. A. (2000). Plant transformation technology. *Molecular biotechnology*, 16(1), 53-65.
- Nguyen, D. Q., Van Eck, J., Eamens, A. L., & Grof, C. P. (2020). Robust and reproducible *Agrobacterium*-mediated transformation system of the C4 genetic model species *Setaria viridis*. *Frontiers in Plant Science*, 11, 513095.
- Nishimura, A., Ashikari, M., Lin, S., Takashi, T., Angeles, E. R., Yamamoto, T., & Matsuoka, M. (2005). Isolation of a rice regeneration quantitative trait loci gene and its application to transformation systems. *Proceedings of the National Academy of Sciences*, 102(33), 11940-11944.
- Nishimura, A., Aichi, I., & Matsuoka, M. (2006). A protocol for *Agrobacterium*-mediated transformation in rice. *Nature protocols*, 1(6), 2796-2802.
- Nissen, S. J., & Sutter, E. G. (1990). Stability of IAA and IBA in nutrient medium to several tissue culture procedures. *HortScience*, 25(7), 800-802.
- Nolan, K. E., & Rose, R. J. (2010). Plant regeneration-somatic embryogenesis. *Plant cell culture*, 39-60.
- Özkul, M., Özel, Ç. A., Yüzbaşıoğlu, D., & Ünal, F. (2016). Does 2, 4-dichlorophenoxyacetic acid (2,4-d) induce genotoxic effects in tissue cultured *Allium* roots?. *Cytotechnology*, 68(6), 2395-2405.
- Pandey, S. K., Ramesh, B., & Gupta, P. K. (1994). Study on effect of genotype and culture medium on callus formation and plant regeneration in rice (*Oryza sativa* L.). *Indian Journal of Genetics and Plant Breeding*, 54(03), 293-299.
- Parizot, B., Laplaze, L., Ricaud, L., Boucheron-Dubuisson, E., Bayle, V., Bonke, M., De Smet, I., Poethig, S.R., Helariutta, Y., Haseloff, J. & Chriqui, D. (2008). Diarch symmetry of the vascular bundle in *Arabidopsis* root encompasses the pericycle and is reflected in distich lateral root initiation. *Plant Physiology*, 146(1): 140-148.
- Park, S. (2021). *Plant tissue culture: techniques and experiments*. Academic Press.

- Pasternak, T. P., & Steinmacher, D. (2024). Plant Growth Regulation in Cell and Tissue Culture In Vitro. *Plants*, 13(2), 327.
- Pratiwi, R., & Purwestri, Y. A. (2017). Black rice as a functional food in Indonesia. *Functional Foods in Health and Disease*, 7(3), 182-194.
- Purugganan, M.D., & Fuller, D.Q. (2009). The nature of selection during plant domestication. *Nature*, 457(7231): 843-848.
- Purwestri, Y.A., Ogaki, Y., Tamaki, S., Tsuji, H., & Shimamoto, K. (2009). The 14-3-3 protein GF14c acts as a negative regulator of flowering in rice by interacting with the florigen *Hd3a*. *Plant Cell Physiol*, 50:429-438.
- Purwestri, Y. A., Sari, R. D. K., Anggraeni, L. N., & Sasongko, A. B. (2015). *Agrobacterium tumefaciens* mediated transformation of rolC:: *Hd3a*-GFP in black rice (*Oryza sativa* L. cv. Cempo Ireng) to promote early flowering. *Procedia Chemistry*, 14, 469-473.
- Purwestri, Y.A., Susanto, F.A., and Tsuji, H. (2017). *Hd3a* Florigen Recruits Different Proteins to Reveal Its Function in Plant Growth and Development. In Sniežana J. (Ed). *Plant Engineering*. IntechOpen. <https://doi.org/10.5772/intechopen.70263>.
- Rahman, Z. A., Seman, Z. A., Basirun, N., Julkifle, A. L., Zainal, Z., & Subramaniam, S. (2011). Preliminary investigations of *Agrobacterium*-mediated transformation in indica rice MR219 embryogenic callus using *gusA* gene. *African Journal of Biotechnology*, 10(40), 7805-7813.
- Rahman, Z. A., RamLi, A., Kamaruzaman, R., Seman, Z. A., Othman, A. N., Zainal, Z., Uddain, J. & Subramaniam, S. (2015). Efficient plant regeneration of malaysian aromatic rice (*Oryza sativa* L.) via improved somatic embryogenesis pathway. *Emirates journal of Food and Agriculture*, 857-863.
- Raja, N. I., Bano, A., Rashid, H., Khan, M. H., & Chaudhry, Z. (2009). Effect of age of embryogenic callus on plant regeneration in local cultivars of wheat (*Triticum aestivum* L.). *Pak. J. Bot*, 41(6), 2801-2806.
- Rao, M. V. R., & Rao, G. J. N. (2007). *Agrobacterium*-mediated transformation of indica rice under Acetosyringone-free conditions. *Plant Biotechnology*, 24(5), 507-511.
- Ratanasut, K., Rod-In, W., & Sujipuli, K. (2017). In planta *Agrobacterium*-mediated transformation of rice. *Rice Science*, 24(3), 181-186.
- Ratnayake, R. M. L. K., & Hettiarachchi, G. H. C. M. (2010). Development of an efficient *Agrobacterium* mediated transformation protocol for Sri Lankan rice variety-Bg 250.
- Reiner, K. (2012). Carbohydrate fermentation protocol. *Energy*, 11, 12.

- Rojo, F. P., Seth, S., Erskine, W., & Kaur, P. (2021). An improved protocol for *Agrobacterium*-mediated transformation in subterranean Clover (*Trifolium subterraneum* L.). *International Journal of Molecular Sciences*, 22(8), 4181.
- Roly, Z.Y., Islam, M.M., Shaekh, M.P.E., Arman, M.S.I., Shahik, S.M., Das, D., Haamem, M.M.E. and Khalekuzzaman, M., (2014). In vitro callus induction and regeneration potentiality of aromatic rice (*Oryza sativa* L.) cultivars in differential growth regulators. *International Journal of Applied Sciences and Biotechnology*, 2(2): 160-167.
- Rueb, S., Leneman, M., Schilperoort, R. A., & Hensgens, L. A. M. (1994). Efficient plant regeneration through somatic embryogenesis from callus induced on mature rice embryos (*Oryza sativa* L.). *Plant Cell, Tissue and Organ Culture*, 36, 259-264.
- Sangma, H.C.R. & Parameshwari, S. (2021). Health benefits of black rice (*Zizania aquatica*) - a review. *Materials Today: Proceedings*.
- Sari, Y.P., Kusumawati, E., Saleh, C., Kustiawan, W. and Sukartingsih, S. (2018). Effect of sucrose and plant growth regulators on callogenesis and preliminary secondary metabolic of different explant *Myrmecodia tuberosa*. *Nusantara Bioscience*, 10(3): 183-192.
- Shimomura, O., Johnson, F. H., & Saiga, Y. (1962). Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusan, *Aequorea*. *Journal of cellular and comparative physiology*, 59(3), 223-239.
- Smith, R.H. (2012). *Plant tissue culture: techniques and experiments*. Academic Press.
- Solanki, A. and Gupta, D. (2013). Effect of Antioxidants and Carbon source on Callus Culture of *Gymnema sylvestre* R. Br. *Research Journal of Chemistry and Environment*, 1(5): 121-125
- Solfanelli, C., Poggi, A., Loreti, E., Alpi, A., & Perata, P. (2006). Sucrose-specific induction of the anthocyanin biosynthetic pathway in *Arabidopsis*. *Plant physiology*, 140(2), 637-646.
- Song, Y., Bai, X., Dong, S., Yang, Y., Dong, H., Wang, N., ... & Li, S. (2020). Stable and efficient *agrobacterium*-mediated genetic transformation of larch using embryogenic callus. *Frontiers in Plant Science*, 11, 584492.
- Sugimoto, K., Jiao, Y. & Meyerowitz, E.M. (2010). *Arabidopsis* regeneration from multiple tissues occurs via a root development pathway. *Developmental cell*, 18(3): 463-471.
- Tamaki, S., Matsuo, S., Wong, H. L., Yokoi, S., & Shimamoto, K. (2007). *Hd3a* protein is a mobile flowering signal in rice. *Science*, 316(5827), 1033-1036.

- Taoka, K-i., Ohki, I., Tsuji, H., Furuita, K., Hayashi, K., Yanase, T., Yamaguchi, M., Nakashima, C., Purwestri, Y.A., Tamaki, S., Ogaki, Y., Shimada, C., Nakagawa, A., Kojima, C., & Shimamoto, K. (2011). 14-3-3 proteins act as intracellular receptors for rice *Hd3a* florigen. *Nature*, 476:332–335.
- Tariq, M., Ali, G., Hadi, F., Ahmad, S., Ali, N., & Shah, A. A. (2008). Callus Induction and in vitro Plant Regeneration of Rice (*Oryza sativa* L.). *Pakistan Journal of Biological Sciences*, 11(2), 255-259.
- Thorpe, T.A. (2007). History of plant tissue culture. *Molecular biotechnology*, 37(2): 169-180.
- Toki, S., Hara, N., Ono, K., Onodera, H., Tagiri, A., Oka, S., & Tanaka, H. (2006). Early infection of scutellum tissue with *Agrobacterium* allows high-speed transformation of rice. *The Plant Journal*, 47(6), 969-976.
- Tsuji, H., Tachibana, C., Tamaki, S., Taoka, K. I., Kyoizuka, J., & Shimamoto, K. (2015). *Hd3a* promotes lateral branching in rice. *The Plant Journal*, 82(2), 256-266.
- Tsuji, H., Taoka, K. I., & Shimamoto, K. (2011). Regulation of flowering in rice: two florigen genes, a complex gene network, and natural variation. *Current opinion in plant biology*, 14(1), 45-52.
- Upadhyaya, G., Sen, M., & Roy, A. (2015). In vitro callus induction and plant regeneration of rice (*Oryza sativa* L.) var. ‘Sita’, ‘Rupali’ and ‘Swarna Masuri’. *Asian Journal of Plant Science and Research*, 5(5), 24-27.
- Valvekens, D., Montagu, M.V. & Lijsebettens, M.V. (1988). *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis thaliana* root explants by using kanamycin selection. *Proceedings of the National Academy of Sciences*, 85(15): 5536-5540.
- Wahyuni, D.K., Huda, A., Faizah, S., Purnobasuki, H. and Wardoyo, B.P.E. (2020). Effects of light, sucrose concentration and repetitive subculture on callus growth and medicinally important production in *Justicia gendarussa* Burm. f. *Biotechnology Reports*, 27: e00473.
- Wanichananan, P., Teerakathiti, T., Roytrakul, S., Kirdmanee, C., & Peyachoknagul, S. (2010). A highly efficient method for *Agrobacterium* mediated transformation in elite rice varieties (*Oryza sativa* L. spp. indica). *African Journal of Biotechnology*, 9(34).
- Wang, X.D., Nolan, K.E., Irwanto, R.R., Sheahan, M.B. & Rose, R.J. (2011). Ontogeny of embryogenic callus in *Medicago truncatula*: the fate of the pluripotent and totipotent stem cells. *Annals of botany*, 107(4): 599-609.
- Wang, Y. & Li, J., (2005). The plant architecture of rice (*Oryza sativa*). *Plant molecular biology*, 59(1): 75-84.

- Weiss, D. (2000). Regulation of flower pigmentation and growth: multiple signaling pathways control anthocyanin synthesis in expanding petals. *Physiol Plant*, 110(2):152–157.
- Welsh, S., & Kay, S. A. (1997). Reporter gene expression for monitoring gene transfer. *Current opinion in biotechnology*, 8(5), 617-622.
- White, P. R. (1939). Potentially unlimited growth of excised plant callus in an artificial nutrient. *American Journal of Botany*, 26: 59–64.
- Wongsa, P. 2020. Phenolic Compounds and Potential Health Benefits of Pigmented Rice. In Ansari, M.R. (Ed.), *Recent Advances in Rice Research*. IntechOpen. <https://doi.org/10.5772/intechopen.93876>.
- Xia, M., Ling, W.H., Ma, J., Kitts, D.D. & Zawistowski, J. (2003). Supplementation of diets with the black rice pigment fraction attenuates atherosclerotic plaque formation in apolipoprotein E deficient mice. *The Journal of nutrition*, 133(3): 744-751.
- Yano, M., Katayose, Y., Ashikari, M., Yamanouchi, U., Monna, L., Fuse, T., Baba, T., Yamamoto, K., Umehara, Y., Nagamura Y. & Sasaki, T. (2000). *Hd1*, a major photoperiod sensitivity quantitative trait locus in rice, is closely related to the *Arabidopsis* flowering time gene *CONSTANS*. *The Plant Cell*, 12(12), 2473-2483.
- Yaseen, M., Ahmad, T., Sablok, G., Standardi, A. and Hafiz, I.A. (2013). Role of carbon sources for in vitro plant growth and development. *Molecular biology reports*, 40(4): 2837-2849.
- Yildiz, M., Onde, S. and Ozgen, M. (2007). Sucrose effects on phenolic concentration and plant regeneration from sugarbeet leaf and petiole explants. *Journal of Sugar Beet Research*, 44(1): 1-15.
- Yoshida, S. (1981). *Fundamentals of Rice Crop Science*. The International Rice Research Institute: Manila, Philippines.
- Young, J. M., Kerr, A., & Sawada, H. (2005). *Agrobacterium*. *Bergey's Manual of Systematics of Archaea and Bacteria*, 1-15.
- Zhang, M.W., Guo, B.J., Zhang, R.F., Chi, J.W., Wei, Z.C., Xu, Z.H., Zhang, Y. & Tang, X.J. (2006). Separation, purification and identification of antioxidant compositions in black rice. *Agricultural Sciences in China*, 5(6): 431-440.
- Zhang, X., Shen, Y., Prinyawiwatkul, W., King, J.M. & Xu, Z. (2013). Comparison of the activities of hydrophilic anthocyanins and lipophilic tocopherols in black rice bran against lipid oxidation. *Food chemistry*, 141(1): 111-116.
- Zimmer, M. (2002). Green fluorescent protein (GFP): applications, structure, and related photophysical behavior. *Chemical reviews*, 102(3), 759-782.