



DAFTAR PUSTAKA

- Abadi, Fitra, Q.A., Nurdiani, R., Firdaus, M. 2021. Aktivitas Antioksidan dan Antibakteri Hidrolisat Protein dari Hasil Samping Ikan Kakatua (*Chlorurus sordidus*). Universitas Brawijaya.
- Abraha, B., M. Abdu, S. Melake, Y. Weyni, K. Selamwit, dan H. Weldemariam. 2017. Production of fish protein hydrolysate from silver catfish (*Arius thalassinus*). MOJ Food Processing & Technology 5(4): 328 – 335.
- Aitken, I. M Mackie, J. H Merrit and M.L Windsor. (1982). Fish Handling and Processing. Ministry of Agriculture, Fisheries, and Food, Torry Research Station Press. Edinburgh.
- Aryani, F., Yulianti, dan T. Martati. 2004. Studi perubahan kadar histamin pada pindang tongkol (*Euthynnus affinis*) selama penyimpanan. Jurnal Penelitian Perikanan Indonesia 10(3): 35-46.
- Baehaki A., D.L. Shanti, dan R.R. Achmad. 2015. Hidrolisis protein ikan patin menggunakan enzim papain dan aktivitas antioksidan hidrolisatnya. Jurnal Pengolahan Hasil Perikanan Indonesia 18(3): 230-239.
- Baharuddin, N. A., N. R. A. Halim, dan N. M. Sarbon. 2016. Effect of degree of hydrolysis (DH) on the functional properties and angiotensin i-converting enzyme (ACE) inhibitory activity of eel (*Monopterus albus*) protein hydrolysate. International Food Research Journal 23(4): 1424-1431.
- Bhaskar, N., Mahendrakar, N.S. 2008. Protein hydrolysate from visceral waste protein of catla (*Catla catla*): optimization of hydrolysis condition for a commercial neutral protease. Journal Bioresource Technology 99: 4105-4111.
- Charoenphun, N., Benjamas, C., Nualpun, S., and Wirote. 2013. Calcium- binding peptides derived from tilapia (*Oreochromis niloticus*) protein hydrolysate. European Food Research and Technology, 236(1): 57-63.
- Chen, et al. (2012). Graphene Oxide: Preparation, Functionalization, and Electrochemical Application. Chemical Review, 112, 6027-6053.
- Dali, S., Natsir, H. Usman, H. dan Ahmad, A. 2011. Bioaktivitas Antibakteri Fraksi protein Alga Merah *Gelidium amansii* dari Perairan Cikoang Kabupaten Takalar, Sulawesi Selatan. 15 (1):47–52.
- Foh, M.B.K., Tamara, M.T., Amadou, I., Foh, B.M., Wenshui, X. 2011. Chemical and physicochemical properties of tilapia (*Oreochromis niloticus*) fish protein hydrolysate and concentrate. Journal International Biological Chemistry 10: 1-15.
- Haryati, S. dan Aris Munandar. 2012. Peenerapan konsep *zero waste* pada pengolahan abon ikan bandeng (*Chanos chanos*). Jurnal Perikanan dan Kelautan. 2(2): 127-130.



Hasnaliza, H., Maskat, M.Y., Wan, A.W.M., Mamot, S. 2010. The effect of enzyme concentration, temperature and incubation time on nitrogen content and degree of hydrolysis of protein precipitate from cockle (*Anadara granosa*) meat wash water. International Food Research Journal 17:147-152.

Hudzicki, J. (2009). Kirby-Bauer Disk Diffusion Susceptibility Test Protocol. Retrieved 15 Mei, 2014.

Intarasirisawat, R., S. Benjakul, W. Visessanguan, dan J. Wu. 2012. Antioxidative and functional properties of protein hydrolysate from defatted skipjack (*Katsuwonus pelamis*) roe. Journal Food Chemistry 135(4): 3039-3048.

Jahwetz, E., Melnick, J.L., Adelberg, E.A. 2005. Mikrobiologi Kedokteran, diterjemahkan oleh Mudihardi, E., Kuntaman, Wasito, E.B., Mertaniasih, N.M., Harsono, S., Alimsardjono, L., Edisi XXII, 327-335, 336-363. Penerbit Salemba Medika, Jakarta.

Jemil, I., M. Jridi, R. Nasri, N. Ktari, R.B.S Salem, M. Mehiri, M. Hajji, M. Nasri. 2014. Functional, antioxidant and antibacterial properties of protein hydrolysates prepared from fish meat fermented by *Bacillus subtilis* A26. Process Biochemistry 49(6): 963-972.

Kerr M., P. Lawicki, S. Aguirre, dan C. Rayner 2002. Effect of Storage Conditions on Histamine Formation in Fresh and Canned Tuna. Victorian Government Departement of Human Services, Werribee. Research Report.

Kim, S.K., Y.R. Choi, P.J. Park, J.H. Choi, dan S.H. Moon. 1999. Screening of biofunctional peptides from cod processing wastes. Journal of Korean Society of Agricultural Chemistry and Biotechnology 43: 225-227.

Koesoemawardani, D., Nurainy, F., Hidayati, S. 2011. Proses pembuatan hidrolisat protein ikan rucah. Jurnal Natur Indonesia Vol 13(3): 256-261.

Kung H.F, Wang T.Y, Huang Y.R, Lin C.S, Wu S.W, Lin C.M, Tsai Y.H. 2009. Isolation and identification of histamine-forming bacteria in tuna sandwiches. Journal Food Control 20(3):1013-1017.

Kristiansson, H.G., dan B.A. Rasco. 2000. Biochemical and functional properties of atlantic salmon (*Salmo salar*) muscle proteins hydrolyzed with various alkaline proteases. Journal of Agricultural and Food Chemistry. Vol.48(3): 657-666.

Kusmawarti, A., & Indriati, N. (2008). Daya hambat ekstrak bahan aktif biji picung (*Pangium edule Reinw.*) terhadap pertumbuhan bakteri penghasil histamin. Jurnal Pascapanen dan Bioteknologi Kelautan dan Perikanan, 3, 29-35.

Liceaga-Gesualdo, A.M., and Li-chan, E.C.Y. 1999. Functional Properties of Fish Protein Hydrolisate from Herring (*Chupea harengus*). Journal of Food Science 64(6): 1000-1004.

Lin, C.-S., Liu, F.-L., Lee, Y.-C., Hwang, C.-C., & Tsai, Y.-H. 2012. Histamine contents of salted seafood products in Taiwan and isolation of halotolerant histamine-forming bacteria. Food Chemistry, 131(2), 574–579.



Matondang, S.E. 2022. Comparison of protein levels in freshwater fish and saltwater fish. Chemistry Education Journal Vol 1(1).

Mc Lauchin J., C.L. Little, K.A. Grant, dan V. Mithani. 2005. Scombrotoxic fish poisoning. Journal of Public Health Andvance 10(1):10-93.

Nahla T.K. dan H.E. Farag. 2005. Histamine and histamine producing bacteria in some local and imported fish and their public health significance. Research Journal of Agriculture and Biological Sciences 1(4): 329-336.

Ndaw A, et al. 2007. Assessment of histamine formation during fermentation of sardine (*Sardina pilchardus*) with lactic acid bacteria. World Journal of Diary and Food Science 2(2): 42-48.

Najafian, L., dan A.S Babji. 2012. A review of fish derived antioxidant and antimicrobial peptides: their production, assessment and applications. Peptides 33(2): 178-185.

Niven, C.F. Jr., Jeffrey, M.B. and Corlet, D.A. Jr. 1981. Differential plating medium for quantitative detection of histamine-producing bacteria. App Environ. Microbiol. 41: 321-322.

Nurhayati, T., Salamah, E., Cholifah, Nugraha, R. 2014. Optimasi proses pembuatan hidrolisat jeroan ikan kakap putih. JPHPI Vol 17(1): 42-52.

Ovissipour, M., A.A. Kenari, A. Motamedzadegan, B. Rasco, dan R.M. Nazari. 2011. Optimization of protein recovery during hydrolysis of yellow fin tuna (*Thunnus albacares*) visceral proteins. Journal of Aquatic Food Product Technology 20: 148-159.

Pigot, G. M. dan Tucker, B.W. 1990. Utility Fish Flesh Eff Ectively While Maintaining Nutritional Qualities. Seafood Efecct of Technology on Nutrition. New York.

Pinto, C.A.S.O., Green, D., Baby, A.R., Ruas, G.W., Kaneko, T.M., Marana, S.R., Velasco, M.V.R. 2007. Determination of papain activity in topical dosage forms: single laboratory validation assay. Latin American Journal of Pharmacy, 26 (5) : 771-775.

Putra, M.M.P., Al-Hammam, M.Y., Ahsan, G., Chandra, K.K.B. and Puspita, I.D., Antibacterial potency of cell free supernatant produced by lactic acid bacteria isolated from Indonesian fermented fish products against histamine-producing bacteria. Saintek Perikanan: Indonesian Journal of Fisheries Science and Technology, 18(1), pp.13-18.

Rahmawati, L.S. 2013. Pemanfaatan Belut (*Monopterus albus* Zuieuw) dalam Pembuatan Bakso. Fakultas Ekologi Manusia. Institut Pertanian Bogor. Skripsi.

Reed, G. 1975. Enzymes in Food Processing. Academic Press. New York.

Roy, R. 2009. Budidaya Bisnis Belut. Jakarta: Agro Media Pustaka. hal. 126.



UNIVERSITAS
GADJAH MADA

Pemgaruh Lama Hidrolisis Terhadap Aktivitas Antibakteri Hidrolisat Protein Jeroan Belut Pada

Bakteri

Pembentuk Histamin

Anas Zulhanafi, Mgs. M. Prima Putra S.Pi., M.Sc., Ph.D.

Universitas Gadjah Mada, 2024 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Sally, H.A., Price, R.S. and Brown, W. 1980. Histamine formation by bacteria isolated from skipjack tuna. Bulletin of the Japanese Society of Scientific Fisheries. 46: 991-995.

Ulacesan, S., Kuppusamy, A., & Kim, H.J. (2018). Antimicrobial and antioxidant activities of protein hydrolysate from terrestrial snail *Cryptozona bistriata*. Journal of Applied Pharmaceutical Science. Vol 8(12):012-019.

Wei, C.I., Chen, J.A., Koburger, J.A., Otwell, W.S. and Marshall, M.R. 1990. Bacterial growth and histamine production on vacuum package tuna. J. Food Sci. 55: 59-63.

Zahid, M., Ashraf, M., Arshad, M., Ghulam, M., Yasmin, A., Hameed H.M.A. 2015. Antimicrobial activity of bacteriocins isolated acid bacteria against resistant pathogenic strain. International Journal of Nutrition and Food Sciences. 4(3): 326-331.