



## DAFTAR PUSTAKA

- Adeyemo, O.S., Chavarriaga, P., Tohme, J., Fregene, M., Davis, J.D., & Setter, T.L. (2017). Overexpression of Arabidopsis *FLOWERING LOCUS T (FT)* gene improves floral development in cassava (*Manihot esculenta*, Crantz). *PLOS ONE*, 12(7): 1-15.
- Adeyemo, O.S., Hyde, P.T., & Setter, L.T. (2019). Identification of FT family genes that respond to photoperiod, temperature and genotype in relation to flowering in cassava (*Manihot esculenta*, Crantz). *Plant Reproduction*, 32(2): 181-191.
- Allifah, A.N. & Rijal, M. (2018). Lama penyimpanan stek terhadap pertumbuhan tanaman ubi kayu (*Manihot esculenta* Crantz). *Jurnal Biology Science & Education*, 7(2): 118-126.
- Alves, A. A. C. (2002). *Cassava Biology and Physiology*. Chap. 5." In Cassava: Biology, Production and Utilization. New York: CABI Publishing. pp. 67–89.
- Amanda, U.D. & Cartealy, I.C. (2015). Isolasi RNA total dari mesokarp buah kelapa sawit (*Elaeis guineensis* Jacq. var. Tenera). *Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia*, 1(2): 171-176.
- Andrés, F. & Coupland, G. (2012). The genetic basis of flowering responses to seasonal cues. *Nature reviews. Genetics*, 13(9): 627–639.
- Ariani, L.N., Estiasih, T., & Martati. (2017). Karakteristik sifat fisiko kimia ubi kayu berbasis kadar sianida. *Jurnal Teknologi Pertanian*, 18(2): 119-128.
- Banunaek, A.F., Hendrik, A. Ch., Ngginak, J., Sabuna, A.Ch., & Meha, A.M. (2023). Inventarisasi keragaman kultivar ubi kayu (*Manihot esculenta* Crantz) dan ubi jalar (*Ipomoea batatas* L.) di Kabupaten Timor Tengah Selatan. *Jurnal Pendidikan dan Sains Biologi*, 6(1): 15-28.
- Behnam, B., Higo, A., Yamaguchi, K., Tokunaga, H., Utsumi, Y., Selvaraj, M.G., Seki, M., Ishitani, M., Lopez-Lavalle, L.A.B., & Tsuji, H. (2021). Field-transcriptome analyses reveal developmental transitions during flowering in cassava (*Manihot esculenta* Crantz). *Plant Molecular Biology*, 106(3): 285-296.
- Blümel, M., Dally, N., & Jung, C. (2015). Flowering time regulation in crops—what did we learn from *Arabidopsis*??. *Current Opinion In Biotechnology*, 32: 121–129.
- Buschmann, H., Potter, U.J. & Beeching, J.R. (2002). Ultrastructure of cassava root by TEM & SEM. *Specimen Processing Microscopy and Analysis*. pp. 9-11.
- Cai, W. M., Chionh, Y. H., Hia, F., Gu, C., Kellner, S., McBee, M. E., Ng, C. S., Pang, Y. L., Prestwich, E. G., Lim, K. S., Babu, I. R., Begley, T. J., & Dedon, P. C. (2015). A platform for discovery and quantification of modified ribonucleosides in RNA: application to stress-induced reprogramming of tRNA modifications. *Methods in Enzymology*, 560: 29–71.
- Caniago, M., Roslim, D.I., & Herman. (2014). Deskripsi karakter morfologi ubi kayu (*Manihot esculenta* Crantz) juray dari Kabupaten Rokan Hulu. *JOM FMIPA*, 1(2): 613-619.
- Cao, H. & Shockley, J.M. (2012). Comparison of TaqMan and SYBR Green qPCR methods for quantitative gene expression in tung tree tissues. *Journal of agricultural and food chemistry*, 60(50): 12296-12303.



- Ceballos, H., Morante., N., Calle, F., Lenis, J. I., Jaramillo, G., & J.C. Perez. (2002). *Mejoramiento genetico de la Yuca en La Yuca en el Tercer Milenio : Sistemas Modernos de Produccion, Procesamiento, Utilizacion y Comercializacion*. Colombia: Centro Internacional de Agricultura Tropical. pp. 295–325.
- Ceballos, H. & De la Cruz, A. (2012). *Cassava taxonomy and morphology*. In: Ospina Patiño, Bernardo; Ceballos, Hernán (eds.). *Cassava in the third millennium: Modern production, processing, use, and marketing systems*. Cali: Technical Centre for Agricultural and Rural Cooperation (CTA). pp. 15-28.
- Chacón, J., Madriñán, S., Debouck, D., Rodriguez, F., & Tohme, J. (2008). Phylogenetic patterns in the genus *Manihot* (Euphorbiaceae) inferred from analyses of nuclear and chloroplast DNA regions. *Molecular Phylogenetics and Evolution*, 49(1): 260–267.
- Chen, Q., Payyavula, R.S., Chen, L., Zhang, J., Zhang, C., & Turgeon, R. (2018). *FLOWERING LOCUS T* mRNA is synthesized in specialized companion cells in *Arabidopsis* and maryland mammoth tobacco leaf veins. *PNAS*, 115(11): 2830-2835.
- Cheng, X.F. & Wang, Z.Y. (2005). Overexpression of *COL9*, a *CONSTANS-LIKE* gene, delays flowering by reducing expression of *CO* and *FT* in *Arabidopsis thaliana*. *The Plant Journal*, 43(5): 758-768.
- Chong, G. O., Han, H. S., Lee, S. D., & Lee, Y. H. (2020). Improvement in RNA quantity and quality in cervico-vaginal cytology. *Virology journal*, 17(1): 8.
- Colasanti, J. & Sundaresan, V. (2000). Florigen enters the molecular age: long-distance signals that cause plants to flower. *Trends in Biochemical Sciences*, 25(5): 236-240.
- Corbesier, L., Vincent, C., Jang, S., Fornara, F., Fan, Q., Searle, I., Giakountis, A., Farrona, S., Gissot, L., Turnbull, C., & Coupland, G. (2007). FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science*, 316(5827): 1030–1033.
- Craciun, L., Spinette, S. A., Rassy, M., Salgado, R., de Wind, A., Demetter, P., Verset, L., Gomez-Galdon, M., Chintinne, M., Sirtaine, N., de St Aubain, N., Laios, I., Roy, F., & Larsimont, D. (2019). Tumor banks: a quality control scheme proposal. *Frontiers in medicine*, 6: 225.
- Daohong, W., Bochu, W., Biao, L. Chuanren, D., & Jin, Z. (2004). Extraction of total RNA from *Chrysanthemum* containing high levels of phenolic and carbohydrates. *Colloids and surfaces. B, Biointerfaces*, 36(2): 111–114.
- Dash P. K. (2013). High quality RNA isolation from ployphenol-, polysaccharide- and protein-rich tissues of lentil (*Lens culinaris*). *3 Biotech*, 3(2): 109–114.
- Dominguez, M.S.C.E. (1983). *Morphology of the Cassava Plant*. Colombia: Centro Internacional de Agricultura Tropical. pp. 6-26.
- Ekanayake, I.J., Osiru, D.S.O., & Porto, M.C.M. (1997). *Morphology of Cassava*. Nigeria: International Institute of Tropical Agriculture (IITA). pp. 3-30.
- Escobar, M.D. & Hunt, J.L. (2017). A cost-effective RNA extraction technique from animal cells and tissue using silica columns. *Journal of Biological Methods*, 4(2): 1-5.
- Faatih, M. (2009). Isolasi dan digesti DNA kromosom. *Jurnal Penelitian Sains dan Teknologi*, 10(1): 61-67.



- Farrell, R.E. (2010). *cDNA Synthesis in RNA Methodologies*. pp. 367-384.
- Firdaus, N.R., Hayati, P.K. D., & Yusniwati. (2016). Karakteristik fenotipik ubi kayu (*Manihot esculenta* Crantz) lokal Sumatera Barat. *Jurnal Agroteknologi*, 10(1): 104-116.
- Fitriani, H., Rahman, N., Kurniawati, S., Aryaningrum, P.D., & Hartati, N.S. (2019). Isolasi dan kultur protoplas mesofil daun dari beberapa genotip ubi kayu (*Manihot esculenta* Crantz). *Biopropal Industri*, 10(1): 1-13.
- Food and Agriculture Organizations of the United Nations (FAO). (2013). *Save and Grow: Cassava*. Rome: FAO of The United Nations. pp. 1-46.
- Fornara, F., de Montaigu, A., & Coupland, G. (2010). SnapShot: control of flowering in *Arabidopsis*. *Cell*, 141(3): 550.
- Guan, L., Ma, X., Zhou, X., Tan, B., & Wang, Z. Y. (2019). An optimized method to obtain high-quality RNA from cassava storage root. *3 Biotech*, 9(4): 118.
- Halsey, M.E., Olsen, K.M., Taylor, N.J., & Chavarriaga-Aguirre, P. (2008). Reproductive biology of cassava (*Manihot esculenta* Crantz) and isolation of experimental field trials, *Crop Science Society of America*, 48(1): 49-58.
- Hasanah, L.M., Puspito, A.N., Arimurti, S., & Su'udi M. (2021). Identifikasi WIN1 (wax inducer1) pada tanaman ubi kayu (*Manihot esculenta* Crantz). *Al-Hayat: Journal of Biology and Applied Biology*, 4(1): 95-103.
- Hasegawa, T., Takahashi, J., & Iwahashi, H. (2018). RNA Quality Control Using External Standard RNA. *Polish Journal of Microbiology*, 67(3): 347–353.
- Hasim, Falah, S., & Dewi, L.K. (2016). Effect of boiled cassava leaves (*Manihot esculenta* Crantz) on total phenolic, flavonoid and its antioxidant activity. *Current Biochemistry*, 3(3): 116-127.
- Hayama, R. & Coupland, G. (2003). Shedding light on the circadian clock and the photoperiodic control of flowering. *Current opinion in plant biology*, 6(1): 13–19.
- Hermanto, Sadikin W., D., & Hikmat, E. (2009). *Deskripsi Varietas Unggul Palawija 1918-2009*. Bogor: Pusat Penelitian dan Pengembangan Tanaman Pangan. pp. 321- 330.
- Howeler, R., Lutaladio, N., & Thomas, G. (2013). *Save and Grow: Cassava, a Guide to Sustainable Production Intendification*. Rome: FAO of The United Nations. pp. 1-18.
- Howeler, R. & T. M. Aye. (2014). *Sustainable Management of Cassava in Asia: From Research to Practice*. Colombia: International Center for Tropical Agriculture (CIAT). pp. 25-40.
- Hyde, P.T. & Setter, T.L. (2022). Long-day photoperiod and cool temperature induce flowering in cassava: Expression of signaling genes. *Frontiers in Plant Science*, 13: 1-17.
- Integrated Taxonomy Information System (ITIS). (2023). “*Manihot esculenta* Crantz”. [https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=503688#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=503688#null). [Diakses 3 April 2023].
- Joshi, C. J., Ke, W., Drangowska-Way, A., O'Rourke, E. J., & Lewis, N. E. (2022). What are housekeeping genes?. *PLOS Computational Biology*, 18(7): 1-19.
- Junistia, R.A., Kardhinata, E. H., & Hanafiah, D. S. (2018). Pertumbuhan dan pekembangan beberapa genotip ubi kayu (*Manihot esculenta* Crantz).



- Jurnal Agroekoteknologi FP USU*, 6(1): 68-76.
- Khanna, R., Kronmiler, B., & Maszle, D.R. (2009). The *Arabidopsis* B-Box zinc finger family. *Plant Cell*, 21(11): 3416-3420.
- Kim, S.Y., Yu, X., & Michaels, S.D. (2008). Regulation of *CONSTANS* and *FLOWERING LOCUS T* expression in response to changing light quality. *Plant Physiology*, 148(1): 269-279.
- Koonjul, P. K., Brandt, W. F., Farrant, J. M., & Lindsey, G. G. (1999). Inclusion of polyvinylpyrrolidone in the polymerase chain reaction reverses the inhibitory effects of polyphenolic contamination of RNA. *Nucleic acids research*, 27(3): 915-916.
- Kuete, V. (2014). *Physical, Hematological, and Histopathological Signs of Toxicity Induced by African Medicinal Plants in Toxicological Survey of African Medicinal Plants*. Amsterdam: Elsevier. pp. 635-657.
- Leelawijitkul, S., Kongsil, P., Kittipadakul, P., & Juntawong, P. (2022). Correlation between relative gene expression patterns of two *FLOWERING LOCUS T* (MeFT1 and MeFT2) in cassava leaf and flowering traits under different flowering induction conditions. *Pakistan Journal of Biological Sciences: PJBS*, 25(5): 369-379.
- Liu, L., Farrona, S., Klemme, S., & Turck, F.K. (2014). Post fertilization expression of *FLOWERING LOCUS T* suppresses reproductive reversion. *Frontiers in Plant Science*, 5(164): 1-7.
- Livak, K.J. & Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods*, 25(4): 402-408.
- McHale, J.S. & Dove, L.D. (1967). Ribonuclease activity in tomato leaves as related to development and senescence. *The New Phytologist*, 67(3): 505-515.
- Mattheus, N., Ekramoddoullah, A. K., & Lee, S. P. (2003). Isolation of high-quality RNA from white spruce tissue using a three-stage purification method and subsequent cloning of a transcript from the PR-10 gene family. *Phytochemical analysis: PCA*, 14(4): 209-215.
- Moreno, I., Gruissem, W., & Vanderschuren, H. (2011). Reference genes for reliable potyvirus quantitation in cassava and analysis of cassava brown streak virus load in host varieties. *Journal of Virological Methods*, 177(1): 49-54.
- Muñoz, P. & Munné-Bosch, S. (2018). Photo-Oxidative stress during leaf, flower and fruit development. *Plant physiology*, 176(2): 1004-1014.
- Novaldi, A.A., Miranda, C., & Nurhayati, A.D. (2022). Teknik budi daya dan karakteristik ubi kayu (*Manihot esculenta Crantz*) di Desa Leuwisadeng, Kecamatan Leuwisadeng, Kabupaten Bogor, Jawa Barat. *Jurnal Pusat Inovasi Masyarakat*, 4(1): 8-16.
- Nwachukwu, I.D. & Simonyan, K.J. (2015). Some engineering properties of cassava tuber related to its peeling mechanization. *Umudike Journal of Engineering and Technology (UJET)*, 1(1): 12-24.
- Odipio, J., Getu, B., Chauhan, R.D., Alicai, T., Bart, R., Nusinow, D.A., & Taylor, N.J. (2020). Transgenic overexpression of endogenous *FLOWERING LOCUS T*-like gene MeFT1 produces early flowering in cassava. *PLOS ONE*, 15(1): 1-16.



- Okai, E. (2001). Genetic diversity in some local cassava cultivars in Ghana.
- Oluwasanya, D.N., Gisel, A., Stavlone, L., & Setter, T.L. (2021). Environmental responsiveness of flowering time in cassava genotypes and associated transcriptome changes. *PLOS ONE*, 16(7): 1-24.
- Oñate-Sánchez, L. & Vicente-Carabajosa, J. (2008). DNA-free RNA isolation protocols for *Arabidopsis thaliana*, including seeds and siliques. *BMC research notes*, 1(1): 1-7.
- Pahlevi, T.R., Buaton, R., & Nurhayati. (2021). Identifikasi jenis bunga menggunakan ekstraksi ciri orde satu dan algoritma multi support-vector machines (multisvm). *Jurnal Informatika Kaputama (JIK)*, 5(1): 116-128.
- Pin, P.A. & Nilsson, O. (2012). The multifaceted roles of *FLOWERING LOCUS T* in plant development. *Plant, Cell & Environment*, 35(10): 1742-1755.
- Pramesti, F.S., Rahayu, E.S., & Agustono. (2017). Analisis daya saing ubi kayu Indonesia di pasar internasional. *SEPA*, 14(1): 1-7.
- Pratiwi, E., & Widodo, L. (2020). Kuantifikasi hasil ekstraksi gen sebagai faktor kritis untuk keberhasilan pemeriksaan RT PCR. *Indonesian Journal for Health Sciences*, 4(1): 1-9.
- Purnomo, B.H., Subayri, A., & Kuswardhani, N. (2015). Model sistem dinamik ketersediaan singkong bagi industri tape di Kabupaten Jember. *Jurnal Agroteknologi*, 9(2): 162-173.
- Putterill, J., Robson, F., Lee, K., Simon, R., & Coupland, G. (1995). The *CONSTANS* gene of *Arabidopsis* promotes flowering and encodes a protein showing similarities to zinc finger transcription factors. *Cell*, 80(6): 847-857.
- Putterill, J. & Varkonyi-Gasic, E. (2016). FT and florigen long-distance flowering control in plants. *Current Opinion in Plant Biology*, 33: 77-82.
- Rahman, A. Y., Usharraj, A. O., Misra, B. B., Thottathil, G. P., Jayasekaran, K., Feng, Y., Hou, S., Ong, S. Y., Ng, F. L., Lee, L. S., Tan, H. S., Sakaff, M. K., Teh, B. S., Khoo, B. F., Badai, S. S., Aziz, N. A., Yuryev, A., Knudsen, B., Dionne-Laporte, A., Mchunu, N. P., Yu, Q., Langston, B.J., Freitas, T.A., Young, A.G., Chen, R., Wang, L., Najimudin, N., Saito, J.A., & Alam, M. (2013). Draft genome sequence of the rubber tree *Hevea brasiliensis*. *BMC genomics*, 14: 75.
- Rahmawati, Siswoyo, T.A., Restanto, D.P., Hartatik, S., Soeparjono, S., & Avivi, S. (2017). Morphological and physiological characters of cassava (*Manihot esculenta* Crantz) which wet tolerant. *Proceeding The 1st International Basic Science Conference: Towards The Extended Use of Basic Science for Enhancing*. pp. 32-35.
- Ramadanti, N.A., Hartati, Putri, D.H., & Hartati, N.S. (2020). Karakteristik molekuler variasi somaklonal ubi kayu varietas Carvita 25 menggunakan marka SSR. *Jurnal ILMU DASAR*, 21(2): 87-96.
- Rauwane, M. E., Odeny, D. A., Millar, I., Rey, C., & Rees, J. (2018). The early transcriptome response of cassava (*Manihot esculenta* Crantz) to mealybug (*Phenacoccus manihoti*) feeding. *PLOS ONE*, 13(8): 1-26.
- Restiani, R., Roslim, D.I., & Herman. (2014). Karakter morfologi ubi kayu (*Manihot esculenta* Crantz) hijau dari Kabupaten Pelalawan. *JOM FMIPA*, 1(2): 619-623.
- Reuben-Kalu, J.I., Eswaran, K., Muthurajan, R., Doraiswamy, U., Venkatasamy,



- B., & Shanmugam, K.P. (2023). Precise isolation of high-quality RNA from leaves and storage roots of cassava (*Manihot esculenta* Crantz) for gene expression studies. *Bulletin of the National Research Centre*, 47: 84.
- Robson, F., Costa, M. M., Hepworth, S. R., Vizir, I., Piñeiro, M., Reeves, P. H., Putterill, J., & Coupland, G. (2001). Functional importance of conserved domains in the flowering-time gene *CONSTANS* demonstrated by analysis of mutant alleles and transgenic plants. *The Plant journal: For Cell and Molecular Biology*, 28(6), 619–631.
- Rodríguez-Falcón, M., Bou, J., & Prat, S. (2006). Seasonal control of tuberization in potato: conserved elements with the flowering response. *Annual Review of Plant Biology*, 57, 151–180.
- Rogers, D.J. & Appan, S.G. (1973). *Manihot Manihotoides (Euphorbiaceae) in Flora neotropica Vol.13*. Newyork: Hafner Press. pp. 1-272.
- Roy, S.S., Dasgupta, R., & Bagchi, A. (2014). A review on phylogenetic analysis: a journey through modern era. *Computational Molecular Bioscience*, 4(3): 39-45.
- Rozi, F. & Heriyanto. (2012). Peluang produksi ubi kayu Madura dalam pengembangan agroindustri berbasis komoditas lokal. *Jurnal Cakrawala*, 7(1): 1-8.
- Samach, A. & Coupland, G. (2000). Time measurement and the control of flowering in plants. *BioEssays*, 22(1): 38–47.
- Steinbach, Y. (2019). The *Arabidopsis thaliana* CONSTANS-LIKE 4 (COL4) - a modulator of flowering time. *Frontiers in Plant Science*, 10: 651.
- Shalit-Kaneh, A., Eviatar-Ribak, T., Horev, G., Suss, N., Aloni, R., Eshed, Y., & Lifschitz. (2019). The flowering hormone florigen accelerates secondary cell wall biogenesis to harmonize vascular maturation with reproductive development. *PNAS*, 116(32): 16127-16136.
- Shiade, S.R.G., Zand-Silakhoor, A., Fathi, A., Rahimi, R., Minkina, T., Rajput, V.D., Zulfiqar, U., & Chaudhary T. (2024). Plant metabolites and signaling pathways in response to biotic and abiotic stresses: Exploring bio stimulant applications. *Plant Stress*, 12: 100454.
- Shim, J.S., Kubota, A., & Imaizumi, T. (2017). Circadian clock and photoperiodic flowering in *Arabidopsis*: *CONSTANS* is a hub for signal integration. *Plant Physiology*, 173(1): 5-15.
- Siebers, T., Catarino, B., & Agusti, J. (2017). Identification and expression analyses of new potential regulators of xylem development and cambium activity in cassava (*Manihot esculenta*). *Planta*, 245(3): 539-548.
- Song, Y.H., Estrada, D.A., Johnson, R.S., Kim, S.K., Lee, S.Y., MacCoss, M.J., & Imaizumi, T. (2014). Distinct roles of *FKF1*, *GIGANTEA*, and *ZEITLUPE* proteins in the regulation of *CONSTANS* stability in *Arabidopsis* photoperiodic flowering. *PNAS*, 111(49): 17672-17677.
- Suárez-López, P., Wheatley, K., Robson, F., Onouchi, H., Valverde, F., & Coupland, G. (2001). *CONSTANS* mediates between the circadian clock and the control of flowering in *Arabidopsis*. *Nature*, 410(6832): 1116–1120.
- Tiwari, S.B., Shen, Y., Chang, H.C., Hou, Y., Harris, A., Ma, S.F., McPartland, M., Hymus, G.J., Adam, L., Marion, C., Belachew, A., Repetti, P.P., Reuber, T.L., & Ratcliffe, O.J. (2010). The flowering time regulator *CONSTANS* is recruited to the *FLOWERING LOCUS T* promoter via a



- unique cis-element. *New Phytologist*, 187(1): 57-66.
- Tonukari, N. J., Ezedom, T., Enuma, C.C., Sakpa, S.O., Avwioroko, O.J., Eraga, L., & Odiyoma, E. (2015). White gold: cassava as an industrial base. *American Journal of Plant Sciences*, 6(7): 972-979.
- Tsuji, H. (2017). Molecular function of florigen. *Breeding science*, 67(4): 327-332.
- Turck, F., Fornara, F., & Coupland G. (2008). Regulation and identity of florigen: flowering locus t moves center stage. *Annual review of plant biology*, 59: 573-594.
- Utama, Y.A.K. & Rukismono, M. (2018). Singkong-Man Vs Gadung-Man. Papua: Aseni. pp. 15-22.
- Villamayor-Jr., F. G. (1987). Ecological and cultur requirements of cassava. *Philippine Root Crops Information Service*, 2(1): 1-4.
- Waluyo, T. (2020). Pemanfaatan hormon tumbuh organik untuk meningkatkan produktivitas singkong hasil eksplorasi seleksi bibit unggul. *Jurnal Ilmu dan Budaya*, 41(70): 8207-8217.
- Wan, P. & Che, D. (2013). Constructing phylogenetic trees using interacting pathways. *Bioinformation*, 9(7): 363-367.
- Wang, H., Zhang, Z., Li, H., Zhao, X., Liu, X., Ortiz, M., Lin, C., & Liu, B. (2013). CONSTANS-LIKE 7 regulates branching and shade avoidance response in *Arabidopsis*. *Journal of Experimental Botany*, 64(4), 1017-1024.
- Xu, Z., Huang, J., Gao, M., Guo, G., Zeng, S., Chen, X., Wang, X., Gong, Z., & Yan, Y. (2021). Current perspectives on the clinical implications of oxidative RNA damage in aging research: challenges and opportunities. *GeroScience*, 43(2): 487-505.
- Yoshino, K., Nishijima, R., & Kawakatsu, T. (2020). Low-cost RNA extraction method for highly scalable transcriptome studies. *Breeding science*, 70(4): 481-486.
- Zannou, I.D., Hanna, R., de Moraes, G.J., Kreiter, S., Phiri, G., & Jone, A. (2014). Mites of cassava (*Manihot esculenta* Crantz) habitats in Southern. *International Journal of Acarology*, 31(2): 149-164.