

DAFTAR PUSTAKA

- Abdallah, M. M. S., Abdelgawad, Z. A., & El-Bassiouny, H. M. S. (2016). Alleviation of the adverse effects of salinity stress using trehalose in two rice varieties. *South African Journal of Botany*, 103, 275–282. <https://doi.org/10.1016/j.sajb.2015.09.019>
- Abdullah, Z., Khan, M. A., & Flowers, T. J. (2001). Causes of sterility in seed set of rice under salinity stress. *Journal of Agronomy and Crop Science*, 187(1), 25–32. <https://doi.org/10.1046/j.1439-037X.2001.00500.x>
- Aher, A. N., Pal, S. C., Yadav, S. K., Patil, U. K., & Bhattacharya, S. (2009). Antioxidant activity of isolated phytoconstituents from *Casuarina equisetifolia* Frost (Casuarinaceae). In *Journal of Plant Sciences* (Vol. 4, Issue 1, pp. 15–20). <https://doi.org/10.3923/jps.2009.15.20>
- Ahn, C., Park, U., & Park, P. B. (2011). Increased salt and drought tolerance by D-ononitol production in transgenic *Arabidopsis thaliana*. *Biochemical and Biophysical Research Communications*, 415(4), 669–674. <https://doi.org/10.1016/j.bbrc.2011.10.134>
- Amirjani, M. R. (2010). Effect of NaCl on Some Physiological Parameters of Rice. *Ejbs*, 3(June), 6–16.
- Anantha, M. S., Patel, D., Quintana, M., Swain, P., Dwivedi, J. L., Torres, R. O., Verulkar, S. B., Variar, M., Mandal, N. P., Kumar, A., & Henry, A. (2016). Trait combinations that improve rice yield under drought: Sahbhagi Dhan and new drought-tolerant varieties in South Asia. *Crop Science*, 56(1), 408–421. <https://doi.org/10.2135/cropsci2015.06.0344>
- André, P., Freitas, F. De, Souza, R. De, Elton, M., Marques, C., & Tarquinio, J. (2018). Salt Tolerance Induced by Exogenous Proline in Maize Is Related to Low Oxidative Damage and Favorable Ionic Homeostasis. *Journal of Plant Growth Regulation*, 0(0), 0. <https://doi.org/10.1007/s00344-018-9787-x>
- Arora, A., Sairam, R. K., & Srivastava, G. C. (2002). Oxidative stress and antioxidative system in plants. *Current Science*, 82(10), 1227–1238.
- Basuki, T. (2014). *Kontribusi Senyawa Kompatibel ekstrak daun tumbuhan halofit sebagai anti-salinitas dalam amandemen fisiologis padi (*Oryza sativa* L.)*.
- Bates, L. S., Waldren, R. C., & Teare. (1973). SHORT COMMUNICATION Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39, 205–207.
- Berwal, O. K., & Ram, C. (2018). Superoxide Dismutase: A Stable Biochemical Marker for Abiotic Stress Tolerance in Higher Plants. *Abiotic and Biotic Stress in Plants*, 1, 1–10. doi: <http://dx.doi.org/10.5772/intechopen.82079>
- Bhattacharjee, S. (2005). Reactive oxygen species and oxidative burst: Roles in

stress, senescence and signal transduction in plants. *Current Science*, 89(7), 1113–1121.

Brennan, J. P., & Malabayabas, A. (2011). International Rice Research Institute's contribution to rice varietal yield improvement in South-East Asia. *Acicar*, 111. https://books.google.com.sg/books?id=X-ibnQAACAAJ&dq=International+Rice+Research+Institute's+contribution+to+rice+varietal+yield+improvement+in+South-East+Asia&hl=en&sa=X&redir_esc=y

Carillo, P., Grazia, M., Pontecorvo, G., Fuggi, A., & Woodrow, P. (2011). Salinity Stress and Salt Tolerance. *Abiotic Stress in Plants - Mechanisms and Adaptations*, September. <https://doi.org/10.5772/22331>

Cha-um, S., Siringam, K., Juntawong, N., & Kirdmanee, C. (2010). Water relations, pigment stabilization, photosynthetic abilities and growth improvement in salt stressed rice plants treated with exogenous potassium nitrate application. *International Journal of Plant Production*, 4(3)(January 2016), 187–198.

Chanda, S. V, & Singh, Y. D. (2002). Estimation of leaf area in wheat using linear measurements. *Plant Breeding and Seed Science*, 46(2), 75–79.

Chen, H., & Jiang, J. G. (2010). Osmotic adjustment and plant adaptation to environmental changes related to drought and salinity. *Environmental Reviews*, 18(1), 309–319. <https://doi.org/10.1139/A10-014>

Chutipaijit, S., Cha-um, S., & Sompornpailin, K. (2011). High contents of proline and anthocyanin increase protective response to salinity in *Oryza sativa* L. spp. indica. *Australian Journal of Crop Science*, 5(10), 1191–1198.

Del Rio, L. A., Corpas, F. J., Sandalio, L. M., Palma, M., Go, M., & Barroso, J. B. (2002). *Reactive oxygen species, antioxidant systems and nitric oxide in peroxisomes*. 53(372), 1255–1272.

Elavarthi, S., & Martin, B. (2010). Spectrophotometric Assays for Antioxidant Enzymes in Plants Sathya. R. Sunkar (Ed.), *Plant Stress Tolerance, Methods in Molecular Biology* 639, R. Sunkar, 273.

FAO. (2018). World Food and Agriculture Statistical Pocketbook. In *The Lancet* (Vol. 274, Issue 7102).

Fichman, Y., Gerdes, S. Y., Kovács, H., Szabados, L., Zilberstein, A., & Csonka, L. N. (2015). Evolution of proline biosynthesis: Enzymology, bioinformatics, genetics, and transcriptional regulation. *Biological Reviews*, 90(4), 1065–1099. <https://doi.org/10.1111/brv.12146>

Fitzgerald, M. A., McCouch, S. R., & Hall, R. D. (2009). Not just a grain of rice: the quest for quality. *Trends in Plant Science*, 14(3), 133–139. <https://doi.org/10.1016/j.tplants.2008.12.004>

- Foyer, C. H., & Noctor, G. (2009). Redox Regulation in Photosynthetic Organisms. *Regulation*, 11(4), 861–905.
<https://doi.org/10.1089/ars.2008.2177>
- Funck, D., Stadelhofer, B., & Koch, W. (2008). Ornithine- δ -aminotransferase is essential for Arginine Catabolism but not for Proline Biosynthesis. *BMC Plant Biology*, 8, 1–14. <https://doi.org/10.1186/1471-2229-8-40>
- Galvani, A. (2007). The challenge of the food sufficiency through salt tolerant crops Adriana. *Rev Environ Sci Biotechnol*, 6(August), 3–16.
<https://doi.org/10.1007/978-1-4020-6285-8>
- Ghosh, B., & Ali Md, N. (2016). Response of Rice under Salinity Stress: A Review Update. *Rice Research: Open Access*, 4(2), 2–9.
<https://doi.org/10.4172/2375-4338.1000167>
- Gill, S. S., & Tuteja, N. (2010). Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry*, 48(12), 909–930. <https://doi.org/10.1016/j.plaphy.2010.08.016>
- Guo, Z., Ou, W., Lu, S., & Zhong, Q. (2006). Differential responses of antioxidative system to chilling and drought in four rice cultivars differing in sensitivity. *Plant Physiology and Biochemistry*, 44(11–12), 828–836.
<https://doi.org/10.1016/j.plaphy.2006.10.024>
- Gupta, B., & Huang, B. (2014). Mechanism of salinity tolerance in plants: physiological, biochemical, and molecular characterization. *International Journal of Genomics*, 2014, 1–18.
<https://doi.org/http://dx.doi.org/10.1155/2014/701596>
- Halliwell, B. (2006). Reactive species and antioxidants. Redox biology is a fundamental theme of aerobic life. *Plant Physiology*, 141(2), 312–322.
<https://doi.org/10.1104/pp.106.077073>
- Harborne. (1993). Phytochemical Methods. A guide to modern techniques of plant analysis. In *Chapman and Hall* (Vol. 21, Issue 8).
[https://doi.org/10.1016/0305-1978\(93\)90098-c](https://doi.org/10.1016/0305-1978(93)90098-c)
- Havaux, M. (1998). Carotenoids as membrane stabilizers in chloroplasts. *Trends in Plant Science*, 3(4), 147–151. [https://doi.org/10.1016/S1360-1385\(98\)01200-X](https://doi.org/10.1016/S1360-1385(98)01200-X)
- Hendriyani, I. S., & Setiari, N. (2009). KANDUNGAN KLOROFIL DAN PERTUMBUHAN KACANG PANJANG ($Vigna W = (T_b - T_k) \times 100 \%$). *J. Sains & Mat.*, 17 no.3, 145–150.
- Hnilickova, H., Kraus, K., Vachova, P., & Hnilicka, F. (2021). Salinity stress affects photosynthesis, malondialdehyde formation, and proline content in portulaca oleracea l. *Plants*, 10(5). <https://doi.org/10.3390/plants10050845>
- Hoang, T. M. L., Tran, T. N., Nguyen, T. K. T., Williams, B., Wurm, P., Bellairs,

- S., & Mundree, S. (2016). Improvement of salinity stress tolerance in rice: Challenges and opportunities. *Agronomy*, 6(4), 1–23. <https://doi.org/10.3390/agronomy6040054>
- Horie, T., Karahara, I., & Katsuhara, M. (2012). Salinity tolerance mechanisms in glycophytes: An overview with the central focus on rice plants. *Rice*, 5(1), 1–18. <https://doi.org/10.1186/1939-8433-5-11>
- Huang, Y., Bie, Z., Liu, Z., Zhen, A., & Wang, W. (2009). Protective role of proline against salt stress is partially related to the improvement of water status and peroxidase enzyme activity in cucumber. *Soil Science and Plant Nutrition*, 55(5), 698–704. <https://doi.org/10.1111/j.1747-0765.2009.00412.x>
- Ighodaro, O. M., & Akinloye, O. A. (2018). First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defence grid. *Alexandria Journal of Medicine*, 54(4), 287–293. <https://doi.org/10.1016/j.ajme.2017.09.001>
- Jain, M. (2013). Emerging Role of Metabolic Pathways in Abiotic Stress Tolerance. *Journal of Plant Biochemistry & Physiology*, 01(02), 10–11. <https://doi.org/10.4172/2329-9029.1000108>
- Kavi Kishor, P. B., Sangam, S., Amrutha, R. N., Sri Laxmi, P., Naidu, K. R., Rao, K. R. S. S., Rao, S., Reddy, K. J., Theriappan, P., & Sreenivasulu, N. (2005). Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: Its implications in plant growth and abiotic stress tolerance. *Current Science*, 88(3), 424–438.
- Kementrian pertanian. (2020). Statistik Pertanian Agricultural Statistics 2020. In *Kementrian Pertanian Republik Indonesia*. Pusat data dan sistem informasi pertanian kementrian pertanian republik indonesia. https://satudata.pertanian.go.id/assets/docs/publikasi/Statistik_Pertanian_Tahun_2020.pdf
- Khedra, J., Agarwal, P., & Agarwal, P. K. (2019). Deciphering hydrogen peroxide-induced signalling towards stress tolerance in plants. *3 Biotech*, 9(11), 1–13. <https://doi.org/10.1007/s13205-019-1924-0>
- Lang, F. (2007). Mechanisms and Significance of Cell Volume Regulation. *Journal of the American College of Nutrition*, 26(January 2015), 613S-623S. <https://doi.org/10.1080/07315724.2007.10719667>
- Le, T. N., & McQueen-Mason, S. J. (2006). Desiccation-tolerant plants in dry environments. *Reviews in Environmental Science and Biotechnology*, 5(2–3), 269–279. <https://doi.org/10.1007/s11157-006-0015-y>
- Li, R. hua, GUO, P. guo, Michael, B., Stefania, G., & Salvatore, C. (2006). Evaluation of Chlorophyll Content and Fluorescence Parameters as Indicators of Drought Tolerance in Barley. *Agricultural Sciences in China*,

5(10), 751–757. [https://doi.org/10.1016/S1671-2927\(06\)60120-X](https://doi.org/10.1016/S1671-2927(06)60120-X)

- Liang, X., Zhang, L., Natarajan, S. K., & Becker, D. F. (2013). Proline mechanisms of stress survival. *Antioxidants and Redox Signaling*, 19(9), 998–1011. <https://doi.org/10.1089/ars.2012.5074>
- Liu, P., Sun, F., Gao, R., & Dong, H. (2012). RAP2.6L overexpression delays waterlogging induced premature senescence by increasing stomatal closure more than antioxidant enzyme activity. *Plant Molecular Biology*, 79(6), 609–622. <https://doi.org/10.1007/s11103-012-9936-8>
- Mackill, D. J., & Khush, G. S. (2018). IR64: a high-quality and high-yielding mega variety. *Rice*, 11(1). <https://doi.org/10.1186/s12284-018-0208-3>
- Maoka, T. (2020). Carotenoids as natural functional pigments. *Journal of Natural Medicines*, 74(1), 1–16. <https://doi.org/10.1007/s11418-019-01364-x>
- Marklund, S., & Marklund, G. (1974). Involvement of the Superoxide Anion Radical in the Autoxidation of Pyrogallol and a Convenient Assay for Superoxide Dismutase. *European Journal of Biochemistry*, 47(3), 469–474. <https://doi.org/10.1111/j.1432-1033.1974.tb03714.x>
- Mehta, D., & Vyas, S. (2023). Comparative bio-accumulation of osmoprotectants in saline stress tolerating plants: A review. *Plant Stress*, 9(June), 100177. <https://doi.org/10.1016/j.stress.2023.100177>
- Miller, G., Shulaev, V., & Mittler, R. (2008). *Reactive oxygen signaling and abiotic stress*. 481–489. <https://doi.org/10.1111/j.1399-3054.2008.01090.x>
- Munns, R., & Tester, M. (2008). Mechanisms of salinity tolerance. *Annual Review of Plant Biology*, 59, 651–681. <https://doi.org/10.1146/annurev.arplant.59.032607.092911>
- Nakano, Y., & Asada, K. (1981). APX Nakano & Asada 1981.pdf. *Plant & Cell Physiology*, 22(May), 867–880.
- Nash, R. J., Thomas, P. I., Waigh, R. D., Fleet, G. W. J., Wormald, M. R., Paul, P. M., & Watkin, D. J. (1994). Casuarine: A very highly oxygenated pyrrolizidine alkaloid. *Tetrahedron Letters*, 35(42), 7849–7852. [https://doi.org/10.1016/0040-4039\(94\)80134-7](https://doi.org/10.1016/0040-4039(94)80134-7)
- Nawaz, K., & Ashraf, M. (2007). Improvement in salt tolerance of maize by exogenous application of glycinebetaine: Growth and water relations. *Pakistan Journal of Botany*, 39(5), 1647–1653.
- Nayyeripasand, L., Garosi, G. A., & Ahmadikhah, A. (2019). Selection for salinity tolerance in an international rice collection at vegetative stage. *Australian Journal of Crop Science*, 13(6), 837–846. <https://doi.org/10.21475/ajcs.19.13.06.p1327>
- Ogunwande, I. A., Flamini, G., Adefuye, A. E., Lawal, N. O., Moradeyo, S., & Avoseh, N. O. (2011). Chemical compositions of *Casuarina equisetifolia* L.,

- Eucalyptus torelliana* L. and *Ficus elastica* Roxb. ex Hornem cultivated in Nigeria. *South African Journal of Botany*, 77(3), 645–649.
<https://doi.org/10.1016/j.sajb.2011.02.001>
- Pandey, S., Fartyal, D., Agarwal, A., & Shukla, T. (2017). *Abiotic Stress Tolerance in Plants : Myriad Roles of Ascorbate Peroxidase*. 8(April), 1–13.
<https://doi.org/10.3389/fpls.2017.00581>
- Pask, A., Pietragalla, J., & Mullan, D. (2012). Physiological Breeding II: A Field Guide to Wheat Phenotyping. In *Chemistry & ...* (Issue November).
<http://onlinelibrary.wiley.com/doi/10.1002/cbdv.200490137/abstract>
- Pham, J., & Desikan, R. (2009). *ROS Signalling in Stomata*. 55–71.
https://doi.org/10.1007/978-3-642-00390-5_4
- Prayoga, M. ., Rostini, N., Setiawati, M. R., Simarmata, T., Stoeber, S., & Adinata, K. (2018). Preferences of farmers to superior rice (*Oryza sativa*) for rice fields in Pangandaran and Cilacap regions. *Jurnal Kultivasi*, 17(1)(022), 523–530.
- Ramadhan, G. R., Avivi, S., Sugiharto, B., & Fanata, W. I. D. (2021). Unfolded protein response in rice (*Oryza sativa* L.) varieties with different level of salt stress tolerance. *Indonesian Journal of Biotechnology*, 26(3), 152–158.
<https://doi.org/10.22146/ijbiotech.67039>
- Ranganayakulu, Veeranagamallaiah, G., & Sudhakar, C. (2013). Effect of salt stress on osmolyte accumulation in two groundnut cultivars (*Arachis hypogaea* L.) with contrasting salt tolerance. *African Journal of Plant Science*, 7(12), 586–592. <https://doi.org/10.5897/ajps11.063>
- Reddy, A. R., Chaitanya, K. V., & Vivekanandan, M. (2004). Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of Plant Physiology*, 161(11), 1189–1202.
<https://doi.org/10.1016/j.jplph.2004.01.013>
- Reddy, Kim, B. K., Yoon, I. S., Kim, K. H., & Kwon, T. R. (2017). Salt Tolerance in Rice: Focus on Mechanisms and Approaches. *Rice Science*, 24(3), 123–144. <https://doi.org/10.1016/j.rsci.2016.09.004>
- Reguera, M., Peleg, Z., & Blumwald, E. (2012). Targeting metabolic pathways for genetic engineering abiotic stress-tolerance in crops. *Biochimica et Biophysica Acta - Gene Regulatory Mechanisms*, 1819(2), 186–194.
<https://doi.org/10.1016/j.bbagrm.2011.08.005>
- Rossatto, T., do Amaral, M. N., Benitez, L. C., Vighi, I. L., Braga, E. J. B., de Magalhães Júnior, A. M., Maia, M. A. C., & da Silva Pinto, L. (2017). Gene expression and activity of antioxidant enzymes in rice plants, cv. BRS AG, under saline stress. *Physiology and Molecular Biology of Plants*, 23(4), 865–875. <https://doi.org/10.1007/s12298-017-0467-2>
- Rothenberg, S. E., Feng, X., Zhou, W., Tu, M., Jin, B., & You, J. (2012).

- Environment and genotype controls on mercury accumulation in rice (*Oryza sativa* L.) cultivated along a contamination gradient in Guizhou, China. *Science of the Total Environment*, 426, 272–280.
<https://doi.org/10.1016/j.scitotenv.2012.03.024>
- Shahzad, A. N., Qureshi, M. K., Ullah, S., Latif, M., Ahmad, S., & Bukhari, S. A. H. (2020). Exogenous Trehalose Improves Cotton Growth by Modulating Antioxidant Defense Under Salinity-Induced Osmotic Stress. *Pakistan Journal of Agricultural Research*, 33(2), 270–279.
<https://doi.org/10.17582/journal.pjar/2020/33.2.270.279>
- Shawon, R. A., Kang, B. S., Lee, S. G., Kim, S. K., Ju Lee, H., Katrich, E., Gorinstein, S., & Ku, Y. G. (2020). Influence of drought stress on bioactive compounds, antioxidant enzymes and glucosinolate contents of Chinese cabbage (*Brassica rapa*). *Food Chemistry*, 308(October 2019), 125657.
<https://doi.org/10.1016/j.foodchem.2019.125657>
- Singh, M., Kumar, J., Singh, S., Singh, V. P., & Prasad, S. M. (2015). Roles of osmoprotectants in improving salinity and drought tolerance in plants: a review. *Reviews in Environmental Science and Biotechnology*, 14(3), 407–426. <https://doi.org/10.1007/s11157-015-9372-8>
- Sunarto. (2001). Toleransi Kedelai terhadap Tanah Salin. *Jurnal Agronomi Indonesia (Indonesian Journal of Agronomy)*, 29(1), 27–30.
- Székel, G., Ábrahám, E., Cséplő, Á., Rigó, G., Zsigmond, L., Csiszár, J., Ayaydin, F., Strizhov, N., Jásik, J., Schmelzer, E., Koncz, C., & Szabados, L. (2008). Duplicated P5CS genes of *Arabidopsis* play distinct roles in stress regulation and developmental control of proline biosynthesis. *Plant Journal*, 53(1), 11–28. <https://doi.org/10.1111/j.1365-313X.2007.03318.x>
- Tani, C., & Sasakawa, H. (2006). Proline accumulates in *Casuarina equisetifolia* seedlings under salt stress. *Soil Science and Plant Nutrition*, 52(1), 21–25.
<https://doi.org/10.1111/j.1747-0765.2006.00005.x>
- Teixeira, F. K., Menezes-Benavente, L., Galvão, V. C., Margis, R., & Margis-Pinheiro, M. (2006). Rice ascorbate peroxidase gene family encodes functionally diverse isoforms localized in different subcellular compartments. *Planta*, 224(2), 300–314. <https://doi.org/10.1007/s00425-005-0214-8>
- Tiwari, J. K., Munshi, A. D., Kumar, R., Pandey, R. N., Arora, A., Bhat, J. S., & Sureja, A. K. (2010). Effect of salt stress on cucumber: Na⁺-K⁺ ratio, osmolyte concentration, phenols and chlorophyll content. *Acta Physiologiae Plantarum*, 32(1), 103–114. <https://doi.org/10.1007/s11738-009-0385-1>
- Tyagi, S., Shumayla, Madhu, Singh, K., & Upadhyay, S. K. (2021). Molecular characterization revealed the role of catalases under abiotic and arsenic stress in bread wheat (*Triticum aestivum* L.). *Journal of Hazardous Materials*, 403(April 2020), 123585. <https://doi.org/10.1016/j.jhazmat.2020.123585>

- Verbruggen, N., & Hermans, C. (2008). Proline accumulation in plants: A review. *Amino Acids*, 35(4), 753–759. <https://doi.org/10.1007/s00726-008-0061-6>
- Vighi, I. L., Benitez, L. C., Amaral, M. N., Moraes, G. P., Auler, P. A., Rodrigues, G. S., Deuner, S., Maia, L. C., & Braga, E. J. B. (2017). Functional characterization of the antioxidant enzymes in rice plants exposed to salinity stress. *Biologia Plantarum*, 61(3), 540–550. <https://doi.org/10.1007/s10535-017-0727-6>
- Vranova, E., Inze, D., & Breusegem, F. Van. (2002). Signal transduction during oxidative stress. *Journal of Experimental Botany*, 53(June 2002), 1227–1236. <https://doi.org/10.1093/jexbot/53.372.1227>
- Wang, F., Wang, Q., Kwon, S., Kwak, S., & Su, W. (2005). *Enhanced drought tolerance of transgenic rice plants expressing a pea manganese superoxide dismutase*. 162. <https://doi.org/10.1016/j.jplph.2004.09.009>
- Wang, Z. Q., Yuan, Y. Z., Ou, J. Q., Lin, Q. H., & Zhang, C. F. (2007). Glutamine synthetase and glutamate dehydrogenase contribute differentially to proline accumulation in leaves of wheat (*Triticum aestivum*) seedlings exposed to different salinity. *Journal of Plant Physiology*, 164(6), 695–701. <https://doi.org/10.1016/j.jplph.2006.05.001>
- Wani, A. S., Ahmad, A., Hayat, S., & Fariduddin, Q. (2013). Salt-induced modulation in growth, photosynthesis and antioxidant system in two varieties of *Brassica juncea*. *Saudi Journal of Biological Sciences*, 20(2), 183–193. <https://doi.org/10.1016/j.sjbs.2013.01.006>
- Yan, Z., Guo, S., Shu, S., Sun, J., & Tezuka, T. (2011). Effects of proline on photosynthesis, root reactive oxygen species (ROS) metabolism in two melon cultivars (*Cucumis melo* L.) under NaCl stress. *African Journal of Biotechnology*, 10(80), 18381–18390. <https://doi.org/10.5897/AJB11.1073>
- Youssef, N. Ben, Al-Moaikal, R. M. S., Al-Hawas, G. H. S., & Alrammah, F. M. (2020). Up-modulation of membrane lipid composition and functionality by seed priming under salinity in the hasawi rice variety. *Journal of Applied Botany and Food Quality*, 93, 159–167. <https://doi.org/10.5073/JABFQ.2020.093.020>
- Zhang, L., & Becker, D. F. (2015). Connecting proline metabolism and signaling pathways in plant senescence. *Frontiers in Plant Science*, 6(JULY), 1–8. <https://doi.org/10.3389/fpls.2015.00552>
- Zhang, L. H., Ye, G. F., Lin, Y. M., Zhou, H. C., & Zeng, Q. (2009). Seasonal changes in tannin and nitrogen contents of *Casuarina equisetifolia* branchlets. *Journal of Zhejiang University: Science B*, 10(2), 103–111. <https://doi.org/10.1631/jzus.B0820217>
- Zhao, M., Lin, Y., & Chen, H. (2020). Improving nutritional quality of rice for human health. *Theoretical and Applied Genetics*, 133(5), 1397–1413.

<https://doi.org/10.1007/s00122-019-03530-x>