

## DAFTAR PUSTAKA

- Aenugu, H. P. R., Sathis Kumar, D., Srisudharson, Parthiban, N., Ghosh, S. S., & Banji, D. (2011). Near infra red spectroscopy- An overview. *International Journal of ChemTech Research*, 3(2), 825–836.
- Afandiyah, G. (2019). Pengaruh metode ekstraksi terhadap viabilitas dan vigor benih cabai rawit (*Capsicum frutescents L.*) *Skripsi*. Fakultas Pertanian. Universitas Brawijaya, Malang.
- Afifah, N., Widajati, E., & Palupi, E. R. (2020). Development of tetrazolium test as vigor analysis method of true seed of shallot. *Jurnal Hortikultura Indonesia*, 11(2), 120–130. <https://doi.org/10.29244/jhi.11.2.120-130>
- Ahmad, U., & Sabihah, . (2018). Prediction of ripeness parameters of melon fruit using Near Infra-red Spectroscopy. *Jurnal Ilmu Pertanian Indonesia*, 23(3), 183–189. <https://doi.org/10.18343/jipi.23.3.183>
- Amanah, H. Z., Joshi, R., Masithoh, R. E., Choung, M. G., Kim, K. H., Kim, G., & Cho, B. K. (2020). Nondestructive measurement of anthocyanin in intact soybean seed using Fourier Transform Near-Infrared (FT-NIR) and Fourier Transform Infrared (FT-IR) spectroscopy. *Infrared Physics and Technology*, 111, 103477. <https://doi.org/10.1016/j.infrared.2020.103477>
- Amanah, H. Z., Rahayoe, S., Harmayani, E., Adhitama, R., & Hernanda, P. (2024). Construction of a sustainable model to predict the moisture content of porang powder (*Amorphophallus oncophyllus*) based on pointed- scan visible near-infrared spectroscopy. *Open Agriculture*, 9(1), 1-12. <https://doi.org/10.1515/opag-2022-0268>
- Ambrose, A., Lohumi, S., Lee, W. H., & Cho, B. K. (2016). Comparative nondestructive measurement of corn seed viability using Fourier transform near-infrared (FT-NIR) and Raman spectroscopy. *Sensors and Actuators, B: Chemical*, 224, 500–506. <https://doi.org/10.1016/j.snb.2015.10.082>
- Aminah, A., & Siregar, N. (2019). Pengaruh waktu pengunduhan dan warna kulit buah terhadap daya berkecambah dan pertumbuhan bibit mindi (*Melia azedarach Linn*). *Jurnal Perbenihan Tanaman Hutan*, 7(1), 21-30.
- Badan Pangan Nasional. (2023). Konsumsi cabai per kapita Indonesia. BAPANAS RI.
- Badan Pusat Statistik. (2022). *Distribusi perdagangan komoditas cabai merah indonesia*. BPS RI.
- Badan Pusat Statistik. (2022). *Statistik hortikultura 2022*. BPS RI.
- Belo, S. M., & Suwarno, F. C. (2012). Penurunan viabilitas benih padi (*Oryza sativa L.*) melalui beberapa metode pengusangan cepat. *J. Agron. Indonesia*, 40(1), 29–35. <https://doi.org/10.24831/jai.v40i1.14948>
- Daneshvar, A., Tigabu, M., Karimidoost, A., & Odén, P. C. (2015). Single seed near infrared spectroscopy discriminates viable and non-viable seeds of *Juniperus polycarpus*. *Silva Fennica*, 49(5), 1–14. <https://doi.org/10.14214/sf.1334>
- Delouche, J. C. (1971). Determinants of Seed Quality. *Proceedings of the Short Course for Seedsmen*. <https://scholarsjunction.msstate.edu/seedsman-short-course/255/>

- Dewi, N. N. T. K., Nyana, I. D. N., & Raka, I. N. (2018). Pengaruh rhizobakteria terhadap hasil dan mutu benih kacang tanah (*Arachis hypogaea* L.). *E-Jurnal Agroekoteknologi Tropika*, 7(4), 593–603. <https://ojs.unud.ac.id/index.php/JAT>
- Dharmawan, A., Masithoh, R. E., & Amanah, H. Z. (2023). Development of PCA-MLP model based on visible and shortwave near infrared spectroscopy for authenticating arabica coffee origins. *Foods*, 12(11). <https://doi.org/10.3390/foods12112112>
- Diansari, P., & Rukmana, D. (2023). *Examining the Economic Impact of Cayenne Pepper in Traditional Markets of Makassar City : A Price Analysis Study*. 6(01), 34–41.
- El Orche, A., Bouatia, M., & Mbarki, M. (2020). Rapid analytical method to characterize the freshness of olive oils using fluorescence spectroscopy and chemometric algorithms. *Journal of Analytical Methods in Chemistry*, 2020. <https://doi.org/10.1155/2020/8860161>
- Ellis, R. H., Hong, T. D., & Roberts, E. H. (1991). Seed moisture content, storage, viability and vigour. *Seed Science Research*, 1(4), 275–279. <https://doi.org/10.1017/S0960258500001008>
- Environment Directorate. (2006). *Consensus Document on the Biology of the Capsicum annum Complex (Chili peppers, Hot peppers and Sweet peppers)*. Organisation for Economic Co-operation and Development, 36.
- Fitri, Z. E. Nuhanatika, U., Madjid, A., & Imron, A. M. N. (2020). Penentuan tingkat kematangan cabe rawit (*Capsicum frutescens* L.) berdasarkan Gray Level Co-Occurrence Matrix. *Jurnal Teknologi Informasi dan Terapan (J-TIT)*, 7(1), 1–5. <https://doi.org/10.25047/jtit.v7i1.121>
- Foroozani, B., Bagherpour, H., Caporaso, N., & Zaboli, K. (2022). Classification of wheat varieties by PLS-DA and LDA models and investigation of the spatial distribution of protein content using NIR spectroscopy. *CIGR Journal*, 24(2), 184–193.
- Golic, M., Walsh, K., & Lawson, P. (2003). Short-wavelength near-infrared spectra of sucrose, glucose, and fructose with respect to sugar concentration and temperature. *Applied Spectroscopy*, 57(2), 139–145. <https://doi.org/10.1366/000370203321535033>
- Irawan, M., Yusmanizar, Y., & Munawar, A. A. (2022). Kalibrasi dan Validasi Model Spektrum Inframerah Biji Kopi untuk Pengujian Kadar Air (Moisture Content) dan Asam Klorogenat (Cholorogenic Acid) Biji Kopi Secara Non-Destruktif. *Jurnal Ilmiah Mahasiswa Pertanian*, 7(4), 838–845. <https://doi.org/10.17969/jimfp.v7i4.22229>
- Isik, H., Tasdemir, S., Taspinar, Y. S., Kursun, R., Cinar, I., Yasar, A., Yasin, E. T., & Koklu, M. (2024). Maize seeds forecasting with hybrid directional and bi-directional long short-term memory models. *Food Science and Nutrition*, 12(2), 786–803. <https://doi.org/10.1002/fsn3.3783>
- Junaidy, R. B., & Shahrudin, S. (2022). Germinability and Seedling Growth Performance of Chilli (*Capsicum annum*) Seeds in Response to Different Gibberellic Acid Concentrations Pre-Treatment. *AgroTech Food Science, Technology and Environment*, 1(1), 10–16.

- <https://doi.org/10.53797/agrotech.v1i1.2.2022>
- Kathuria, D., Hamid, Chavan, P., Jaiswal, A. K., Thakur, A., & Dhiman, A. K. (2024). A Comprehensive Review on Sprouted Seeds Bioactives, the Impact of Novel Processing Techniques and Health Benefits. *Food Reviews International*, 40(1), 370–398. <https://doi.org/10.1080/87559129.2023.2169453>
- Kementerian Pertanian. (2023). *Laporan kinerja Direktorat Jenderal Perkebunan Triwulan I*. Direktorat Jenderal Perkebunan.
- Kerr, A., Rafuse, H., Sparkes, G., Hinchey, J., & Sandeman, H. (2011). Visible/Infrared Spectroscopy (Virs) As a Research Tool in Economic Geology: Background and Pilot Studies From Newfoundland and Labrador. *Current Research*, March 2011, 145–166. [www.ausspec.com](http://www.ausspec.com)
- Khuriyati, N., Falah, M. A. F., Ushada, M., Kristiawan, B., & Wicaksono, N. A. (2018). *Color as a predictor of chili content*. Proceeding of the 2nd International Conference on Tropical Agriculture (Issue December). <https://doi.org/10.1007/978-3-319-97553-5>
- Kusandriani, Y. & Muharam, A. (2005). *Produksi benih cabai*. Badan Penelitian Tanaman Sayuran.
- Kusumaningrum, D., Lee, H., Lohumi, S., Mo, C., Kim, M. S., & Cho, B. K. (2017). Nondestructive technique for determining the viability of soybean (*Glycine max*) seeds using FT-NIR Spectroscopy. *Journal of the Science of Food and Agriculture*, 98(5), 1734–1741. <https://doi.org/10.1002/jsfa.8646>
- Kusumawardana, A., Pujiasmanto, B., & Pardono, P. (2018). Short communication: tetrazolium test for evaluating viability of capsicum annum seeds. *Nusantara Bioscience*, 10(3), 142–145. <https://doi.org/10.13057/nusbiosci/n100302>
- Lagiman & Supriyanta, B. (2021). *Karakterisasi morfologi dan pemuliaan tanaman cabai*. LPPM UPN Veteran Yogyakarta.
- Lasalvia, M., Capozzi, V., & Perna, G. (2022). *applied sciences A Comparison of PCA-LDA and PLS-DA Techniques for Classification of Vibrational Spectra*.
- Listanti, R., Evi Masithoh, R., Dwi Saputro, A., & Zuhrotul Amanah, H. (2023). Identification of Maturity Stage of Cacao using Visible Near Infrared (Vis-NIR) and Shortwave Near Infrared (SW-NIR) Reflectance Spectroscopy. *BIO Web of Conferences*, 80. <https://doi.org/10.1051/bioconf/20238006003>
- Liu, W., Luo, B., Kang, K., Xia, Y., & Zhang, H. (2024). Non-destructive detection of single corn seed vigor based on visible/near-infrared spatially resolved spectroscopy combined with chemometrics. *Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy*, 312(October 2023), 124089. <https://doi.org/10.1016/j.saa.2024.124089>
- Marbun, K. S., Pramono, E., & Hadi, M. S. (2014). Pengaruh suhu deraan dan lama penderaan pada viabilitas benih tomat (*Lycopersicon Esculentum* Mill.) varietas oval. *Jurnal Agrotek Tropika*, 2(1), 55–60. <https://doi.org/10.23960/jat.v2i1.1930>
- Masithoh, R. E., Pahlawan, M. F. R., & Wati, R. K. (2021). Non-destructive determination of SSC and pH of banana using a modular Vis/NIR spectroscopy: Comparison of Partial Least Square (PLS) and Principle Component Regression (PCR). *IOP Conference Series: Earth and*

- Environmental Science*, 752(1). <https://doi.org/10.1088/1755-1315/752/1/012047>
- Metrohm. (2013). *A guide to near-infrared spectroscopic analysis of industrial manufacturing processes*. Metrohm Ltd.
- Mo, C., Kim, G., Lee, K., Kim, M. S., Cho, B. K., Lim, J., & Kang, S. (2014). Non-destructive quality evaluation of pepper (*Capsicum annuum* L.) seeds using LED-induced hyperspectral reflectance imaging. *Sensors (Switzerland)*, 14(4), 7489–7504. <https://doi.org/10.3390/s140407489>
- Munira, Utami, K., & Nasir, M. (2019). Uji aktivitas antibakteri cabai rawit hijau dan cabai rawit merah (*Capsicum frutescens* L) serta kombinasinya terhadap bakteri *Staphylococcus aureus*. *Jurnal Bioleuser*, 3(1), 13–17. <https://doi.org/10.24815/j.%20bioleuser.v0i0.18815>
- Nadarajan, J., Walters, C., Pritchard, H. W., Ballesteros, D., & Colville, L. (2023). Seed Longevity—The evolution of knowledge and a conceptual framework. *Plants*, 12, 471. <https://doi.org/10.3390/plants12030471>
- Nasr, M., & Farouk, H. (2017). Performance evaluation of agricultural drainage water using modeling and statistical approaches. *The Egyptian Journal of Aquatic Research*, 42(2), 141–148. <https://doi.org/10.1016/j.ejar.2016.04.006>
- Nizaruddin, Suwarno, F. C., Widajati, E., & Qadir, A. (2014). Metode deteriorasi terkontrol untuk pendugaan daya simpan benih kedelai. *Jurnal Agron. Indonesia*, 42(1), 24–31.
- Nurhafidah, Rahmat, A., Karre, A., & Juraeje, H. H. (2021). Uji daya kecambah berbagai jenis varietas jagung (*Zea mays*) dengan menggunakan metode yang berbeda. *Agroplantae*, 10(8), 30–39.
- O'Shea, M. G., Donald, D. A., Simpson, J. M., Keefe, E. C., & Staunton, S. P. (2013). Development of laboratory Near Infrared (NIR) spectroscopy system for sugar mill and refinery applications. *Proc. Int. Soc. Sugar Cane Technology*, 28, 1771–1881. <https://cabidigitallibrary.org>
- Pahlawan, M. F. R., Murti, B. M. A., & Masithoh, R. E. (2022). The potency of Vis/NIR spectroscopy for classification of soybean based of colour. *IOP Conference Series: Earth and Environmental Science*, 1018(1). <https://doi.org/10.1088/1755-1315/1018/1/012015>
- Pahlawan, M. F. R., Wati, R. K., & Masithoh, R. E. (2021). Development of a low-cost modular VIS/NIR spectroscopy for predicting soluble solid content of banana. *IOP Conference Series: Earth and Environmental Science*, 644(1). <https://doi.org/10.1088/1755-1315/644/1/012047>
- Pirutin, S. K., Jia, S., Yusipovich, A. I., Shank, M. A., Parshina, E. Y., & Rubin, A. B. (2023). Vibrational spectroscopy as a tool for bioanalytical and biomonitoring studies. *International Journal of Molecular Sciences*, 24(8). <https://doi.org/10.3390/ijms24086947>
- Pradhan, N., Fan, X., Martini, F., Chen, H., Liu, H., Gao, J., & Goodale, U. M. (2022). Seed viability testing for research and conservation of epiphytic and terrestrial orchids. *Botanical Studies*, 63(1), 1–14. <https://doi.org/10.1186/s40529-022-00333-0>
- Pu, Y. Y., & Sun, D. W. (2015). Vis-NIR hyperspectral imaging in visualizing moisture distribution of mango slices during microwave-vacuum drying. *Food*



- Chemistry*, 188, 271–278. <https://doi.org/10.1016/j.foodchem.2015.04.120>
- Radhikaa, C. (2020). A Study on various modes of vibration in spectroscopy. *Malaya Journal of Matematik*, S(2), 1103–1105. <https://doi.org/10.26637/MJM0S20/0285>
- Rai, A. K., & Vikram, A. (2016). Genetic characterization of tomato germplasm for yield and quality traits through principal component analysis. *Research Journal of Agricultural Sciences*, 8(5), 1171–1174. <https://www.rjas.org>
- Sada, D. Y. (2023). *Analisis perilaku konsumen dalam pembelian cabai merah keriting (Capsicum annum L) di SayurYuk.com Kota Samarinda*. Prosiding Seminar Nasional Pertanian, 3(1), 20–24.
- Sanchez, P. M., Pauli, E. D., Scheel, G. L., Rakocevic, M., Bruns, R. E., & Scarminio, I. S. (2018). Irrigation and light access effects on coffea arabica l. Leaves by FTIR-Chemometric analysis. *Journal of the Brazilian Chemical Society*, 29(1), 168–176. <https://doi.org/10.21577/0103-5053.20170125>
- Santoso, M. B. (2018). Kajian permasalahan usaha tani dan penerapan teknologi budidaya cabai di Kecamatan Bajuin Kabupaten Tanah Laut. *Jurnal AgroSainTa*. 2(2), 249–257.
- Saputri, D. A. S., Pahlawan, M. F. R., Murti, B. M. A., & Masithoh, R. E. (2022). *Vis / NIR spectroscopy for non-destructive method in detecting soybean seeds viability*. *IOP Conference Series: Earth and Environmental Science*, 1038(1). <https://doi.org/10.1088/1755-1315/1038/1/012043>
- Sathish, S., Ahamed, R., Senthil, N., Arulkumar, N., Park, H. S., Kalaiselvi, S., Umarani, R., Raveendran, M., Bhaskaran, M., & Kim, G. S. (2015). Proteomic analysis of ageing in black gram (*Vigna mungo L.*) seeds and its relation to seed viability. *Plant OMICS*, 8(3), 201–211.
- Seo, Y., Ahn, C. K., Lee, H., Park, E., Mo, C., & Cho, B. (2016). Non-destructive sorting techniques for viable pepper (*Capsicum annum L.*) seeds using Fourier Transform Near-Infrared and Raman Spectroscopy. *Journal of Biosystem Engineering*, 41(1), 51–59. <http://dx.doi.org/10.5307/JBE.2016.41.1.051>
- Shrestha, S., Deleuran, L. C., & Gislum, R. (2017). Separation of viable and non-viable tomato (*Solanum lycopersicum L.*) seeds using single seed near-infrared spectroscopy. *Computers and Electronics in Agriculture*, 142, 348–355. <https://doi.org/10.1016/j.compag.2017.09.004>
- Sukrasno, S., & Kusmardiyani, S. (1997). Kandungan kapsaisin dan dihidrokapsaisin pada berbagai buah capsicum. *Jurnal Matematika dan Sains*, 2(1), 28–34.
- Tazi, I. (2023). *Linear Discriminant Analysis (LDA) and Partial Least Square (PLS) of chemometric in mixture of lard and palm oil-based on Ftir-Spectroscopy data*. Proceedings of the 12<sup>th</sup> International Conference on Green Technology, 378–385. <https://doi.org/10.2991/978-94-6463-148-7>
- Wakholi, C., Kandpal, L. M., Lee, H., Bae, H., Park, E., Kim, M. S., Mo, C., Lee, W. H., & Cho, B. K. (2018). Rapid assessment of corn seed viability using short wave infrared line-scan hyperspectral imaging and chemometrics. *Sensors and Actuators, B: Chemical*, 255, 498–507. <https://doi.org/10.1016/j.snb.2017.08.036>

- Wen, B. (2019). Seed germination ecology of Alexandra palm (*Archontophoenix alexandrae*) and its implication on invasiveness. *Scientific Reports*, 9(1), 1–9. <https://doi.org/10.1038/s41598-019-40733-0>
- Westerhuis, J. A., Hoefsloot, H. C. J., Smit, S., Vis, D. J., Smilde, A. K., Velzen, E. J. J., Duijnhoven, J. P. M., & Dorsten, F. A. (2008). Assessment of PLSDA cross validation. *Metabolomics*, 4(1), 81–89. <https://doi.org/10.1007/s11306-007-0099-6>
- Yasmin, J., Ahmed, M. R., Lohumi, S., Wakholi, C., Kim, M. S., & Cho, B. (2019). Classification method for viability screening of naturally aged watermelon seeds using FT-NIR spectroscopy. *Sensors*, 19. <https://doi.org/10.3390/s19051190>
- Zwinkels, J. (2015). Light, electromagnetic spectrum. *Encyclopedia of Color Science and Technology*, 1–8. <https://doi.org/10.1007/978-3-642-27851-8>