

## DAFTAR PUSTAKA

- Abdel-Farid, I. B., Kim, H. K., Choi, Y. H., & Verpoorte, R. (2007). Metabolic Characterization of *Brassica rapa* Leaves by NMR Spectroscopy. *J. Agric. Food Chem*, 55, 7936–7943. <https://doi.org/10.1021/jf071294b>
- Abubakar, R., Suleiman, M., & Wagini, N. H. (2021). Phytochemical and insecticidal properties of some botanical extracts against the lesser grain borer , *Rhyzopertha dominica* F . ( Coleoptera : Bostrichidae). 9(2), 6–13. <https://doi.org/10.22271/j.ento.2021.v9.i2a.8452>
- Adamski, Z., Radtke, K., Kopiczko, A., Chowański, S., Marciniak, P., Szymczak, M., Spochacz, M., Falabella, P., Lelario, F., Scrano, L., & Bufo, S. A. (2016). Ultrastructural and developmental toxicity of potato and tomato leaf extracts to beet armyworm, *Spodoptera exigua* (Lepidoptera: noctuidae). *Microscopy Research and Technique*, 79(10), 948–958. <https://doi.org/10.1002/jemt.22726>
- Adfa, M., Wiradimafan, K., Febri, R., & Angga, P. (2023). Anti-Termite Activity of *Azadirachta excelsa* Seed Kernel and Its Isolated Compound against *Coptotermes curvignathus*. 51(3), 157–172. <https://doi.org/10.5658/WOOD.2023.51.3.157>
- Aflatoon, H., Ebrahimi, M., Ramshini, H., & Akbari, G. (2018). Evaluation of Some Morphological Characteristics of Different Basil Masses ( *Ocimum basilicum* L .) *Gene Technology*. 1–5. <https://doi.org/10.24015/2329-6682.8.148>
- Agustian, A., & Rachman, B. (2013). Penerapan teknologi pengendalian hama terpadu pada komoditas perkebunan rakyat. *Prespektif*, 8(1), 30–41.
- Agustina, N., Purwestri, Y. A., & Nugroho, L. H. (2016). Antioxidant Activity and Histochemical Analysis of *Acalypha indica* L . and *Acalypha wilkesiana* Muell . Arg . *Vegetative and Generative Organs*. 8(10), 1657–1662.
- Ahmad, M., Farid, A., & Saeed, M. (2018). Resistance to new insecticides and their synergism in *Spodoptera exigua* (Lepidoptera: Noctuidae) from Pakistan. *Crop Protection*, 107(December 2017), 79–86. <https://doi.org/10.1016/j.cropro.2017.12.028>
- Al-maawali, S. S., Al-sadi, A. M., Khalifa, S. A., Al-sabahi, J. N., & Velazhahan, R. (2021). The potential of antagonistic yeasts and bacteria from tomato phyllosphere and fructoplane in the control of *Alternaria* fruit rot of tomato. *All Life*, 14(1), 34-48. <https://doi.org/10.1080/26895293.2020.1858975>
- Al-snafi, P. A. E., & Mill, F. (2018). The chemical constituents and pharmacological effects of *Foeniculum vulgare* - A review. *IOSR Journal of Pharmacy*, 8(5), 81–96.

- Aldini, G. M., Wijonarko, A., Witjaksono, De Putter, H., Hengsdijk, H., & Trisyono, Y. A. (2021). Insecticide Resistance in *Spodoptera exigua* (Lepidoptera: Noctuidae) Populations in Shallot Areas of Java, Indonesia. *Journal of Economic Entomology*, 114(6), 2505–2511. <https://doi.org/10.1093/jee/toab183>
- Anlas, C., Ustuner, O., Alkan, F. U., Bakirel, T., Aydogan, M. N., & Erel, S. B. (2017). A comparative study on the antioxidant activities and phenolic contents of different extracts of *Achillea nobilis* subsp. *Sipylea* and *Alcea apterocarpa* (fenzl) boiss, *Endemic plants*. 26(2), 1423–1430
- Anwaar, M. A. (2020). Phytochemical and insecticidal evaluation of agro-waste of *Lagenaria siceraria* (Cucurbitaceae) plant against the cotton leafworm *Spodoptera littoralis* (Lepidoptera: Noctuidae). *Egypt. J. Plant Prot. Res. Inst.*, 3(2), 777–785.
- Ardrey, R. E. (2003). *Liquid chromatography-mass spectrometry: an introduction* (Vol. 2). John Wiley & Sons.
- Arivoli, S., & Tennyson, S. (2012). Antifeedant Activity of Plant Extracts Against *Spodoptera litura* (Fab.) (Lepidoptera: Noctuidae). *American-Eurasian J. Agric. & Environ. Sci*, 12(6), 764–768. <https://doi.org/10.5829/idosi.aejaes.2012.12.06.63178>
- Arulkumar, G., Manisegaran, S., Nalini, R., & Mathialagan, M. (2017). Seasonable abundance of beet armyworm *Spodoptera exigua* (Hubner) infesting Onion with weather factors in Madurai district of Tamil Nadu. *Journal of Entomology and Zoology Studies*, 5(6), 1157–1162.
- Askari-Khorasgani, O., & Pessarakli, M. (2020). Evaluation of cultivation methods and sustainable agricultural practices for improving shallot bulb production—a review. *Journal of Plant Nutrition*, 43(1), 148–163. <https://doi.org/10.1080/01904167.2019.1659329>
- Atkins, L. E. (1960). The Beet Armyworm, *Spodoptera exigua*; an Economic Pest of Citrus in California1. *Journal of Economic Entomology*, 53(4), 616–619. <https://doi.org/10.1093/jee/53.4.616>
- Azmir, J., Zaidul, I. S. M., Rahman, M. M., Sharif, K. M., Mohamed, A., Sahena, F., Jahurul, M. H. A., Ghafoor, K., Norulaini, N. A. N., & Omar, A. K. M. (2013). Techniques for extraction of bioactive compounds from plant materials: A review. *Journal of Food Engineering*, 117(4), 426–436. <https://doi.org/10.1016/j.jfoodeng.2013.01.014>
- Azwanida, N. N. (2015). A review on the extraction methods use in medicinal plants, principle, strength and limitation. *Med Aromat Plants*, 4(196), 2167–0412.
- Badawy, M. E. I., El-Arami, S. A. A., & Abdelgaleil, S. A. M. (2010). Acaricidal and quantitative structure activity relationship of monoterpenes against the

- two-spotted spider mite, *Tetranychus urticae*. *Experimental and Applied Acarology*, 52(3), 261–274. <https://doi.org/10.1007/s10493-010-9363-y>
- Baker, B. P., Green, T. A., & Loker, A. J. (2020). Biological control and integrated pest management in organic and conventional systems. *Biological Control*, 140(September 2019), 104095. <https://doi.org/10.1016/j.biocontrol.2019.104095>
- Barbieri, G., Vallone, S., Orsini, F., Paradiso, R., De Pascale, S., Negre-Zakharov, F., & Maggio, A. (2012). Stomatal density and metabolic determinants mediate salt stress adaptation and water use efficiency in basil (*Ocimum basilicum* L.). *Journal of Plant Physiology*, 169(17), 1737–1746. <https://doi.org/10.1016/j.jplph.2012.07.001>
- Barzalona, M., & Casanova, J. (2008). Chemical variability of the leaf oil of 113 hybrids from *Citrus clementina* (Commun)× *Citrus deliciosa* (Willow Leaf). *Flavour and fragrance journal*, 23(3), 152–163. <https://doi.org/10.1002/ffj>
- Belwal, T., Ezzat, S. M., Rastrelli, L., Bhatt, I. D., Daglia, M., Baldi, A., Devkota, H. P., Orhan, I. E., Patra, J. K., Das, G., Anandharamakrishnan, C., Gomez-Gomez, L., Nabavi, S. F., Nabavi, S. M., & Atanasov, A. G. (2018). A critical analysis of extraction techniques used for botanicals: Trends, priorities, industrial uses and optimization strategies. *TrAC - Trends in Analytical Chemistry*, 100, 82–102. <https://doi.org/10.1016/j.trac.2017.12.018>
- Bharti, S. K., & Roy, R. (2012). Quantitative <sup>1</sup>H NMR spectroscopy. *TrAC - Trends in Analytical Chemistry*, 35, 5–26. <https://doi.org/10.1016/j.trac.2012.02.007>
- Bhavya, M. L., Chandu, A. G. S., & Devi, S. S. (2018). *Ocimum tenuiflorum* oil, a potential insecticide against rice weevil with anti-acetylcholinesterase activity. *Industrial Crops and Products*, 126(August), 434–439. <https://doi.org/10.1016/j.indcrop.2018.10.043>
- Bibiano, C. S., Alves, D. S., Freire, B. C., Vilela Bertolucci, S. K., & Carvalho, G. A. (2022). Toxicity of essential oils and pure compounds of Lamiaceae species against *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and their safety for the nontarget organism *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae). *Crop Protection*, 158(October 2021), 106011. <https://doi.org/10.1016/j.cropro.2022.106011>
- Biru, F. N. (2015). Effect of spacing and nitrogen fertilizer on the yield and yield component of shallot (*Allium ascalonium* L.). *Journal of Agronomy*, 14(4), 220–226. <https://doi.org/10.3923/ja.2015.220.226>
- Bilal, A., Jahan, N., Ahmed, A., Brito, S. N., Habib, S., & Hajra, S. (2012). Phytochemical and pharmacological studies on *Ocimum basilicum* Linn-A review. *International Journal of Current Research and Review*, 4(23), 73–

83.

- Blokhina, O., virolainen, E., & Fagerstedt, F. K. V. (2003). Antioxidants , Oxidative Damage and Oxygen Deprivation Stress : a Review. *Annals of Botany*, 91, 179–194. <https://doi.org/10.1093/aob/mcf118>
- Brito, V. D., Achimon, F., Dambolena, J. S., Pizzolitto, R. P., & Zygadlo, J. A. (2019). Trans-2-hexen-1-ol as a tool for the control of *Fusarium verticillioides* in stored maize grains. *Journal of Stored Products Research*, 82, 123–130. <https://doi.org/10.1016/j.jspr.2019.05.002>
- Brito, V. D., Achimon, F., Dambolena, J. S., Pizzolitto, R. P., & Zygadlo, J. A. (2019). Trans-2-hexen-1-ol as a tool for the control of *Fusarium verticillioides* in stored maize grains. *Journal of Stored Products Research*, 82, 123–130. <https://doi.org/10.1016/j.jspr.2019.05.002>
- Cai, L. (2014). Thin layer chromatography. *Current Protocols in Essential Laboratory Techniques*, 2014(February), 6.3.1-6.3.18. <https://doi.org/10.1002/9780470089941.et0603s08>
- Caldwell, M. M., Robberecht, R., Flint, S. D., & Internal, S. D. (1983). Internal filters: prospects for UV-acclimation in higher plants. *Physiologia Plantarum*, 58(3), 445–450. <https://doi.org/10.1111/j.1399-3054.1983.tb04206.x>
- Campos, E. V. R., Proença, P. L. F., Oliveira, J. L., Pereira, A. E. S., De Moraes Ribeiro, L. N., Fernandes, F. O., Gonçalves, K. C., Polanczyk, R. A., Pasquoto-Stigliani, T., Lima, R., Melville, C. C., Della Vechia, J. F., Andrade, D. J., & Fraceto, L. F. (2018). Carvacrol and linalool co-loaded in  $\beta$ -cyclodextrin-grafted chitosan nanoparticles as sustainable biopesticide aiming pest control. *Scientific Reports*, 8(1), 1–14. <https://doi.org/10.1038/s41598-018-26043-x>
- Cantino, P. D. (1990). The phylogenetic significance of stomata and trichomes in the labiatae and verbenaceae. *Journal of the Arnold Arboretum*, 71( 3), 323-370. <https://doi.org/10.5962/p.184532>
- Castro, M. D. M., & Demarco, D. (2008). Phenolic compounds produced by secretory structures in plants: a brief review. *Natural Product Communications*, 3(8), 1273-1284. <https://doi.org/10.1177/1934578X0800300809>
- Chaaban, S. Ben, Hamdi, S. H., Mahjoubi, K., & Jemâa, J. M. Ben. (2019). Composition and insecticidal activity of essential oil from *Ruta graveolens*, *Mentha pulegium* and *Ocimum basilicum* against *Ectomyelois ceratoniae* Zeller and *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae). *Journal of Plant Diseases and Protection*, 0123456789. <https://doi.org/10.1007/s41348-019-00218-8>
- Chaudhary, A., Bala, K., Thakur, S., Kamboj, R., & Dumra, N. (2018). Plant defenses against herbivorous insects: A Review. *International Journal of*

*Chemical Studies*, 6(5), 681–688.  
<https://www.researchgate.net/publication/327703671>

- Chemat, F., & Boutekedjiret, C. (2015). Extraction // Steam Distillation☆. In *Reference Module in Chemistry, Molecular Sciences and Chemical Engineering* (Issue March). Elsevier Inc. <https://doi.org/10.1016/b978-0-12-409547-2.11557-4>
- Chen, W., & Viljoen, A. M. (2010). Geraniol — A review of a commercially important fragrance material. *South African Journal of Botany*, 76(4), 643–651. <https://doi.org/10.1016/j.sajb.2010.05.008>
- Chwil, M., Nurzyńska, R., Chwil, S., Matraszek, R., & Neugebauerová, J. (2016). Histochemistry and micromorphological diversity of glandular trichomes in *Melissa officinalis* L. leaf epidermis. *Acta Scientiarum Polonorum Hortorum Cultus*, 15(3), 153–172.
- Coll, J. C., & Bowden, B. F. (1986). The application of vacuum liquid chromatography to the separation of terpene mixtures. *Journal of Natural Products*, 49(5), 934–936. <https://doi.org/10.1021/np50047a033>
- Copping, L. G., & Menn, J. J. (2000). Biopesticides: A review of their action, applications and efficacy. *Pest Management Science*, 56(8), 651–676. [https://doi.org/10.1002/1526-4998\(200008\)56:8<651::AID-PS201>3.0.CO;2-U](https://doi.org/10.1002/1526-4998(200008)56:8<651::AID-PS201>3.0.CO;2-U)
- Cruz, G. S., Wanderley-Teixeira, V., Oliveira, J. V., Lopes, F. S. C., Barbosa, D. R. S., Breda, M. O., Dutra, K. A., Guedes, C. A., Navarro, D. M. A. F., & Teixeira, A. A. C. (2016). Sublethal effects of essential oils from *Eucalyptus staigeriana* (Myrtales: Myrtaceae), *Ocimum gratissimum* (Lamiales: Lamiaceae), and *Foeniculum vulgare* (Apiales: Apiaceae) on the biology of *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Journal of Economic Entomology*, 109(2), 660–666. <https://doi.org/10.1093/jee/tow005>
- Curzi, M. J., Zavala, J. A., Spencer, J. L., & Seufferheld, M. J. (2012). Abnormally high digestive enzyme activity and gene expression explain the contemporary evolution of a diabrotica biotype able to feed on soybeans. *Ecology and Evolution*, 2(8), 2005–2017. <https://doi.org/10.1002/ece3.331>
- da Silva Moura, E., D'Antonino Faroni, L. R., Fernandes Heleno, F., Aparecida Zinato Rodrigues, A., Figueiredo Prates, L. H., & Lopes Ribeiro de Queiroz, M. E. (2020). Optimal extraction of *Ocimum basilicum* essential oil by association of ultrasound and hydrodistillation and its potential as a biopesticide against a major stored grains pest. *Molecules*, 25(12), 2781. <https://doi.org/10.3390/molecules25122781>
- Dai, H., Zhang, G., & Zhang, W. (2017). Temperature dependent development parameters and population life table of beet armyworm, *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae). *Arthropods*, 6(4), 117–125. [https://www.researchgate.net/publication/321414212\\_Temperature\\_depen](https://www.researchgate.net/publication/321414212_Temperature_depen)

dent\_development\_parameters\_and\_population\_life\_table\_of\_beet\_armyworm\_Spodoptera\_exigua\_Hubner\_Lepidoptera\_Noctuidae

- Dai, L. Shang, Tian, H. Fei, Hang, Y., Wen, C. Wei, Huang, Y. hao, Wang, B. Feng, Hu, J. Wei, Xu, J. Ping, & Deng, M. Jie. (2020). <sup>1</sup>H NMR-based metabonomic evaluation of the pesticides camptothecin and matrine against larvae of *Spodoptera litura*. *Pest Management Science*, April. <https://doi.org/10.1002/ps.6009>
- Dancewicz, K., Szumny, A., Wawrzńczyk, C., & Gabryś, B. (2020). Repellent and Antifeedant Activities of Citral-Derived Lactones against the Peach Potato Aphid. *International Journal of Molecular Sciences*, 21(21), 1–16. <https://doi.org/10.3390/ijms21218029>
- Day, T. A., & Howells, B. W. (1994). Ultraviolet absorption and epidermal-transmittance spectra in foliage. *Physiologia Plantarum*, 92(2), 207–218. <https://doi.org/10.1111/j.1399-3054.1994.tb05328.x>
- Deewatthanawong, R., Kongchinda, P., Deewatthanawong, P., Pumnuan, J., & Insung, A. (2019). GC-MS analysis and biopesticide properties of different crude extracts of *Annona squamosa* and *Annona muricata*. *International Journal of Agricultural Technology*, 15(6), 859–868.
- Deng, M. J., Lin, X. D., Lin, Q. T., Wen, D. F., Zhang, M. L., Wang, X. Q., Gao, H. C., & Xu, J. P. (2015). A <sup>1</sup>H-NMR based study on hemolymph metabolomics in Eri silkworm after oral administration of 1-deoxynojirimycin. *PLoS ONE*, 10(7), 1–15. <https://doi.org/10.1371/journal.pone.0131696>
- Deschamps, C., & Simon, J. E. (2006). Terpenoid Essential Oil Metabolism in Basil (*Ocimum basilicum* L.) Following Elicitation. *Journal of Essential Oil Research*, 18(6), 618–621. <https://doi.org/10.1080/10412905.2006.9699183>
- Dimetry, N. Z. (2014). Different plant families as bioresource for pesticides. *Advances in plant biopesticides*, 1–20. [https://doi.org/10.1007/978-81-322-2006-0\\_1](https://doi.org/10.1007/978-81-322-2006-0_1)
- Dirar, A. I., Alsaadi, D. H. M., Wada, M., Mohamed, M. A., Watanabe, T., & Devkota, H. P. (2019). Effects of extraction solvents on total phenolic and flavonoid contents and biological activities of extracts from Sudanese medicinal plants. *South African Journal of Botany*, 120, 261–267. <https://doi.org/10.1016/j.sajb.2018.07.003>
- Dmitruk, M., Sulborska, A., Żuraw, B., Stawiarz, E., & Weryszko-Chmielewska, E. (2019). Sites of secretion of bioactive compounds in leaves of *Dracocephalum moldavica* L.: anatomical, histochemical, and essential oil study. *Revista Brasileira de Botanica*, 42(4), 701–715. <https://doi.org/10.1007/s40415-019-00559-6>

- Draz, K. A., Tabikha, R. M., Eldosouky, M. I., Darwish, A. A., & Abdelnasser, M. (2022). Biototoxicity of essential oils and their nano-emulsions against the coleopteran stored product insect pests *Sitophilus oryzae* L. and *Tribolium castaneum* herbst. *International Journal of Pest Management*, 0(0), 1–15. <https://doi.org/10.1080/09670874.2022.2036862>
- Duarte, J. L., Amado, J. R. R., Oliveira, A. E. M. F. M., Cruz, R. A. S., Ferreira, A. M., Souto, R. N. P., Falcão, D. Q., Carvalho, J. C. T., & Fernandes, C. P. (2015). Evaluation of larvicidal activity of a nanoemulsion of *Rosmarinus officinalis* essential oil. *Revista Brasileira de Farmacognosia*, 25(2), 189–192. <https://doi.org/10.1016/j.bjp.2015.02.010>
- Dutra, K. de A., de Oliveira, J. V., Navarro, D. M. do A. F., Barbosa, D. R. e. S., & Santos, J. P. O. (2016). Control of *Callosobruchus maculatus* (FABR.) (Coleoptera: Chrysomelidae: Bruchinae) in *Vigna unguiculata* (L.) WALP. with essential oils from four *Citrus* spp. plants. *Journal of Stored Products Research*, 68, 25–32. <https://doi.org/10.1016/j.jspr.2016.04.001>
- El-Wakeil, N. E. (2013). Botanical Pesticides and Their Mode of Action. *Gesunde Pflanzen*, 65(4), 125–149. <https://doi.org/10.1007/s10343-013-0308-3>
- El Makarem, H. A., El Kholy, S. E., Abdel-Latif, A., & Seif, A. I. (2015). Physiological and biochemical effects of some essential oils on the granary weevil, *Sitophilus granarius* (L.) (Coleoptera: curculionidae). *Journal of Experimental Biology*, 11(2), 117–123.
- Elumalai, K., Krishnappa, K., Anandan, A., Govindarajan, M., & Mathivanan, T. (2010). Larvicidal and ovicidal activity of seven essential oil against lepidopteran pest *S. litura* (Lepidoptera: Noctuidae). *Int J Recent Sci Res*, 1, 8–14.
- Fatima, S., Farooqi, A. H. A., Ansari, S. R., & Sharma, S. (2011). Effect of water stress on growth and essential oil metabolism in *Cymbopogon martinii* (palmarosa) cultivars. *Journal of Essential Oil Research*, 11(4), 491–496. <https://doi.org/10.1080/10412905.1999.9701193>
- Ferreira, F. T. R., Vendramim, J. D., & Forim, M. R. (2012). Bioatividade de nanoformulações de nim sobre a traça-do-tomateiro. *Ciência Rural*, 42(8), 1347–1353. <https://doi.org/10.1590/s0103-84782012000800003>
- Figueiredo, A. C., Barroso, J. G., Pedro, L. G., & Scheffer, J. J. (2008). Factors affecting secondary metabolite production in plants: volatile components and essential oils. *Flavour and Fragrance Journal*, 23(4), 213–226. <https://doi.org/10.1002/ffj.1875>
- Fitriana, N., & Susandarini, R. (2019). Morphology and taxonomic relationships of shallot (*Allium cepa* L. group aggregatum) cultivars from Indonesia. *Biodiversitas*, 20(10), 2809–2814. <https://doi.org/10.13057/biodiv/d201005>
- Fürstenberg-Hägg, J., Zagrobelny, M., & Bak, S. (2013). Plant defense against

- insect herbivores. In *International Journal of Molecular Sciences* (Vol. 14, Issue 5). <https://doi.org/10.3390/ijms140510242>
- Gang, D. R., Simon, J., Lewinsohn, E., & Pichersky, E. (2002). Peltate glandular trichomes of *Ocimum basilicum* L. (sweet basil) contain high levels of enzymes involved in the biosynthesis of phenylpropenes. *Journal of Herbs, Spices and Medicinal Plants*, 9(2–3), 189–195. [https://doi.org/10.1300/J044v09n02\\_27](https://doi.org/10.1300/J044v09n02_27)
- Gaspar-Pintiliescu, A., Mihai, E., Ciucan, T., Popescu, A. F., Luntraru, C., Tomescu, J., & Craciunescu, O. (2022). Antioxidant and acetylcholinesterase inhibition capacity of hyrosols from lamiaceae plants for biopesticide use: role of phenolics. *International Journal of Food Properties*, 25(1), 996–1008. <https://doi.org/10.1080/10942912.2022.2071289>
- Gökçe, A., Whalon, M. E., & Yanar, Y. (2007). Contact and residual toxicities of 30 plant extracts to Colorado potato beetle larvae. *Archives of Phytopathology and Plant Protection*, 40(6), 441–450. <https://doi.org/10.1080/03235400600628013>
- Gonzales, W. L., Negritto, A., Sua, L. H., & Gonza, W. L. (2008). Induction of glandular and non-glandular trichomes by damage in leaves of *Madia sativa* under contrasting water regimes. *Acta oecologica*, 33(1), 128–132. <https://doi.org/10.1016/j.actao.2007.10.004>
- Govindaraju, V., Young, K., & Maudsley, A. A. (2000). Proton NMR chemical shifts and coupling constants for brain metabolites. *NMR in Biomedicine: An International Journal Devoted to the Development and Application of Magnetic Resonance In Vivo*, 13(3), 129–153. [https://doi.org/10.1002/1099-1492\(200005\)13:3<129::AID-NBM619>3.0.CO;2-V](https://doi.org/10.1002/1099-1492(200005)13:3<129::AID-NBM619>3.0.CO;2-V)
- Govindharaj, G., Jena, M., Annamalai, M., Basana-gowda, G., Muthiah, C., Patil, N., Chandra, P., & Adak, T. (2022). Industrial Crops & Products Toxicity of water pepper , *Persicaria hydropiper* ( L . ) extracts against *Nilaparvata lugens* ( Stål ) and non-targeted effect on earthworm. *Industrial Crops & Products*, 187(PA), 115309. <https://doi.org/10.1016/j.indcrop.2022.115309>
- Greenacre, M., Groenen, P. J. F., Hastie, T., Enza, A. I. D., Markos, A., & Tuzhilina, E. (2022). Principal component analysis. *Nature Reviews Methods Primers*, 2(1), 100. <https://doi.org/10.1038/s43586-022-00184-w>
- Gul, S., Ahmad, M., Zafar, M., Bahadur, S., Sultana, S., Ashfaq, S., Ullah, F., Kilic, O., Hassan, F. ul, & Siddiq, Z. (2019). Foliar epidermal anatomy of Lamiaceae with special emphasis on their trichomes diversity using scanning electron microscopy. *Microscopy Research and Technique*, 82(3), 206–223. <https://doi.org/10.1002/jemt.23157>
- Hage, D. S. (2018). Chromatography. In *Principles and Applications of Clinical Mass Spectrometry: Small Molecules, Peptides, and Pathogens*.

<https://doi.org/10.1016/B978-0-12-816063-3.00001-3>

Halliru, M., & Suleiman, M. (2022). Potentiality of some botanical extracts as biopesticides against the maize weevil, *Sitophilus zeamais* Motsch.(Coleoptera: Curculionidae). *J. Entomol. Zool. Stud*, 10, 34-41. <https://doi.org/10.22271/j.ento.2022.v10.i1a.8920>

Handayani, P. A., & Nurcahyanti, H. (2014). Ekstraksi Minyak Atsiri Daun Zodia (*Evodia Suaveolens*) dengan Metode Maserasi dan Distilasi Air. *Jurnal Bahan Alam Terbarukan*, 3(1), 1-7.

Hussain, A., Rizwan-Ul-Haq, M., AlJabr, A. M., & Al-Ayedh, H. (2019). Lethality of sesquiterpenes reprogramming red palm weevil detoxification mechanism for natural novel biopesticide development. *Molecules*, 24(9), 1–13. <https://doi.org/10.3390/molecules24091648>

Iason, G. (2005). The role of plant secondary metabolites in mammalian herbivory: ecological perspectives. *Proceedings of the Nutrition Society*, 64(1), 123–131. <https://doi.org/10.1079/pns2004415>

Ibrahim, H. A. H. (Ed.). (2019). *Fractionation*. IntechOpen.

Ibrahim, N., Abbas, H., Sayed, N. S. El, & Gad, H. A. (2022). *Rosmarinus officinalis* L . hexane extract : phytochemical analysis , nanoencapsulation , and in silico , in vitro , and in vivo anti - photoaging potential evaluation. *Scientific Reports*, 1–20. <https://doi.org/10.1038/s41598-022-16592-7>

Iijima, Y., Gang, D. R., Fridman, E., Lewinsohn, E., & Pichersky, E. (2004). Characterization of Geraniol Synthase from the Peltate Glands of Sweet Basil. *Plant Physiology*, 134(1), 370–379. <https://doi.org/10.1104/pp.103.032946>

Ikawati, S., Himawan, T., Abadi, A. L., Tarno, H., & Fajarudin, A. (2022). In Silico Study of Eugenol and Trans-Caryophyllene also Clove Oil Fumigant Toxicity on *Tribolium castaneum*. *Journal of Tropical Life Science*, 12(3), 339–349. <https://doi.org/10.11594/jtls.12.03.07>

Ilmiah, H. H., Nuringtyas, T. R., & Nugroho, L. H. (2018). Accumulation of potential photo-protective compound groups in mangrove (*Sonneratia caseolaris* (L.) engler.) leaves. *Pharmacognosy Journal*, 10(3), 576–580. <https://doi.org/10.5530/pj.2018.3.94>

Indriyanti, D. R., Fauzi, B. A., & Maretta, Y. A. (2017). The pathogenicity of entomopathogenic nematodes against *Spodoptera exigua*. *ARPN Journal of Engineering and Applied Sciences*, 12(24), 7161–7164.

ITIS, (2020a). *Allium cepa* L.

[https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=42720#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=42720#null)

ITIS, (2020b). *Spodoptera exigua* Hübner.

[https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=117471#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=117471#null)

ITIS, (2020c). *Ocimum* spp.

[https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=32626#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=32626#null)

Jaleel, C. A., Manivannan, P., Wahid, A., Farooq, M., Al-Juburi, H., Jasim, Somasundaram, R., & Panneerselvam, R. (2009). Drought Stress in Plants : A Review on Morphological Characteristics and Pigments Composition. *International Journal Of Agriculture & Biology*, 11(1), 100–105.

Jankowska, M., Rogalska, J., Wyszowska, J., & Stankiewicz, M. (2018). Molecular targets for components of essential oils in the insect nervous system—a review. *Molecules*, 23(1). <https://doi.org/10.3390/molecules23010034>

Jaramillo-Colorado, B. E., Pino-Benitez, N., & González-Coloma, A. (2019). Volatile composition and biocidal (antifeedant and phytotoxic) activity of the essential oils of four Piperaceae species from Choco-Colombia. *Industrial Crops and Products*, 138(January), 111463. <https://doi.org/10.1016/j.indcrop.2019.06.026>

John, G., Parsons, I., Lees, R. S., Balaska, S., & Vontas, J. (2022). A practical insecticide resistance monitoring bioassay for orally ingested dinotefuran in Anopheles malaria vectors. *Insects*, 13(4), 311. <https://doi.org/10.3390/insects13040311>

Jönsson, M., & Anderson, P. (1999). Electrophysiological response to herbivore-induced host plant volatiles in the moth *Spodoptera littoralis*. *Physiological Entomology*, 24(4), 377–385. <https://doi.org/10.1046/j.1365-3032.1999.00154.x>

Juárez, Z. N., Fortuna, A. M., Sánchez-Arreola, E., López-Olguin, J. F., Bach, H., & Hernández, L. R. (2014). Antifeedant and phagostimulant activity of extracts and pure compounds from *Hymenoxys robusta* on *Spodoptera exigua* (Lepidoptera: Noctuidae) larvae. *Natural Product Communications*, 9(7), 895–898. <https://doi.org/10.1177/1934578x1400900703>

Junhirun, P., Pluempanupat, W., Yooboon, T., Ruttanaphan, T., Koul, O., & Bullangpoti, V. (2018). The Study of Isolated Alkane Compounds and Crude Extracts from *Sphagneticola trilobata* (Asterales: Asteraceae) as a Candidate Botanical Insecticide for Lepidopteran Larvae. *Journal of Economic Entomology*, 111(6), 2699–2705. <https://doi.org/10.1093/jee/toy246>

Kaleeswaran, G., Firake, D. M., Sanjukta, R., Behere, G. T., & Ngachan, S. V. (2018). Bamboo-Leaf Prickly Ash extract: A potential bio-pesticide against oriental leaf worm, *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae). *Journal of Environmental Management*, 208, 46–55.

<https://doi.org/10.1016/j.jenvman.2017.12.017>

- Kamaraj, C., Rahuman, A. A., & Bagavan, A. (2008). Screening for antifeedant and larvicidal activity of plant extracts against *Helicoverpa armigera* (Hübner), *Sylepta derogata* (F.) and *Anopheles stephensi* (Liston). *Parasitology Research*, 103(6), 1361–1368. <https://doi.org/10.1007/s00436-008-1142-x>
- Kanehisa, M., & Goto, S. (2000). KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research*, 28(1), 27–30. <https://doi.org/10.3892/ol.2020.11439>
- Kariyat, R. R., Smith, J. D., Stephenson, A. G., Moraes, C. M. De, & Mescher, M. C. (2017). Non-glandular trichomes of *Solanum carolinense* deter feeding by *Manduca sexta* caterpillars and cause damage to the gut peritrophic matrix. *Proceedings of the Royal Society B: Biological Sciences*, 284(1849), 20162323. <https://doi.org/10.1098/rspb.2016.2323>
- Khair-ul-Bariyah, S., Ahmed, D., & Aujla, M. I. (2012). Comparative Analysis of *Ocimum basilicum* and *Ocimum sanctum*: Extraction Techniques and Urease and alpha-Amylase inhibition Department of Chemistry, Faculty of Organic Chemistry, Forman Christian College, Lahore. Department of Chemistry, Faculty of B. *Pakistan Journal of Chemistry*, 2(3), 134–141. <https://doi.org/10.15228/2012.v02.i03.p06>
- Khalid, K. A. (2006). Influence of water stress on growth, essential oil, and chemical composition of herbs (*Ocimum* sp.). *International Agrophysics*, 20(4), 289–296.
- Kiferle, C., Ascrizzi, R., Martinelli, M., Gonzali, S., Mariotti, L., Pistelli, L., Flamini, G., & Perata, P. (2020). Erratum: Effect of Iodine treatments on *Ocimum basilicum* L.: Biofortification, phenolics production and essential oil composition (PLoS ONE (2019) 14:12 (e0226559) DOI: 10.1371/journal.pone.0226559). *PLoS ONE*, 15(2), 229016. <https://doi.org/10.1371/journal.pone.0229016>
- Kim, H. J., Chen, F., Wang, X., & Rajapakse, N. C. (2005). Effect of chitosan on the biological properties of sweet basil (*Ocimum basilicum* L.). *Journal of Agricultural and Food Chemistry*, 53(9), 3696–3701. <https://doi.org/10.1021/jf0480804>
- Kim, Y., & Hong, Y. (2015). Regulation of hemolymph trehalose level by an insulin-like peptide through diel feeding rhythm of the beet armyworm, *Spodoptera exigua*. *Peptides*, 68, 91–98. <https://doi.org/10.1016/j.peptides.2015.02.003>
- Klowden, M. J. (2013). *Physiological systems in insects*. Academic press.
- Kostić, M., Popović, Z., Brkić, D., Milanović, S., Sivčev, I., & Stanković, S. (2008). Larvicidal and antifeedant activity of some plant-derived compounds to *Lymantria dispar* L. (Lepidoptera: Limantriidae). *Bioresource Technology*,

99(16), 7897–7901. <https://doi.org/10.1016/j.biortech.2008.02.010>

- Kostyukovsky, M., Rafaeli, A., Gileadi, C., Demchenko, N., & Shaaya, E. (2002). Activation of octopaminergic receptors by essential oil constituents isolated from aromatic plants: Possible mode of action against insect pests. *Pest Management Science*, 58(11), 1101–1106. <https://doi.org/10.1002/ps.548>
- Koul. (2008). Essential Oils as Green Pesticides: Potential and Constraints. *Biopestic. Int.*, 4(1), 63–84. <https://doi.org/10.1303/aez.32.437>
- Krauss, P., Markstadter, C., & Riederer, M. (1997). Attenuation of UV radiation by plant cuticles from woody species. *Plant, Cell and Environment*, 20, 1079–1085. <https://doi.org/10.1111/j.1365-3040.1997.tb00684.x>
- Krzysko-Lupicka, T., Walkowiak, W., & Bialon, M. (2019). Comparison of the Fungistatic Activity of Selected Essential Oils Relative to *Fusarium graminearum* Isolates. *Molecules*, 23(311), 1–14. <https://doi.org/10.3390/molecules24020311>
- Kuster, V. C., & Vale, F. H. A. (2016). Leaf histochemistry analysis of four medicinal species from Cerrado. *Revista Brasileira de Farmacognosia*, 26(6), 673–678. <https://doi.org/10.1016/j.bjp.2016.05.015>
- Lachowicz, K. J., Jones, G. P., Briggs, D. R., Bienvenu, F. E., Palmer, M. V, Mishra, V., & Hunter, M. M. (1997). Characteristics of plants and plant extracts from five varieties of basil (*Ocimum basilicum* L.) grown in Australia. *Journal of Agricultural and Food Chemistry*, 45(7), 2660–2665. <https://doi.org/10.1021/jf960791h>
- Lade, B. D., Patil, A. S., Paikrao, H. M., Kale, A. S., & Hire, K. K. (2014). A comprehensive working, principles and applications of thin layer chromatography. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 5(4), 486–503.
- Lessire, R., & Stumpf, P. K. (1983). Nature of the fatty acid synthetase systems in parenchymal and epidermal cells of *Allium porrum* L. leaves. *Plant physiology*, 73(3), 614–618. <https://doi.org/10.1104/pp.73.3.614>
- Lewinsohn, E., Dudai, N., Tadmor, K., Katzir, I., Ravid, U., Putievsky, E., & Joels, D. M. (1998). Histochemical Localization of Citral Accumulation in Lemongrass Leaves (*Cymbopogon citratus* (DC.) Stapf., Poaceae). *Annals of Botany Q*, 81, 35–39. <https://doi.org/10.1006/anbo.1997.0525>
- Li, P., Hu, J. wei, Wen, C. wei, Hang, Y., Zhou, Z. hua, Xie, M., Lv, J. cheng, Wang, C. meng, Huang, Y. H., Xu, J. ping, & Deng, M. jie. (2019). Sanguinarine caused larval lethality and growth inhibition by suppressing energy metabolism in silkworms, *Bombyx mori*. *Pesticide Biochemistry and Physiology*, 160(July), 154–162. <https://doi.org/10.1016/j.pestbp.2019.08.004>
- Li, Y., Kong, D., & Wu, H. (2013). Analysis and evaluation of essential oil

- components of cinnamon barks using GC – MS and FTIR spectroscopy. *Industrial Crops & Products*, 41, 269–278. <https://doi.org/10.1016/j.indcrop.2012.04.056>
- López-Gresa, M. P., Maltese, F., Bellés, J. M., Conejero, V., Kim, H. K., Choi, Y. H., & Verpoorte, R. (2010). Metabolic response of tomato leaves upon different plant–pathogen interactions. *Phytochemical Analysis: An International Journal of Plant Chemical and Biochemical Techniques*, 21(1), 89–94. <https://doi.org/10.1002/pca.1179>
- López, M. D., Jordán, M. J., & Pascual-Villalobos, M. J. (2008). Toxic compounds in essential oils of coriander, caraway and basil active against stored rice pests. *Journal of Stored Products Research*, 44(3), 273–278. <https://doi.org/10.1016/j.jspr.2008.02.005>
- Luthra, R., Srivastava, A. K., & Ganjewala, D. (2017). Histochemical localisation of citral accumulating cite in lemongrass. *Asian Journal of Plant Science*, 6(2), 419–422. <https://doi.org/10.3923/ajps.2007.419.422>
- Makenzi, A. M. M., Manguro, L. O. A., Owuor, P. O., & Opiyo, S. A. (2019). Chemical Constituents of Tobacco. *Bull. Chem. Soc. Ethiop*, 33(3), 527–539. <https://doi.org/10.1271/nogekagaku1924.24.485>
- Mandoulakani, B. A., Eyvazpour, E., & Ghadimzadeh, M. (2017). Phytochemistry The effect of drought stress on the expression of key genes involved in the biosynthesis of phenylpropanoids and essential oil components in basil (*Ocimum basilicum* L.). *Phytochemistry*, 139, 1–7. <https://doi.org/10.1016/j.phytochem.2017.03.006>
- Manikanta, penumudi, & Dokuparthi, sudheer kumar. (2014). A Review On Role Of *Azadirachta indica* A. Juss As A Biopesticide. *International Journal of Universal Pharmacy and Bio Sciences*, 3(2), 10.
- Marotti, M., Piccaglia, R., & Giovanelli, E. (1996). Differences in Essential Oil Composition of Basil (*Ocimum basilicum* L.) Italian Cultivars Related to Morphological Characteristics. *Journal of Agricultural and Food Chemistry*, 44(12), 3926–3929. <https://doi.org/10.1021/jf9601067>
- Matsuda, H., Yamada, T., Yoshida, M., & Nishimura, T. (2015). Flies without trehalose. *Journal of Biological Chemistry*, 290(2), 1244–1255. <https://doi.org/10.1074/jbc.M114.619411>
- Matsumura, F. (2009). See Also the Following Articles. *Encyclopedia of Insects*, all, 502–505. <https://doi.org/10.1016/B978-0-12-374144-8.00143-0>
- Martin, C., Bhatt, K., & Baumann, K. (2001). Shaping in plant cells. *Current Opinion in Plant Biology*, 4(6), 540–549. [https://doi.org/10.1016/S1369-5266\(00\)00213-2](https://doi.org/10.1016/S1369-5266(00)00213-2)
- Mead, D. (2014). Basils (*Ocimum* spp.) in Indonesia. *Sulang Lang Data Work Pap*, 1, 1–10.

- Mead, H. M. I. (2018). Composition and larvicidal action of *Ocimum basilicum* L. essential oil against *Spodoptera littoralis* (Boisd.). *Journal of Plant Protection and Pathology*, 9(2), 139-143. <https://doi.org/10.21608/jppp.2018.41266>
- Miano, R. N., Ayelo, P. M., Musau, R., Hassanali, A., & Mohamed, S. A. (2022). Electroantennogram and machine learning reveal a volatile blend mediating avoidance behavior by *Tuta absoluta* females to a wild tomato plant. *Scientific Reports*, 12(1), 1–16. <https://doi.org/10.1038/s41598-022-13125-0>
- Mirzajani, Z., Hadavi, E., & Kashi, A. (2015). Changes in the essential oil content and selected traits of sweet basil (*Ocimum basilicum* L.) as induced by foliar sprays of citric acid and salicylic acid. *Industrial Crops and Products*, 76, 269–274. <https://doi.org/10.1016/j.indcrop.2015.06.052>
- Monteiro, I. N., Monteiro, O. dos S., Oliveira, A. K. M. de, Favero, S., Garcia, N. Z. T., Fernandes, Y. M. L., Jacinto, G. S. S., Rivero-Wendt, C. L. G., & Matias, R. (2020). Chemical analysis and insecticidal activity of *Ocimum gratissimum* essential oil and its major constituent against *Spodoptera frugiperda* (Smith, 1797) (Lepidoptera: Noctuidae). *Research, Society and Development*, 9(11), e4999119787. <https://doi.org/10.33448/rsd-v9i11.9787>
- Morgan, E. D. (2010). *Biosynthesis in insects*. Royal society of chemistry.
- Mossa, A. T. H. (2016). Green Pesticides: Essential oils as biopesticides in insect-pest management. *Journal of Environmental Science and Technology*, 9(5), 354–378. <https://doi.org/10.3923/jest.2016.354.378>
- Mulugeta, S. M., Sárosi, S., & Radácsi, P. (2023). Physio-morphological trait and bioactive constituents of *Ocimum* species under drought stress. *Industrial Crops and Products*, 205. <https://doi.org/10.1016/j.indcrop.2023.117545>
- Murata, J., Roepke, J., Gordon, H., & De Luca, V. (2008). The leaf epidermome of *Catharanthus roseus* reveals its biochemical specialization. *Plant Cell*, 20(3), 524–542. <https://doi.org/10.1105/tpc.107.056630>
- Murcia-Meseguer, A., Alves, T. J. S., Budia, F., Ortiz, A., & Medina, P. (2018). Insecticidal toxicity of thirteen commercial plant essential oils against *Spodoptera exigua* (Lepidoptera: Noctuidae). *Phytoparasitica*, 46(2), 233–245. <https://doi.org/10.1007/s12600-018-0655-9>
- Naboulsi, I., Aboulmouhajir, A., Kouisni, L., Bekkaoui, F., & Yasri, A. (2018). Plants extracts and secondary metabolites, their extraction methods and use in agriculture for controlling crop stresses and improving productivity: A review. *Acad. J. Med. Plants*, 6, 223-240.
- Nahak, G., Mishra, R., & Sahu, R. (2011). Taxonomic Distribution, Medicinal Properties and Drug Development Potentiality of *Ocimum* (Tulsi). *Drug*

*Invention Today*, 3(6), 95–113.

- Namin, F. R., Naseri, B., & Razmjou, J. (2014). Nutritional performance and activity of some digestive enzymes of the *Cotton bollworm*, *Helicoverpa armigera*, in response to seven tested bean cultivars. *Journal of Insect Science*, 14(93), 1–18. <https://doi.org/10.1673/031.014.93>
- Naseri, B., Abedi, Z., Abdolmaleki, A., Jafary-Jahed, M., Borzoui, E., & Mansouri, S. M. (2017). Fumigant Toxicity and Sublethal Effects of *Artemisia khorassanica* and *Artemisia sieberi* on *Sitotroga cerealella* (Lepidoptera: Gelechiidae). *Journal of Insect Science*, 17(5). <https://doi.org/10.1093/jisesa/iex073>
- Navasero, M. M., Navasero, M. V, Candano, N., & Panis, W. N. De. (2019). Comparative Life History , Fecundity , And Survival Of *Spodoptera exigua* ( *Hübner* ) ( Lepidoptera : Noctuidae ) On *Allium cepa* L . And Other Host Plants in the Philippines. *Philipp. Ent*, 73-84.
- Nerio, L. S., Olivero-Verbel, J., & Stashenko, E. (2010). Repellent activity of essential oils: A review. *Bioresource Technology*, 101(1), 372–378. <https://doi.org/10.1016/j.biortech.2009.07.048>
- Niessen, W. M. A. (2019). Liquid chromatography | Mass spectrometry. In *Encyclopedia of Analytical Science* (Vol. 1). <https://doi.org/10.1016/B978-0-12-409547-2.14213-1>
- Ntalli, N., Kopiczko, A., Radtke, K., Marciniak, P., Rosinski, G., & Adamski, Z. (2014). Biological activity of *Melia azedarach* extracts against *Spodoptera exigua*. *Biologia (Poland)*, 69(11), 1606–1614. <https://doi.org/10.2478/s11756-014-0454-9>
- Nugroho, L. H. (2017). *Structure and products of plant secretory tissues*. Gadjah Mada University Press.
- Nugroho, L. H., & Verpoorte, R. (2002). Secondary metabolism in tobacco. *Plant Cell, Tissue and Organ Culture*, 68, 105-125. <https://doi.org/10.1023/A:1013853909494>
- Nuringtyas, T. R., Choi, Y. H., Verpoorte, R., Klinkhamer, P. G. L., & Leiss, K. A. (2012). Differential tissue distribution of metabolites in *Jacobaea vulgaris*, *Jacobaea aquatica* and their crosses. *Phytochemistry*, 78, 89–97. <https://doi.org/10.1016/j.phytochem.2012.03.011>
- Oliveira, E. R. De, Alves, D. S., Carvalho, G. A., Aazza, S., Kelly, S., & Bertolucci, V. (2018). Toxicity of *Cymbopogon flexuosus* essential oil and citral for *Spodoptera frugiperda*. *Ciência e Agrotecnologia*, 42(4), 408–419.
- Oviedo, A., Van Nieuwenhove, G., Van Nieuwenhove, C., & Rull, J. (2018). Biopesticide effects on pupae and adult mortality of *Anastrepha fraterculus* and *Ceratitis capitata* (Diptera: Tephritidae). *Austral Entomology*, 57(4), 457-464. <https://doi.org/10.1111/aen.12296>

- Padilha De Paula, J., Gomes-Carneiro, M. R., & Paumgarten, F. J. R. (2003). Chemical composition, toxicity and mosquito repellency of *Ocimum selloi* oil. *Journal of Ethnopharmacology*, 88(2–3), 253–260. [https://doi.org/10.1016/S0378-8741\(03\)00233-2](https://doi.org/10.1016/S0378-8741(03)00233-2)
- Pandey, A. K., Singh, P., & Tripathi, N. N. (2014). Chemistry and bioactivities of essential oils of some *Ocimum* species: An overview. *Asian Pacific Journal of Tropical Biomedicine*, 4(9), 682–694. <https://doi.org/10.12980/APJTB.4.2014C77>
- Pandey, V., & Shukla, A. (2015). Acclimation and Tolerance Strategies of Rice under Drought Stress. *Rice Science*, 22(4), 147–161. <https://doi.org/10.1016/j.rsci.2015.04.001>
- Paparang, M., Memah, I. V. V., & Kaligis, I. J. B. (2016). Populasi Dan Persentase Serangan Larva *Spodoptera Exigua* Hubner Pada Tanaman Bawang Daun Dan Bawang Merah Di Desa Ampreng Kecamatan Langowan Barat. *Cocos*, 7(7), 1–10.
- Pardian, P., Noor, T. I., & Kusumah, A. (2016). Analisis Penawaran Dan Permintaan Bawang Merah Di Provinsi Jawa Barat. *Agricore: Jurnal Agribisnis dan Sosial Ekonomi Pertanian Unpad*, 1(2), 189–198. <https://doi.org/10.24198/agricore.v1i2.22711>
- Pareek, S., Sagar, N. A., Sharma, S., & Kumar, V. (2018). Onion (*Allium cepa* L.). *Fruit and Vegetable Phytochemicals: Chemistry and Human Health, 2nd Edition*, 1145–1162. <https://doi.org/10.1002/9781119158042.ch58>
- Pavela, R., & Benelli, G. (2016). Essential Oils as Ecofriendly Biopesticides? Challenges and Constraints. *Trends in Plant Science*, 21(12), 1000–1007. <https://doi.org/10.1016/j.tplants.2016.10.005>
- Pavela, R., Guedes, R. N. C., Maggi, F., Desneux, N., & Benelli, G. (2023). Essential oil antifeedants against armyworms: promises and challenges. *Entomologia Generalis*. <https://doi.org/10.1127/entomologia/2023/1887>
- Peiffer, M., Tooker, J. F., Luthe, D. S., & Felton, G. W. (2009). Plants on early alert: Glandular trichomes as sensors for insect herbivores. *New Phytologist*, 184(3), 644–656. <https://doi.org/10.1111/j.1469-8137.2009.03002.x>
- Pelletier, S. W., Chokshi, H. P., & Desai, H. K. (1986). Separation of Diterpenoid Alkaloid Mixtures using Vacuum Liquid Chromatography. *Journal of Natural Products*, 49(5), 892–900. <https://doi.org/10.1021/np50047a021>
- Polatoğlu, K., Karakoç, Ö. C., Yücel Yücel, Y., Gücel, S., Demirci, B., Demirci, F., & Başer, K. H. C. (2017). Insecticidal activity of *Salvia veneris* Hedge. Essential oil against coleopteran stored product insects and *Spodoptera exigua* (Lepidoptera). *Industrial Crops and Products*, 97, 93–100. <https://doi.org/10.1016/j.indcrop.2016.12.012>

- Popović, Z., Kostić, M., Popović, S., & Skorić, S. (2006). Bioactivities of essential oils from basil and sage to *Sitophilus oryzae* L. *Biotechnology and Biotechnological Equipment*, 20(1), 36–40. <https://doi.org/10.1080/13102818.2006.10817301>
- Popović, Zorica, Kostić, M., Stanković, S., Milanović, S., Sivčev, I., Kostić, I., & Kljajić, P. (2013). Ecologically acceptable usage of derivatives of essential oil of sweet basil, *Ocimum basilicum*, as antifeedants against larvae of the gypsy moth, *Lymantria dispar*. *Journal of Insect Science*, 13, 1–12. <https://doi.org/10.1673/031.013.16101>
- Psychogios, N., Hau, D. D., Peng, J., Guo, A. C., Mandal, R., Bouatra, S., Krishnamurthy, R., Eisner, R., Gautam, B., Young, N., Knox, C., Dong, E., Huang, P., Hollander, Z., Pedersen, T. L., Steven, R., Bamforth, F., Greiner, R., Mcmanus, B., ... Wishart, D. S. (2011). *The Human Serum Metabolome*. *PloS one*, 6(2). <https://doi.org/10.1371/journal.pone.0016957>
- Pusat Data dan Sistem Informasi Pertanian. (2020). <http://pusdatin.setjen.pertanian.go.id/>
- Qin, W., Huang, S., Li, C., Chen, S., & Peng, Z. (2010). Biological activity of the essential oil from the leaves of *Piper sarmentosum* Roxb. (Piperaceae) and its chemical constituents on *Brontispa longissima* (Gestro) (Coleoptera: Hispididae). *Pesticide Biochemistry and Physiology*, 96(3), 132–139. <https://doi.org/10.1016/j.pestbp.2009.10.006>
- Quan, M., Liu, Q. Z., & Liu, Z. L. (2018). Identification of Insecticidal Constituents from the Essential Oil from the Aerial Parts *Stachys riederi* var. *japonica*. *Molecules*, 23(5), 1200. <https://doi.org/10.3390/molecules23051200>
- Quassinti, L., Bramucci, M., Lupidi, G., Barboni, L., Ricciutelli, M., Sagratini, G., Papa, F., Caprioli, G., Petrelli, D., Vitali, L. A., Vittori, S., & Maggi, F. (2013). In vitro biological activity of essential oils and isolated furanosesquiterpenes from the neglected vegetable *Smyrniololus atrum* L. (Apiaceae). *Food Chemistry*, 138(2–3), 808–813. <https://doi.org/10.1016/j.foodchem.2012.11.075>
- Rahmani, S., & Azimi, S. (2020). Fumigant toxicity of three *Satureja* species on tomato leafminers, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Toxin Reviews*, 0(0), 1–12. <https://doi.org/10.1080/15569543.2020.1767651>
- Rajashekar, Y., Raghavendra, A., & Bakthavatsalam, N. (2014). Acetylcholinesterase inhibition by biofumigant (Coumaran) from leaves of *Lantana camara* in stored grain and household insect pests. *BioMed Research International*, 2014. <https://doi.org/10.1155/2014/187019>
- Ramakrishna, A., & Ravishankar, G. A. (2011). Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signaling and Behavior*, 6(11),

1720–1731. <https://doi.org/10.4161/psb.6.11.17613>

Ramírez-Zamora, J., Salinas-Sánchez, D. O., Figueroa-Brito, R., Ramos-López, M., Castañeda-Espinoza, J. D., & Flores-Macías, A. (2020). Botanical extracts from *Dodonaea viscosa* (Sapindales: Sapindaceae) reduce hemocyte counts from *Spodoptera exigua* (Lepidoptera: Noctuidae) with potential insecticidal synergism with *Isaria fumosorosea* (Hypocreales: Cordycipitaceae). *Biocontrol Science and Technology*, 30(12), 1365–1376. <https://doi.org/10.1080/09583157.2020.1826903>

Rautio, P., Markkola, A., Martel, J., Tuomi, J., Ha, E., Kuikka, K., Siitonen, A., Riesco, I. L., & Roitto, M. (2002). Developmental plasticity in birch leaves: defoliation causes a shift from glandular to nonglandular trichomes. *Oikos*, 98(3), 437–446. <https://doi.org/10.1034/j.1600-0706.2002.980308.x>

Regnault-roger, C., Vincent, C., & Arnason, J. T. (2012). Essential oils in insect control: low-risk products in a high-stakes world. *Annual Review of Entomology*, 57, 405–424. <https://doi.org/10.1146/annurev-ento-120710-100554>

Reo, N. V. (2002). NMR-based metabolomics. *Drug and Chemical Toxicology*, 25(4), 375–382. <https://doi.org/10.1081/DCT-120014789>

Rice, P. J., & Coats, J. R. (1994). Insecticidal properties of several monoterpenoids to the house fly (Diptera: Muscidae), red flour beetle (Coleoptera: Tenebrionidae), and southern corn rootworm (Coleoptera: Chrysomelidae). *Journal of Economic Entomology*, 87(5), 1172–1179. <https://doi.org/10.1093/jee/87.5.1172>

Rismayani, Rohimatun, & Laba, I. W. (2018). Hama Utama Pada Pembibitan dan Pengendaliannya. *J. Balitro*, 13(04), 136–144.

Rizwan-Ul-Haq, M., Hu, M. Y., Afzal, M., Bashir, M. H., Gong, L., & Luo, J. (2010). Impact of two medicinal plant extracts on glutathione S-transferase activity in the body tissues of *Spodoptera exigua* (Lepidoptera: Noctuidae). *Pakistan Journal of Botany*, 42(6), 3971–3979.

Rizwan-Ul-Haq, M., Hu, Q. B., Hu, M. Y., Lin, Q. S., & Zhang, W. L. (2009). Biological impact of harmaline, ricinine and their combined effects with *Bacillus thuringiensis* on *Spodoptera exigua* (Lepidoptera: Noctuidae). *Journal of Pest Science*, 82(4), 327–334. <https://doi.org/10.1007/s10340-009-0257-x>

Rossi, P., Cappelli, A., Marinelli, O., Valzano, M., Pavoni, L., Bonacucina, G., Petrelli, R., Pompei, P., Mazzara, E., Ricci, I., Maggi, F., & Nabissi, M. (2020). Mosquitocidal and anti-inflammatory properties of the essential oils obtained from monoecious, male, and female inflorescences of hemp (*Cannabis sativa* L.) and their encapsulation in nanoemulsions. *Molecules*, 25(15), 3451.

<https://doi.org/10.3390/molecules25153451>

- Ruttanaphan, T., Pluempanupat, W., Aungsirirawat, C., Boonyarit, P., Goff, G. Le, Bullangpoti, V., & Isman, M. (2019). Effect of Plant Essential Oils and Their Major Constituents on Cypermethrin Tolerance Associated Detoxification Enzyme Activities in *Spodoptera litura* (Lepidoptera: Noctuidae). *Journal of Economic Entomology*, 112(5), 2167–2176. <https://doi.org/10.1093/jee/toz126>
- Sá, R. D., Santana, A. S. C. O., Silva, F. C. L., Alberto, L., Soares, L., & Randau, K. P. (2016). Anatomical and histochemical analysis of *Dysphania ambrosioides* supported by light and electron microscopy. *Revista Brasileira de Farmacognosia*, 26(5), 533–543. <https://doi.org/10.1016/j.bjp.2016.05.010>
- Sacks, M. M., Silk, W. K., & Burman, P. (1997). Effect of water stress on cortical cell division rates within the apical meristem of primary roots of maize. *Plant physiology*, 114(2), 519–527. <https://doi.org/10.1104/pp.114.2.519>
- Saeed, Q., Ahmad, F., Iqbal, N., & Zaka, S. M. (2019). Chemical control of polyphagous pests on their auxiliary hosts can minimize insecticide resistance: A case study of *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae) in cotton agroecosystem. *Ecotoxicology and Environmental Safety*, 171(November 2018), 721–727. <https://doi.org/10.1016/j.ecoenv.2019.01.038>
- Saharkhiz, M. J., & Ghani, A. (2009). Changes in essential oil content and composition of clary sage (*Salvia sclarea*) aerial parts during different phenological stages. *Medicinal and Aromatic Plant Science and Biotechnology*, 3(1), 90–93.
- Sakul, E. H., Manoppo, J. S., Posumah, D. C., & Tengker, A. C. (2020). Control Of The Beet Armyworm *Spodoptera exigua* (Hübner)(Lepidoptera: Noctuidae) Utilized Four Minahasa Plant Extracts. *Indonesian Biodiversity Journal*, 1(1), 77–90.
- Salsinha, Y. carolina F., Indradewa, D., & Purwestri, Y. A. (2020). Selection of drought-tolerant local rice cultivars from East Nusa Tenggara , Indonesia during vegetative stage. *Biodiversitas*, 21(1), 170–178. <https://doi.org/10.13057/biodiv/d210122>
- Schmelzer, E., Jahnen, W., & Hahlbrock, K. (1988). In situ localization of light-induced chalcone synthase mRNA, chalcone synthase, and flavonoid end products in epidermal cells of parsley leaves. *Proceedings of the National Academy of Sciences*, 85(9), 2989–2993.
- Scott, R. M. (1981). The stationary phase in thin layer chromatography. *Journal of Liquid Chromatography*, 4(12), 2147–2174. <https://doi.org/10.1080/01483918108066850>

- Senthil-Nathan, S., Choi, M. Y., Paik, C. H., & Kalaivani, K. (2008). The toxicity and physiological effect of goniotalamin, a styryl-pyrone, on the generalist herbivore, *Spodoptera exigua* Hübner. *Chemosphere*, 72(9), 1393–1400. <https://doi.org/10.1016/j.chemosphere.2008.03.037>
- Setiawati, W., Hasyim, A., Hudayya, A., & Shepard, B. M. (2014). Evaluation of shade nets and nuclear polyhedrosis virus (SeNPV) to control *Spodoptera exigua* (Lepidoptera: Noctuidae) on shallot in Indonesia. *Advances in Agriculture & Botany*, 6(1), 88–97.
- Shan, L., Chen, L., Gao, F., & Zhou, X. (2019). Diterpenoid alkaloids from *Delphinium naviculare* var. lasiocarpum with their antifeedant activity on *Spodoptera exigua*. *Natural Product Research*, 33(22), 3254–3259. <https://doi.org/10.1080/14786419.2018.1475382>
- Shields, V. D. (Ed.). (2017). *Biological control of pest and vector insects*. BoD–Books on Demand.
- Shiga, T., Shoji, K., Shimada, H., Hashida, S. N., Goto, F., & Yoshihara, T. (2009). Effect of light quality on rosmarinic acid content and antioxidant activity of sweet basil, *Ocimum basilicum* L. *Plant biotechnology*, 26(2), 255–259. <https://doi.org/10.5511/plantbiotechnology.26.255>
- Shorey, H. H. (1963). A Simple Artificial Rearing Medium for the Cabbage Looper. *Journal of economic entomology*, 56(4), 536–537. <https://doi.org/10.1093/jee/56.4.536a>
- Shorey, H. H., & Hale, R. L. (1965). Mass-Rearing of the Larvae of Nine Noctuid Species on a Simple Artificial Medium<sup>12</sup>. *Journal of Economic Entomology*, 58(3), 522–524. <https://doi.org/10.1093/jee/58.3.522>
- Shree, M., Nanda, R. K., & Masakapalli, S. K. (2019). Untargeted metabolite analysis of *Ocimum* leaves shows species specific variations. *BioRxiv*. <https://doi.org/10.1101/673269>
- Siagian, V.J. 2016. Outlook bawang merah. Dalam Nuryati, L., B. Wariant (Eds). Pusat Data dan Sistem Informasi Pertanian Kementerian Pertanian
- Siddhartha, K., C. C. and M. S. (2019). In vitro bioassay of certain botanical oils for their efficacy against maize fall army worm (J.E. Smith) *Spodoptera frugiperda* (Noctuidae: Lepidoptera). *Journal of Entomology and Zoology Studies* 2, 7(5), 606–609.
- Sidou, L.'s F., & Borges, E. M. (2020). Teaching Principal Component Analysis Using a Free and Open Source Software Program and Exercises Applying PCA to Real-World Examples'. *Journal of Chemical Education*, 97, 1666–1676. <https://doi.org/10.1021/acs.jchemed.9b00924>
- Sina, I., Simbolon, S., Sutoro, M., & Ahmad, A. (2021). Assessments: toxicity efficacy of *Tinospora Crispa* (family: Menispermaceae) against *Macrotermes gilvus* (family: Termitidae) in oil palm plantation.

*In Proceedings of the 1st International Conference on Economics Engineering and Social Science, InCEEES 2020, 17-18 July, Bekasi, Indonesia.* <https://doi.org/10.4108/eai.17-7-2020.2302970>

- Singh, P., & Pandey, A. K. (2018). Prospective of Essential Oils of the Genus *Mentha* as Biopesticides: A Review. *Frontiers in Plant Science*, 9(September), 1–14. <https://doi.org/10.3389/fpls.2018.01295>
- Sirousmehr, A., & Asgharipour, M. R. (2014). Effect of drought stress levels and organic manures on yield, essential oil content and some morphological characteristics of sweet basil (*Ocimum basilicum*). *Advances in Environmental Biology*, 8(4), 880-885.
- Sletvold, N., Huttunen, P., Handley, R., Kärkkäinen, K., & Ågren, J. (2010). Cost of trichome production and resistance to a specialist insect herbivore in *Arabidopsis lyrata*. *Evolutionary Ecology*, 24(6), 1307–1319. <https://doi.org/10.1007/s10682-010-9381-6>
- Snart, C. J. P., Hardy, I. C. W., & Barrett, D. A. (2015). Entometabolomics: Applications of modern analytical techniques to insect studies. *Entomologia Experimentalis et Applicata*, 155(1), 1–17. <https://doi.org/10.1111/eea.12281>
- Soumia, P. S., Karuppaiah, V., Mahajan, V., & Singh, M. (2020). Beet Armyworm *Spodoptera exigua*: Emerging Threat to Onion Production. *National Academy Science Letters*, 43(5), 473–476. <https://doi.org/10.1007/s40009-020-00892-5>
- Spochacz, M., Chowański, S., Walkowiak-Nowicka, K., Szymczak, M., & Adamski, Z. (2018). Plant-Derived Substances Used Against Beetles–Pests of Stored Crops and Food—and Their Mode of Action: A Review. *Comprehensive Reviews in Food Science and Food Safety*, 17(5), 1339–1366. <https://doi.org/10.1111/1541-4337.12377>
- Sukirno, S., Tufail, M., Rasool, K. G., El Salamouny, S., Sutanto, K. D., & Aldawood, A. S. (2017). The effectiveness of spinosad and neem extract against *Spodoptera littoralis* (Boisd.) and *Spodoptera exigua* (Hubner): Exploring possibilities to enhance the bio-pesticide persistence with natural UV protectants under field-sunlight conditions of Saudi Arabia. *Pakistan Journal of Agricultural Sciences*, 54(4), 743–751. <https://doi.org/10.21162/PAKJAS/17.5306>
- Sultana, B., Anwar, F., & Ashraf, M. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*, 14(6), 2167–2180. <https://doi.org/10.3390/molecules14062167>
- Sundararajan, B., Moola, A. K., Vivek, K., & Kumari, B. D. R. (2018). Formulation of nanoemulsion from leaves essential oil of *Ocimum basilicum* L. and its antibacterial, antioxidant and larvicidal activities (*Culex quinquefasciatus*). *Microbial Pathogenesis*, 125(May), 475–485.

<https://doi.org/10.1016/j.micpath.2018.10.017>

- Szymanowska, U., Złotek, U., Karaś, M., & Baraniak, B. (2015). Anti-inflammatory and antioxidative activity of anthocyanins from purple basil leaves induced by selected abiotic elicitors. *Food Chemistry*, 172(February 2015), 71–77. <https://doi.org/10.1016/j.foodchem.2014.09.043>
- Targett, N. M., Kilcoyne, J. P., & Green, B. (1979). Vacuum liquid chromatography: an alternative to common chromatographic methods. *The Journal of Organic Chemistry*, 44(26), 4962-4964. <https://doi.org/10.1021/jo00394a045>
- Tattini, M. (2000). Flavonoids accumulate in leaves and glandular trichomes of *Phillyrea latifolia* exposed to excess solar radiation. *The New Phytologist*, 148(1), 69-77. <https://doi.org/10.1046/j.1469-8137.2000.00743.x>
- Teng, H., Yuan, Y., Zhang, T., Chang, X., & Wang, D. (2020). Evaluation of the sublethal effect of tetrachlorantraniliprole on *Spodoptera exigua* and its potential toxicity to two non-target organisms. *PLoS ONE*, 15(11 November), 1–15. <https://doi.org/10.1371/journal.pone.0242052>
- Teshika, J. D., Zakariyyah, A. M., Zaynab, T., Zengin, G., Rengasamy, K. R., Pandian, S. K., & Fawzi, M. M. (2019). Traditional and modern uses of onion bulb (*Allium cepa* L.): a systematic review. *Critical Reviews in Food Science and Nutrition*, 59(0), S39–S70. <https://doi.org/10.1080/10408398.2018.1499074>
- Thakur, M., Bhattacharya, S., Khosla, P. K., & Puri, S. (2019). Improving production of plant secondary metabolites through biotic and abiotic elicitation. *Journal of Applied Research on Medicinal and Aromatic Plants*, 12(November 2018), 1–12. <https://doi.org/10.1016/j.jarmap.2018.11.004>
- Tian, D., Tooker, J., Peiffer, M., Chung, S. H., & Felton, G. W. (2012). Role of trichomes in defense against herbivores: Comparison of herbivore response to woolly and hairless trichome mutants in tomato (*Solanum lycopersicum*). *Planta*, 236(4), 1053–1066. <https://doi.org/10.1007/s00425-012-1651-9>
- To'bungan, N., & Jati, W. N. (2022). Larvicidal activity of Knobweed (*Hyptis capitata*) leaves ethanolic extract and fraction against *Culex quinquefasciatus*. *Biogenesis: Jurnal Ilmiah Biologi*, 10(2), 236-243. <https://doi.org/10.24252/bio.v10i2.31825>
- Tomar, H., Rawat, A., Nagarkoti, K., Prakash, O., Kumar, R., Srivastava, R. M., Rawat, S., & Rawat, D. S. (2023). *Ocimum gratissimum* L. and *Ocimum sanctum* L.: Comparative compositional analysis of essential oils and in-vitro biological activities with in-silico PASS prediction and ADME/Tox studies. *South African Journal of Botany*, 157, 360–371. <https://doi.org/10.1016/j.sajb.2023.04.014>

- Traw, M. B., & Bergelson, J. (2003). Interactive Effects of Jasmonic Acid, Salicylic Acid, and Gibberellin on Induction of Trichomes in *Arabidopsis*. *Plant Physiology*, 133(3), 1367–1375. <https://doi.org/10.1104/pp.103.027086>
- Triwidodo, H., & Tanjung, M. H. (2020). Hama Penyakit Utama Tanaman Bawang Merah (*Allium Ascalonicum*) dan Tindakan Pengendalian di Brebes, Jawa Tengah. *Agrovigor: Jurnal Agroekoteknologi*, 13(2), 149–154. <https://doi.org/10.21107/agrovigor.v13i2.7131>
- Tulashie, S. K., Adjei, F., Abraham, J., & Addo, E. (2021). Potential of neem extracts as a natural insecticide against fall armyworm ( *Spodoptera frugiperda* ( Smith ) ( Lepidoptera : Noctuidae )) Case Studies in Chemical and Environmental Engineering Potential of neem extracts as natural insecticide against fall. *Case Studies in Chemical and Environmental Engineering*, 4, 100–130. <https://doi.org/10.1016/j.cscee.2021.100130>
- Ujiyanti, F., Trisyono, Y. A., Witjaksono, W., & Suputa, S. (2019). Population of *Spodoptera exigua* Hübner during On- and Off-Season of Shallot in Bantul Regency, Yogyakarta. *Jurnal Perlindungan Tanaman Indonesia*, 23(2), 261. <https://doi.org/10.22146/jpti.36740>
- Usman, M., Raheem, Z., Ahsan, T., Iqbal, A., Safaraz, Z. N., & Haq, Z. (2013). Morphological , Physiological and Biochemical Attributes as Indicators for Drought Tolerance in Rice (*Oryza sativa* L .). *European Journal of Biological Sciences*, 5(1), 23–28. <https://doi.org/10.5829/idosi.ejbs.2013.5.1.1104>
- Van Schie, C. C. N., Haring, M. A., & Schuurink, R. C. (2007). Tomato linalool synthase is induced in trichomes by jasmonic acid. *Plant Molecular Biology*, 64(3), 251–263. <https://doi.org/10.1007/s11103-007-9149-8>
- Varmuza, K. (2000). Applied chemometrics: from chemical data to relevant information. In *1st Conference on Chemistry* (6-9).
- Wahba, T. F. (2020). Antifeedant activity of three essential oils and their nanoemulsions against antifeedant activity of three essential oils and their nanoemulsions against the rice weevil *Sitophilus oryzae* (L.). *Egyptian Scientific Journal of Pesticides*, 2(June), 19–31.
- Wang, X., Shen, C., Meng, P., Tan, G., & Lv, L. (2021). Analysis and review of trichomes in plants. *BMC Plant Biology*, 21(1), 1–11. <https://doi.org/10.1186/s12870-021-02840-x>
- War, A. R., Taggar, G. K., Hussain, B., Taggar, M. S., Nair, R. M., & Sharma, H. C. (2018). Special Issue: Using non-model systems to explore plant-pollinator and plant-herbivore interactions: Plant defence against herbivory and insect adaptations. *AoB PLANTS*, 10(4), 1–19. <https://doi.org/10.1093/aobpla/ply037>

- Werker, E., Putievsky, E., Ravid, U., Dudai, N., & Katzir, I. (1993). Glandular hairs and essential oil in developing leaves of *Ocimum basilicum* L.(Lamiaceae). *Annals of Botany*, 71(1), 43-50. <https://doi.org/10.1006/anbo.1993.1005>
- Wibisono, I. I., Trisyono, Y. A., Martono, E., & Purwantoro, A. (2007). Evaluasi Resistensi Terhadap Metoksifenoazida pada *Spodoptera exigua* di Jawa. *Jurnal Perlindungan Tanaman Indonesia*, 13(2), 127–135.
- Wilson, C., & Tisdell, C. (2001). Why farmers continue to use pesticides despite environmental, health and sustainability costs. *Ecological Economics*, 39(3), 449–462. [https://doi.org/10.1016/S0921-8009\(01\)00238-5](https://doi.org/10.1016/S0921-8009(01)00238-5)
- Wu, Q., & Brown, M. R. (2006). Signaling and function of insulin-like peptides in insects. *Annual Review of Entomology*, 51, 1–24. <https://doi.org/10.1146/annurev.ento.51.110104.151011>
- Wulandari, R., Santoso, R. E., Prasetyo, D., Lestari, A., Mizar, A., & Puspitasari, P. (2019). Increasing the weight of onion bulbs due to the reduction of *Spodoptera exigua* using a portable light trap. *IOP Conference Series: Materials Science and Engineering*, 694(1). <https://doi.org/10.1088/1757-899X/694/1/012010>
- Xie, Z., Kapteyn, J., & Gang, D. R. (2008). A systems biology investigation of the MEP/terpenoid and shikimate/phenylpropanoid pathways points to multiple levels of metabolic control in sweet basil glandular trichomes. *Plant Journal*, 54(3), 349–361. <https://doi.org/10.1111/j.1365-3113X.2008.03429.x>
- Xu, J., Sheng, Z., & Palli, S. R. (2013). Juvenile Hormone and Insulin Regulate Trehalose Homeostasis in the Red Flour Beetle, *Tribolium castaneum*. *PLoS Genetics*, 9(6). <https://doi.org/10.1371/journal.pgen.1003535>
- Yan, P. cheng, Wen, C. wei, Zhang, S. zhi, Zhang, Z. da, Xu, J. ping, & Deng, M. jie. (2018). A toxicological, metabonomic and transcriptional analysis to investigate the property of mulberry 1-deoxynojirimycin against the growth of *Samia cynthia ricini*. *Pesticide Biochemistry and Physiology*, 152(March), 45–54. <https://doi.org/10.1016/j.pestbp.2018.08.009>
- Yarou, B. B., Bawin, T., Boullis, A., Heukin, S., Lognay, G., Verheggen, F. J., & Francis, F. (2018). Oviposition deterrent activity of basil plants and their essentials oils against *Tuta absoluta* (Lepidoptera: Gelechiidae). *Environmental Science and Pollution Research*, 25(30), 29880–29888. <https://doi.org/10.1007/s11356-017-9795-6>
- Yasri, A., Naboulsi, I., Kouisni, L., & Bekkaoui, F. (2018). Plants extracts and secondary metabolites, their extraction methods and use in agriculture for controlling crop stresses and improving productivity: A review. *Acad. J. Med. Plants*, 6(8), 223–240. <https://doi.org/10.15413/ajmp.2018.0139>

- Yooboon, T., Pengsook, A., Poonsri, W., Pluempanupat, W., & Bullangpoti, V. (2020). Toxicity of Phenylpropanoids from *Alpinia galanga* (Zingiberaceae) extracts against *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae). *Phytoparasitica*, 48(5), 833–840. <https://doi.org/10.1007/s12600-020-00830-7>
- Zarei, H., Fakheri, B. A., Bahabadi, S. E., & Solouki, M. (2015). Increasing of chavicol o-methyl transferase gene expression (cvomt) and methyl chavicol value of basil (*Ocimum basilicum*) by salicylic acid. *J. Bio. Env. Sci*, 6, 46–53.
- Zhandi, W. A. N. G., La, Z. H. A. N. G., Bailian, D. I. N. G., Yundong, S. H. I., Yani, W. A. N. G., Guangqiu, L. U., & Lin, J. I. A. (2021). Regulation of pakchoi's secondary metabolites on the behavior of female *Plutella xylostella* (Lepidoptera: Plutellidae). *农药学报*, 23(2), 323–330.
- Zhang, J. F., Chen, L., Huang, S., Shan, L. H., Gao, F., & Zhou, X. L. (2017). Diterpenoid Alkaloids from Two Aconitum Species with Antifeedant Activity against *Spodoptera exigua*. *Journal of Natural Products*, 80(12), 3136–3142. <https://doi.org/10.1021/acs.jnatprod.7b00380>
- Zhang, Q. W., Lin, L. G., & Ye, W. C. (2018). Techniques for extraction and isolation of natural products: A comprehensive review. *Chinese Medicine (United Kingdom)*, 13(1), 1–26. <https://doi.org/10.1186/s13020-018-0177-x>
- Zhang, X. G. Z., I, X. L., Ao, Y. L. G., Iu, Y. L., Ong, W. X. D., & Iao, C. X. (2019). Oviposition deterrents in larval frass of potato tuberworm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). *Neotropical Entomology*, 48, 496–502. <https://doi.org/10.1007/s13744-018-0655-y>
- Zhang, Y. N., He, P., Xue, J. P., Guo, Q., Zhu, X. Y., Fang, L. P., & Li, J. B. (2017). Insecticidal activities and biochemical properties of *Pinellia ternata* extracts against the beet armyworm *Spodoptera exigua*. *Journal of Asia-Pacific Entomology*, 20(2), 469–476. <https://doi.org/10.1016/j.aspen.2017.03.003>
- Zhang, Z., & Shao, Z. (2015). An improved K-OPT local search algorithm for the vertex separator problem. *Journal of Computational and Theoretical Nanoscience*, 12(11), 4942–4958. <https://doi.org/10.1166/jctn.2015.4464>
- Zhao, A., Li, Y., Leng, C., Wang, P., & Li, Y. (2019). Inhibitory effect of protease inhibitors on larval midgut protease activities and the performance of *Plutella xylostella* (Lepidoptera: Plutellidae). *Frontiers in Physiology*, 10(JAN), 1–9. <https://doi.org/10.3389/fphys.2018.01963>
- Zheng, S., Henken, B., Wietsma, W., Sofiari, E., Jacobsen, E., Krens, F. A., & Kik, C. (2000). Development of bio-assays and screening for resistance to beet armyworm (*Spodoptera exigua* Hubner) in *Allium cepa* L. and its wild relatives. *Euphytica*, 114(1), 77–85.

<https://doi.org/10.1023/A:1004089424419>

Ziv, C., Zhao, Z., Gao, Y. G., & Xia, Y. (2018). Multifunctional roles of plant cuticle during plant-pathogen interactions. *Frontiers in Plant Science*, 9, 1088. <https://doi.org/10.3389/fpls.2018.01088>

Złotek, U., Michalak-Majewska, M., & Szymanowska, U. (2016). Effect of jasmonic acid elicitation on the yield, chemical composition, and antioxidant and anti-inflammatory properties of essential oil of lettuce leaf basil (*Ocimum basilicum* L.). *Food Chemistry*, 213, 1–7. <https://doi.org/10.1016/j.foodchem.2016.06.052>