

DAFTAR PUSTAKA

- Aguiar, A. J. and Zelmer, J. E., 1969, "Dissolution Behavior of Polymorphs of Chloramphenicol Palmitate and Mefenamic Acid", J. Pharm. Sci., 58, 983 - 987.
- Alam, A. S., Parrot, E. L., 1971, "Effect of Dissolution Media on Disintegration and Dissolution of Hydrochlorothiazide Tablets", J. Pharm. Sci., 60, 795 - 797.
- Allen, L. V., Yanchick, V. A., Maness, D. D., 1977, "Dissolution Rates of Corticosteroids Utilizing Sugar-Glass Dispersions", J. Pharm. Sci., 66, 494 - 496.
- Armstrong, N. A., Haines Nut, R. F., 1973, "Dissolution Rates and Surface Area of Compacted Powder System", J. Pharm. Pharmacol., 25, 147.
- Carstensen, J. T., 1974, "Theories of Dissolution: Single Particulate System", in: Leeson, L. J. and Carstensen, J. T. (Ed.), Dissolution Technology, The Industrial Pharmaceutical Technology Section of Academy of Pharmaceutical Sciences, Washington D.C. 1 - 32.
- Chiou, W. L., 1977, "Pharmaceutical Applications of Solid Dispersion System: X-Ray Diffraction and Aqueous Solubility Studies on Griseofulvin-Polyethylen Glycol 6000 System", J. Pharm. Sci., 66, 989 - 991.
- Chiou, W. L. and Riegelman, S., 1971, "Pharmaceutical Applications of Solid Dispersion System", J. Pharm. Sci., 60, 1281 - 1302.
- Clarke, E. G. C., 1971, Isolation and Identification of Drugs: in Pharmaceutical, Body Fluids and Postmortem Materials, 1st. Ed., The Pharmaceutical Press, London, 1089.
- Finholt, P., 1974, "Influence of Formulation on Dissolution Rate" in: Leeson, L. J. and Carstensen, J. T., Dissolution Technology, The Industrial Pharmaceutical Technology Section of Academy of Pharmaceutical Sciences, Washington D. C. 106 - 146.
- Fudholi, A., 1986, "Disolusi" dalam: Fudholi, A., Marchaban, Yuwono, T., Kursus Penyebaran, Fabrikasi, Kontrol Stabilitas dan Disolusi, Fakultas Farmasi Universitas Gadjah Mada, Yogyakarta, 1 - 3.
- Ghanem, A., Meshali, M., Ibraheem, Y., 1980, "Dissolution Rates of Sulfamethoxazole Utilizing Sugar Glass Dispersions", J. Pharm. Pharmacol., 32, 675 - 677.



- Gibaldi, M., 1970, "Biopharmaceutics", in: Lachman, L., Lieberman, A. H., Kanig, J. L. (Ed.), The Theory and Practice of Industrial Pharmacy, Lea & Febiger, Philadelphia, 246 - 268.
- Gibaldi, M., Fieldman, S., 1970, "Mechanism of Surfactant Effects on Drug Absorption", J. Pharm. Sci., 59, 579 - 589.
- Goldberg, A. H., Gibaldi, M., Kanig, J. L., 1965, "Increasing Dissolution Rates and Gastrointestinal Absorption of Drug via Solid Solution and Eutectic Mixture", J. Pharm. Sci., 54, 1145 - 1148.
- Hanson, W. A., 1982, Handbook of Dissolution Testing, Pharmaceutical Technology Publications, Springfield, Oregon, 13 - 25.
- Higuchi, T. and Ikeda, M., 1974, "Rapidly Dissolving Forms of Digoxin: Hidroquinon Complex", J. Pharm. Sci., 63, 809.
- Kaplan, S., 1974, "Biological Implications of In Vitro Dissolution Testing", in: Leeson, L. J. and Carstensen, J. T. (Ed.), Dissolution Technology, The Industrial Pharmaceutical Technology Section of Academy of Pharmaceutical Sciences, Washington D. C., 163 - 195.
- Lowenthal, W., 1972, "Disintegration of Tablets" J. Pharm. Sci., 61, 1695 - 1709.
- Martin, A., Swarbrick, J., Cammarata, A., 1988, Physical Pharmacy: Physical Chemical Principles in the Pharmaceutical Sciences, 3th., Lea & Febiger, Philadelphia, 89 - 90.
- Riswaka, S., Moechtar, Imono, A. D., 1981, "Hubungan Ukuran Partikel Prednison dengan Kadarnya Dalam Darah", Laporan Penelitian, Lembaga Penelitian Universitas Gadjah Mada.
- Riswaka, S., Moechtar, Mimiek, M., 1981, "Hubungan Viskositas Relatif dengan Kecepatan Pelarutan Bahan Obat Tablet Sulfadiazina", Laporan Penelitian, Lembaga Penelitian Universitas Gadjah Mada.
- Shah, K. A., Needham, T. E., 1979, "Correlation of Urinary Excretion with In Vitro Dissolution Using Several Dissolution Methods for Hydrochlorothiazide Formulations", J. Pharm. Sci., 68, 1486 - 1490.



- Shargel, L., and Andrew, E. C., 1985, Applied Biopharmaceutics and Pharmacokinetics, 2nd. Ed., Appleton-Century-Crofts, Norwalk, Connecticut, 67 - 78.
- Shefter, E., and Higuchi, T., 1963, "Dissolution Behavior of Crystalline Solvated and Nonsolvated Forms of Some Pharmaceutical", J. Pharm. Sci., 52, 781 - 791.
- Simonelli, A. P., Mehta, S. C., and Higuchi, W. I., 1969, "Dissolution Rates of High Energy Polyvinylpyrrolidone(PVP)-Sulfathiazole Coprecipitate", J. Pharm. Sci., 58, 538 - 549.
- Stoll, R. G., Bates, T. R., and Swarbrick, J., 1973, "In-Vitro Dissolution and In Vivo Absorption of Nitrofurantoin from Deoxycholic Acid Coprecipitate", J. Pharm. Sci., 62, 65 - 68.
- Stupak, E. I., and Bates, T. R., 1972, "Enhanced Absorption and Dissolution of Reserpine from Reserpine-Polyvinylpyrrolidone Coprecipitate", J. Pharm. Sci., 61, 400 - 403.
- Stupak, E. I., and Bates, T. R., 1973, "Enhanced Absorption of Digitoxine from Orally Administered Digitoxine-Polyvinylpyrrolidone Coprecipitates", J. Pharm. Sci., 62, 1806 - 1809.
- Tan dan Kirana, R., 1979, Obat-obat Penting: Khasiat dan Penggunaannya, Edisi III, Jakarta, 126 - 127.
- Tawashi, R., Bisailon, S., 1978, "L'ecoulement Des Fluides et Son Implication Vis a Vis de la Dissolution Des Medicaments", Pharm. Acta Helv., 5, 245 - 278.
- Tuladhar, M. D., Carless, J. E., and Summers, M. P., 1983, "The Effect of Polymorphism, Particle Size and Compression Pressure on the Dissolution Rate of Phenylbutazone Tablets", J. Pharm. Pharmacol., 35, 269 - 274.
- Wagner, J. G., 1971, Biopharmaceutics and Relevant Pharmacokinetics, 1st. Ed., Drug Intelligence Publications, Hamilton, Illinois, 98 - 117.
- Wood, J. H., Syarto, J. E., and Letterman, H., 1965, "Improved Holder of Intrinsic Dissolution Rate Studies", J. Pharm. Sci., 54, 1068.
- Wurster, D. E., and Taylor, P. W., 1965, "Dissolution Rates", J. Pharm. Sci., 54, 169 - 175.



Yuwono, T., 1987, "Pengaruh Kafeina terhadap Kelarutan dan Ketersediaan Hayati o-Etoksibenzamida pada Tikus Jantan", Disertasi, Institut Teknologi Bandung, Bandung, 71.

Yuwono, T., 1988, "Pembentukan Dispersi Solid Asam Mefenamat untuk Meningkatkan Kecepatan Disolusinya", Laporan Penelitian, Fakultas Farmasi, Universitas Gadjah Mada, Yogyakarta, 34.