



DAFTAR PUSTAKA

- Aggarwal, B. B., Gupta, S. C., & Kim, J. H. (2012). Historical perspectives on tumor necrosis factor and its superfamily: 25 years later, a golden journey. *Blood*, 119(3): 651–665.
- Akash, M. S. H., Rehman, K., & Liaqat, A. (2018). Tumor Necrosis Factor-Alpha: Role in Development of Insulin Resistance and Pathogenesis of Type 2 Diabetes Mellitus. *J. Cell. Biochem.*, 119(1): 105–110.
- American Diabetes Association (2014). Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*, 37(Supplement 1): 81–90.
- Amoozadeh, Y., Dan, Q., Xiao, J., Waheed, F., & Szászi K. (2015). Tumor necrosis factor- α induces a biphasic change in claudin-2 expression in tubular epithelial cells: role in barrier functions. *Am. J. Physiol. Cell. Physiol.*, 309(1):C38-50.
- Autenrieth, D.M., & Baumgart, D.C. (2012). Toxic megacolon. *Inflamm. Bowel Dis.*, 18(3):584-91.
- Badole, S.L, Chaudhari, S. M., Bagul, P. P., Mahamuni, S. P., Khose, R. D., Joshi, A. C., et al. (2013). Effect of concomitant administration of L-glutamine and cycloart-23-ene-3 β , 25-diol (B2) with sitagliptin in GLP-1 (7-36) amide secretion, biochemical and oxidative stress in streptozotocin - nicotinamide induced diabetic Sprague Dawley rats. *PLoS One*. 8(8):e72817.
- Badr, G., Sayed, L. H., Omar, H. E. M., Abd El-Rahim, A. M., Ahmed, E. A., & Mahmoud M.H. (2017). Camel Whey Protein Protects B and T Cells from Apoptosis by Suppressing Activating Transcription Factor-3 (ATF-3) - Mediated Oxidative Stress and Enhancing Phosphorylation of AKT and I κ B- α in Type I Diabetic Mice, *Cell. Physiol. Biochem.*, 41: 41–54.
- Barman, S., & Srinivasan, K. (2019). Ameliorative effect of zinc supplementation on compromised small intestinal health in streptozotocin-induced diabetic rats. *Chem. Biol. Interac.*, 307: 37–50.
- Barone, F.C., Deegan, J.F., Price, W.J., Fowler, P.J., Fondacaro, J.D., & Ormsbee 3rd, H.S. (1990). Cold-restraint stress increases rat fecal pellet output and colonic transit. *Am. J. Physiol.*, 258, G329-337.
- Barta, B.P., Onhausz, B., Al Doghmi, A., Szalai, Z., Balázs, J., Bagyánszki, M., & Bódi, N. (2023). Gut region-specific TNFR expression: TNFR2 is more affected than TNFR1 in duodenal myenteric ganglia of diabetic rats. *World J. Diabetes*, 14(1):48-61.



- Bayramgürler, D., Karson, A., Ozer, C., & Utkan T. (2013). Effects of long-term etanercept treatment on anxiety- and depression-like neurobehaviors in rats. *Physiol. Behav.*, 2;119:145-8.
- Becker, L., Nguyen, L., Gill, J., Kulkarni, S., Pasricha, P. J., Habtezion, A. (2018). Age-dependent shift in macrophage polarisation causes inflammation-mediated degeneration of enteric nervous system. *Gut*, 67(5):827-836.
- Bodi, N., Talapka, P., Poles, M. Z., Hermesz, E., Jancsó, Z., Katarova, Z., *et al.* (2012). Gut Region-Specific Diabetic Damage to the Capillary Endothelium Adjacent to the Myenteric Plexus. *Microcirculation*, 19(4): 316–326.
- Bódi, N., Chandrakumar, L., Al Doghmi, A., Mezei, D., Szalai, Z., Barta, B. P., *et al.* (2021). Intestinal region-specific and layer-dependent induction of tnf- α in rats with streptozotocin-induced diabetes and after insulin replacement. *Cells*, 10(9): 1–14.
- Browning, K. N., & Travagli, R. A. (2014). Central nervous system control of gastrointestinal motility and secretion and modulation of gastrointestinal functions. *Compr. Physiol.*, 4(4): 1339–1368.
- Chandrasekharan, B., Anitha, M., Blatt, R., Shahnavaz, N., Kooby, D., Staley, C., *et al.* (2011). Colonic motor dysfunction in human diabetes is associated with enteric neuronal loss and increased oxidative stress. *Neurogastroenterol. Motil.*, 23(2):131-8, e26.
- Chandramouli, C., Reichelt, M. E., Curl, C. L., Varma, U., Bienvenu, L. A., Koutsifeli, P., *et al.* (2018). Diastolic dysfunction is more apparent in STZ-induced diabetic female mice, despite less pronounced hyperglycemia. *Sci. Rep.*, 8(2346): 1–13.
- Chang, X., Qin, Y., Jin Z, Xi, T.F, Yang, X., Lu, Z. H., *et al.* (2015). Interleukin-6 (IL-6) mediated the increased contraction of distal colon in streptozotocin-induced diabetes in rats via IL-6 receptor pathway. *Int. J. Clin. Exp. Pathol.*, 8(5): 4514–4524.
- Charan, J., & Kantharia, N. D. (2013). How to calculate sample size in animal studies?. *J. Pharmacol. Pharmacother.*, 4(4): 303–306.
- Chen, P.M., Gregersen, H., & Zhao, J.B. (2015). Advanced glycation end-product expression is upregulated in the gastrointestinal tract of type 2 diabetic rats. *World J. Diabetes*, 6(4): 662–72.
- Chen, W., Lu, C., Hirota, C., Iacucci, M., Ghosh, S., & Gui, X. (2017). Smooth muscle hyperplasia/hypertrophy is the most prominent histological change in Crohn's fibrostenosing bowel strictures: A semiquantitative analysis by using a novel histological grading scheme. *J. Crohn's Colitis.*, 11, 92–104.



- Costa, F. R. C., Françozo, M.C., de Oliveira, G.G., Ignacio, A., Castoldi, A., Zamboni, D.S., *et al.* (2016). Gut microbiota translocation to the pancreatic lymph nodes triggers NOD2 activation and contributes to T1D onset. *J.Exp.Med.*, 213(7): 1223–1239.
- Costa, M., Spencer, N. J., & Brookes, S. J. H. (2021). The role of enteric inhibitory neurons in intestinal motility. *Auton. Neurosci.*, 235: 1–11.
- Crumbie, L. (2021). Histology of the lower digestive. <https://www.kenhub.com/en/library/anatomy/histology-of-the-lower-digestive-tract>. Diakses pada tanggal 19 Maret 2022.
- Dewanjee, S., Das, S., Das, A. K., Bhattacharjee, N., Dihingia, A., Dua, T. K., *et al.* (2018). Molecular mechanism of diabetic neuropathy and its pharmacotherapeutic targets. *Eur. J. Pharmacol.*, 833:472-523.
- de Zoete, M. R., Palm, N. W., Zhu, S., & Flavell, R. A. (2014). Inflammasomes. *Cold Spring Harb. Perspect. Biol.*, 6(12): 1-22.
- Dik, B., Bahcivan, E., Faki H. E., & Uney K. (2018). Combined Treatment with Interlukin-1 and Tumor Necrosis Factor-Alpha Antagonists Improve Type 2 Diabetes in Rats. *Can. J. Physiol. Pharmacol.*, 96(8): 751–756.
- DiPetrillo, K., Coutermash, B., & Gesek, F. A. (2003). Urinary tumor necrosis factor contributes to sodium retention and renal hypertrophy during diabetes. *Am. J. Physiol. Renal. Physiol.*, 284(1): F113-21.
- Dora, D., Ferenczi, S., Stavely, R., Toth, V. E., Varga, Z. V., Kovacs, T., *et al.* (2021). Evidence of a Myenteric Plexus Barrier and Its Macrophage-Dependent Degradation During Murine Colitis: Implications in Enteric Neuroinflammation. *Cell Mol. Gastroenterol. Hepatol.*, 12(5): 1617–1641.
- Du, F., Wang, L., Qian, W., & Liu, S. (2009). Loss of enteric neurons accompanied by decreased expression of GDNF and PI3K / Akt pathway in diabetic rats. *Neurogastroenterol. Motil.*, 21: 1229–1235.
- Du, Y. T., Rayner, C. K., Jones, K. L., Talley, N. J., & Horowitz, M. (2018). Gastrointestinal Symptoms in Diabetes: Prevalence, Assessment, Pathogenesis, and Management. *Diabetes Care*, 41: 627–637.
- Dumitriu, I. L., Gurzu, B., Cojocaru, E., Slătineanu, S. M, & Enea, M. (2011). Validation of GOD / PAP Method for Quantitative Determination of Glucose Concentration in Human Serum. *Rev. Romana. Med. Lab.*, 19(1): 85–100.
- D'Arpino, M.C., Fuchs, A.G., Sanchez, S.S., & Honore, S.M. (2018). Extracellular matrix remodeling and TGF- β 1 / Smad signaling in diabetic colon mucosa. *Cell Biol. Int.*, 42(4):443–56.



- Elmarakby, A. A., & Sullivan, J. C. (2012). Relationship between oxidative stress and inflammatory cytokines in diabetic nephropathy. *Cardiovasc. Ther.*, 30(1): 49–59.
- El-Shaer, N. O., Hegazy, A. M., & Muhammad, M. H. (2023). Protective effect of quercetin on pulmonary dysfunction in streptozotocin-induced diabetic rats via inhibition of NLRP3 signaling pathway. *Environ. Sci. Pollut. Res. Int.* 30(14):42390-42398.
- Feng, Y., Weng, H., Ling, L., Zeng, T., Zhang, Y., Chen, D., et al. (2019). Modulating the gut microbiota and inflammation is involved in the effect of Bupleurum polysaccharides against diabetic nephropathy in mice. *Int. J. Biol. Macromol.*, 132: 1001–1011.
- Fricker, M., Tolkovsky, A. M., Borutaite, V., Coleman, M., & Brown, G. C. (2018). Neuronal cell death. *Physiol. Rev.*, 98(2): 813–880.
- Furman, B. L. (2015). Streptozotocin-Induced Diabetic Models in Mice and Rats. *Curr. Protoc. Pharmacol.*, 70: 5.47.1-5.47.20.
- Ghasemi, A., Khalifi, S., & Jedi, S. (2014). Streptozotocin-nicotinamide-induced rat model of type 2 diabetes (Review). *Acta Physiol. Hung.*, 101(4): 408–420.
- Gheibi, S., Kash, K. & Ghasemi, A. (2017). A practical guide for induction of type-2 diabetes in rat: Incorporating a high-fat diet and streptozotocin. *Biomed. Pharmacother.*, 95: 605–613.
- Gough, P. and Myles, I. A. (2020). Tumor Necrosis Factor Receptors: Pleiotropic Signaling Complexes and Their Differential Effects. *Front. Immunol.*, 11: 1–14.
- Grauballe, M. B., Østergaard, J. A., Schou, S., Flyvbjerg, A., & Holmstrup, P. (2015). Effects of TNF- α blocking on experimental periodontitis and type 2 diabetes in obese diabetic Zucker rats. *J. Clin. Periodontol.*, 42(9): 807–816.
- Guillot, X., Prati, C., Sondag, M., & Wendling, D. (2017). Etanercept for treating axial spondyloarthritis. *Expert. Opin. Biol. Ther.*, 17(9): 1173–1181.
- Guo, M., Chen, Q., Huang, Y., Wu, Q., Zeng, Y., Tan, X., et al. (2023). High Glucose-Induced Kidney Injury via Activation of Necroptosis in Diabetic Kidney Disease. *Oxid. Med. Cell. Longev.*, 2023:2713864.
- Ha C., Kim H., Cha R., Lee J., Lee S., Ryu J.H., Kim H., & Lee O.J. (2021). Effect



- of da-9701 on the gastrointestinal motility in the streptozotocin-induced diabetic mice. *J. Clin. Med.*, 10: 1-12.
- Harberson, J., Thomas, R.M., Harbison, S. P., & Parkman, H. P. (2010). Gastric neuromuscular pathology in gastroparesis: analysis of full-thickness antral biopsies. *Dig. Dis. Sci.*, 55(2):359-70.
- Honoré, S. M., Zelarayan, L. C., Genta, S. B., & Sánchez, S. S. (2011). Neuronal loss and abnormal BMP/Smad signaling in the myenteric plexus of diabetic rats. *Auton. Neurosci.*, 164(1–2): 51–61.
- Horváth, C., Jarabicová, I., Rajtík, T., Bartošová, L., Ferenczyová, K., Kaločayová, B., et al. (2023). Analysis of Signaling Pathways of Necroptotic and Pyroptotic Cell Death in the Hearts of Rats With Type 2 Diabetes Mellitus. *Physiol. Res.*, 72(S1): S23-S29.
- Hosseiniard, E. S., Morshedi, M., Bavafa-Valenlia, K., & Saghafi-Asl, M. (2019). The novel insight into anti-inflammatory and anxiolytic effects of psychobiotics in diabetic rats: possible link between gut microbiota and brain regions. *Eur J Nutr.*, 58(8):3361-3375.
- Ingaramo, P. I., Ronco, M. T., Francés, D. E., Monti, J. A., Pisani, G. B., Ceballos, M. P., et al. (2011). Tumor necrosis factor alpha pathways develops liver apoptosis in type 1 diabetes mellitus. *Mol. Immunol.*, (12-13):1397-407.
- International Diabetes Federation. (2019). *IDF Diabetes Atlas*. Ninth edit. Edited by S. Karuranga et al. Available at: www.diabetesatlas.org.
- Isic, A., Täng, M.S., Haugen, E., & Fu, M. (2008). TNF α -antagonist neither improve cardiac remodelling or cardiac function at early stage of heart failure in diabetic rats. *Autoimmunity*, 41: 473–477.
- Isola, R., Broccia, F., Casti, A., Loy, F., Isola, M., & Vargiu, R. (2021). STZ-diabetic rat heart maintains developed tension amplitude by increasing sarcomere length and crossbridge density. *Exp. Physiol.*, 106(7):1572-1586.
- Iwata, N., Takayama, H., Xuan, M., Kamiuchi, S., Matsuzaki, H., Okazaki, M., et al. (2015). Effects of Etanercept against Transient Cerebral Ischemia in Diabetic Rats. *Biomed. Res. Int.*, 2015:189292.
- Izbeki, F., Wittman, T., Rosztóczy, A., Linke, N., Bódi, N., Fekete, E., et al. (2008). Immediate insulin treatment prevents gut motility alterations and loss of nitrenergic neurons in the ileum and colon of rats with streptozotocin-induced



- diabetes. *Diabetes Res. Clin. Pract.*, 80: 192–198.
- Jain, N., Sudhakar, C., & Swarup, G. (2007). Tumor necrosis factor- α -induced caspase-1 gene expression. Role of p73. *FEBS. J.*, 274(17): 4396–4407.
- Jancsó, Z., Bódi, N., Borsos, B., Fekete, É., & Hermesz E. (2015). Gut region-specific accumulation of reactive oxygen species leads to regionally distinct activation of antioxidant and apoptotic marker molecules in rats with STZ-induced diabetes. *Int. J. Biochem. Cell. Biol.*, 62: 125–131.
- Joussen, A. M., Doecken, S., Le, M. L., Koizumi, K., Radetzky, S., Krohne, T. U., et al. (2009). TNF- α mediated apoptosis plays an important role in the development of early diabetic retinopathy and long-term histopathological alterations. *Mol. Vis.*, 15: 1418–1428.
- Karson, A., Utkan, T., Şahin, T.D., Balcı, F., Arkan, S., & Ateş, N. (2021) Etanercept rescues cognitive deficits, depression-like symptoms, and spike-wave discharge incidence in WAG/Rij rat model of absence epilepsy. *Epilepsy Behav.*, 115:1-7.
- Kern, T.S., Miller, C. M., Tang, J., Du, Y., Ball, S. L., & Berti-Matera L. (2010). Comparison of three strains of diabetic rats with respect to the rate at which retinopathy and tactile allodynia develop. *Mol. Vis.*, 16:1629-39.
- Klein, M., & Varga, I. (2020). Hirschsprung's Disease-Recent Understanding of Embryonic Aspects, Etiopathogenesis and Future Treatment Avenues. *Medicina*, 56(11):611.
- Kostouros, A., Koliarakis, I., Natsis, K., Spandidos, D. A., Tsatsakis, A., & Tsiaouassis, J. (2020). Large intestine embryogenesis: Molecular pathways and related disorders (Review). *Int. J. Mol. Med.*, 46(1): 27–57.
- Kuiken, N. S. S., Rings, E. H. H. M., Alffenaar, J. W. C., Havinga, R., Jurdzinski, A., Groen, A. K., & Tissing, W. J. E. (2017). Tumor Necrosis Factor-Alpha Inhibitor Etanercept Does Not Alter Methotrexate-Induced Gastrointestinal Mucositis in Rats. *J. Pediatr. Gastroenterol. Nutr.*, 65: e28–e34.
- Kulkarni, S., Micci, M. A., Leser, J., Shin, C., Tang, S. C., Fu, Y. Y., et al. (2017). Adult enteric nervous system in health is maintained by a dynamic balance between neuronal apoptosis and neurogenesis. *Proc. Natl. Acad. Sci. U S A.*, 114(18): E3709-E3718.
- Kuznik, E., Dudkowiak, R., Adamiec, R., & Poniewierka, E. (2020). Diabetic



- autonomic neuropathy of the gastrointestinal tract. *Prz. Gastroenterol.*, 15: 89–93.
- Kühn, E. R., Bellon, K., Huybrechts, L., & Heyns, W. (1983). Endocrine differences between the Wistar and Sprague-Dawley laboratory rat: influence of cold adaptation. *Horm. Metab. Res.*, 15(10):491-8.
- Kwak, J. M., Babygirija, R., Gribovskaja-Rupp, I., Takahashi, T., Yamato, S., & Ludwig, K. (2013). Regional difference in colonic motility response to electrical field stimulation in guinea pig. *J. Neurogastroenterol. Motil.*, 19(2): 192–203.
- Le Berre-Scoul, C., Chevalier, J., Oleynikova, E., Cossais, F., Talon, S., Neunlist, M., et al. (2017). A novel enteric neuron–glia coculture system reveals the role of glia in neuronal development. *J. Physiol.*, 595(2): 583–598.
- Lenzen, S. (2008). The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia*, 51: 216–226.
- Liao, S., Apaijai, N., Chattipakorn, N., & Chattipakorn, S. C. (2020). The possible roles of necroptosis during cerebral ischemia and ischemia / reperfusion injury. *Arch. Biochem. Biophys.*, 695: 1-19.
- Liu, T., Wei, H., Zhang, L., Ma, C., Wei, Y., Jiang, T., et al. (2024). Metformin attenuates lung ischemia-reperfusion injury and necroptosis through AMPK pathway in type 2 diabetic recipient rats. *BMC. Pulm. Med.*, 24(1):237.
- Lu, L., Lu, Q., Chen, W., Li, J., Li, C., Zheng, Z. (2018). Vitamin D₃ Protects against Diabetic Retinopathy by Inhibiting High-Glucose-Induced Activation of the ROS/TXNIP/NLRP3 Inflammasome Pathway. *J. Diabetes Res.*, 2018:1-11.
- Luckensmeyer, G. B., & Keast, J. R. (1998). Activation of α- and β-adrenoceptors by sympathetic nerve stimulation in the large intestine of the rat. *J. Physiol.*, 510(2): 549–561.
- Luo, P., Liu, D., Li, C., He, W. X., Zhang, C. L., & Chang, M. J. (2018). Enteric glial cell activation protects enteric neurons from damage due to diabetes in part via the promotion of neurotrophic factor release. *Neurogastroenterol. Motil.*, 30(10): 1–10.



- Madla, C. M., Qin, Y., Gavins, F. K. H., Liu, J., Dou, L., Orlu, M., *et al.* (2022). Sex Differences in Intestinal P-Glycoprotein Expression in Wistar versus Sprague Dawley Rats. *Pharmaceutics*. 14(5):1030.
- Magalhães, D. A., Kume, W. T., Correia, F. S., Queiroz, T. S., Allebrandt Neto, E. W., Santos, M. P. D., *et al.* (2019). High-fat diet and streptozotocin in the induction of type 2 diabetes mellitus: a new proposal. *An. Acad. Bras. Cienc.*, 91(1): 1 - 14.
- Majumder, S., Sreedhara, S. R., Banerjee, S., & Chatterjee, S. (2012). TNF a Signaling Beholds Thalidomide Saga: A Review of Mechanistic Role of TNF-a Signaling Under Thalidomide. *Curr. Top. Med. Chem.*, 12(13): 1456–1467.
- Mamun, A. A., Wu, Y., Nasrin, F., Akter, A., Taniya, M. A., Munir, F., *et al.* (2021). Role of pyroptosis in diabetes and its therapeutic implications. *J. Inflamm. Res.*, 14: 2187–2206.
- Manduca, A., Campolongo, P., Palmery, M., Vanderschuren, L. J., Cuomo, V., & Trezza, V. (2014). Social play behavior, ultrasonic vocalizations and their modulation by morphine and amphetamine in Wistar and Sprague-Dawley rats. *Psychopharmacology*, 231(8):1661-73.
- Markossian, S., & Kreydiyyeh, S. I. (2005). TNF-alpha down-regulates the Na⁺-K⁺ ATPase and the Na⁺-K⁺-2Cl⁻cotransporter in the rat colon via PGE2. *Cytokine*, 30(6):319-27.
- Medichem (2010). Glucose GOD/PAP: Enzymatic colorimetric test *Manual procedure*. www.medichem-me.com (diakses 28-09-2021).
- Meehan, C. J., Fleming, S., Smith, W., & Baird, J.D. (1994). Idiopathic megacolon in the BB rat. *Int. J. Clin. Exp. Pathol.*, 75(1): 37–42.
- Mescher, A. L. (2010). Junqueira's Basic Histology: Text & Atlas, 12th edition, McGraw-Hill company.
- Miranda, J., Eseberri, I., Lasa, A., & Portillo, M. P. (2018). Lipid metabolism in adipose tissue and liver from diet-induced obese rats: a comparison between Wistar and Sprague-Dawley strains. *J. Physiol. Biochem.*, 74(4):655-666.
- Moran, B. J. & Jackson, A. A. (1992). Function of the human colon. *Br. J. Surg.*,



79(11): 1132–1137.

- Nair, D. G., Miller, K. G., Lourensse, S. R., & Blennerhassett, M. G. (2014). Inflammatory cytokines promote growth of intestinal smooth muscle cells by induced expression of PDGF-R β . *J. Cell. Mol. Med.*, 18(3):444-54.
- Noorafshan, A., Yousefi, M., Hosseini, L., & Karbalay-Doust, S. (2019). Can Sertraline and Nortriptyline Protect the Neurons in Submucosal and Myenteric Plexuses of Rat's Colon Against Stress?. *Dig. Dis. Sci.*, 64(9): 2548–2554.
- Ojiro, K., Kitamura, H., Shimada, T., & Nakamura, M. (1993). A morphometrical study of the postnatal development of rat pancreatic islets, with special regard to the differences between Wistar and Sprague-Dawley strains. *Kaibogaku Zasshi.*, 68(2):190-203.
- Pandey, S., & Dvorakova, M. C. (2020). Future Perspective of Diabetic Animal Models. *Endocr. Metab. Immune. Disord. Drug. Targets*, 20(1): 25–38.
- Patel, B. A., Fidalgo, S., Wang, C., Parmar, L., Mandona, K., Panossian, A., et al. (2017). The TNF- α antagonist etanercept reverses age-related decreases in colonic SERT expression and faecal output in mice. *Sci. Rep.*, 7(42754): 1–12.
- Pei, Y., Wang, R., Chen, W., Yi, S., Huang, C., Liang S, et al. (2023). Impaired colonic motility in high-glycemic diet-induced diabetic mice is associated with disrupted gut microbiota and neuromuscular function. *Endocr. Connect.*, 12(9):1-11.
- Piper, M. S., & Saad, R. J. (2017). Diabetes Mellitus and the Colon. *Curr. Treat. Options. Gastroenterol.*, 15: 1–16.
- Qi, R., Yang, W., & Chen, J. (2013). Role of enteric glial cells in gastric motility in diabetic rats at different stages. *J. Huazhong. Univ. Sci. Technolog. Med. Sci.*, 33(4): 496–500.
- Rauf, A., Shah, M., Yellon, D. M., Davidson, S. M. (2019) Role of Caspase 1 in Ischemia/Reperfusion Injury of the Myocardium. *J. Cardiovasc. Pharmacol.*, 74(3):194-200.
- Rebolledo-solleiro, D. & Fernández-guasti, A. (2018). Influence of sex and estrous cycle on blood glucose levels, body weight gain, and depressive-like behavior



- in streptozotocin-induced diabetic rats. *Physiol. Behav.*, 1(194): 560–567.
- Rehman, G., Shehzad, A., Khan, A. L, & Hamayun, M. (2014). Role of AMP-activated protein kinase in cancer therapy. *Arch. Pharm. Chem. Life Sci.*, 347(7): 457-68.
- Roh, J. S., & Sohn, D. H. (2018). Damage-Associated Molecular Patterns in Inflammatory Diseases. *Immune. Netw.*, 18(4): 1–14.
- Rybnikova, E.A., Vetrovoi, O.V. & Zenko, M.Y. (2018). Comparative Characterization of Rat Strains (Wistar, Wistar-Kyoto, Sprague Dawley, Long Evans, LT, SHR, BD-IX) by Their Behavior, Hormonal Level and Antioxidant Status. *J. Evol. Biochem. Phys.*, 54: 374–382.
- Ruder, B., Atreya, R., & Becker C. (2019). Tumour necrosis factor alpha in intestinal homeostasis and gut related diseases. *Int. J. Mol. Sci.*, 20: 1-21.
- Saadane, A., Lessieur, E. M., Du, Y., Liu, H., & Kern, T. S. (2020). Successful induction of diabetes in mice demonstrates no gender difference in development of early diabetic retinopathy. *PLoS One*, 15(9):1–14.
- Sanders, L. M., Henderson, C. E., Hong, M. Y., Barhoumi, R., Burghardt, R. C., Carroll, R. J., et al. (2004). Pro-oxidant environment of the colon compared to the small intestine may contribute to greater cancer susceptibility. *Cancer Lett.*, 208(2): 155–161.
- Serban, A. I., Stanca, L., Geicu, O. I., Munteanu, M. C., & Dinischiotu, A. (2016). RAGE and TGF- β 1 cross-talk regulate extracellular matrix turnover and cytokine synthesis in AGes exposed fibroblast cells. *PLoS One*, 11: 1-19.
- Sha, H., Tong, X., & Zhao, J. (2018). Abnormal expressions of AGEs, TGF- β 1, BDNF and their receptors in diabetic rat colon—Associations with colonic morphometric and biomechanical remodeling. *Sci. Rep.*, 8(1): 1–14.
- Song, N. N., Lu, H. L., Lu, C., Tong, L., Huang, S. Q., Huang, X., et al. (2018). Diabetes-induced colonic slow transit mediated by the up-regulation of PDGFR α^+ cells/SK3 in streptozotocin-induced diabetic mice. *Neurogastroenterol. Motil.*, 30(8): 1-14
- Sun, Q., & Scott, M. J. (2016). Caspase-1 as a multifunctional inflammatory mediator: noncytokine maturation roles. *J. Leukoc. Biol.*, 100(5): 961–967.



- Sun, L., Ding, M., Chen, F., Zhu, D., & Xie, X. (2023). Brevicagipine alleviates podocyte injury by inhibiting NF- κ B/NLRP3-mediated pyroptosis in diabetic nephropathy. *PeerJ.*, 11:e14826.
- Szkudelski, T. (2012). Streptozotocin–nicotinamide-induced diabetes in the rat. Characteristics of the experimental model. *Exp. Biol. Med.*, 237(5): 481–490.
- Takahashi, T., & Owyang, C. (1998). Regional Differences in the Nitrergic Innervation Between the Proximal and the Distal Colon in Rats. *Gastroenterology*, 115(16): 1504–1512.
- Teixeira, A. R., Nitz, N., Guimaro, M. C., Gomes, C., & Santos-Buch, C. A. (2006). Chagas disease. *Postgrad. Med. J.*, 82(974):788-98..
- Tomasello, G., Mazzola, M., Leone, A., Sinagra, E., Zummo, G., Farina, F., et al. (2016). Nutrition, oxidative stress and intestinal dysbiosis: Influence of diet on gut microbiota in inflammatory bowel diseases. *Biomed. Pap. Med. Fac. Univ. Palacky. Olomouc. Czech. Repub.*, 160(4): 461–466.
- Uranga-Ocio, J. A., Bastús-Díez, S., Delkáder-Palacios, D., García-Cristóbal, N., Leal-García, M. Á., & Abalo-Delgado, R. (2015). Enteric neuropathy associated to diabetes mellitus. *Rev. Esp. de Enferm. Dig.*, 107(6): 366–373.
- Van Den Berge, N., Ferreira, N., Mikkelsen, T. W., Alstrup, A. K. O, Tamgüney, G., Karlsson P., et al. (2021). Ageing promotes pathological alpha-synuclein propagation and autonomic dysfunction in wild-type rats. *Brain*, 144: 1853–1868.
- Voss, U. & Ekblad, E. (2014). Lipopolysaccharide-induced loss of cultured rat myenteric neurons - role of AMP-activated protein kinase. *PLoS One*, 9(12): 1-17.
- Wang, X., Guo, Z., Ding, Z., Mehta, J. L. (2018). Inflammation, Autophagy, and Apoptosis After Myocardial Infarction. *J. Am. Heart. Assoc.*, 7(9): 1-13.
- Weerasinghe, P. & Buja, L. M. (2012). Oncosis: An important non-apoptotic mode of cell death. *Exp. Mol. Pathol.*, 93(3): 302–308.
- Wei, J., Zhao, Y., Liang, H., Du, W., & Wang L. (2022). Preliminary evidence for the presence of multiple forms of cell death in diabetes cardiomyopathy. *Acta Pharm. Sin. B.*, 12(1): 1–17.



- Wirth, R., Bódi, N., Maróti, G., Bagyánszki, M., Talapka, P., Fekete, É., *et al.* (2014). Regionally Distinct Alterations in the Composition of the Gut Microbiota in Rats with Streptozotocin-Induced Diabetes. *PloS One*, 9(12): 1–15.
- Wright, E., Bacon, J., & Glass, L. (2006). Oxidative stress in type 2 diabetes : the role of fasting and postprandial glycaemia. *Int. J. Clin. Pract.*, 60(3): 308–314.
- Xu, J., Chen, Y., Liu, S., & Hou, X. (2013). Electroacupuncture regulates apoptosis/proliferation of intramuscular interstitial cells of cajal and restores colonic motility in diabetic constipation rats. *Evid. Based. Complement. Alternat. Med.*, 2013: 1-10.
- Xie, D. P., Li, S., Li, L., Chang, X. W., Xi, T. F., Yang, X., *et al.* (2013). Beta-arrestin2 is involved in the increase of distal colonic contraction in diabetic rats. *Regul. Pept.*, 185:29-33.
- Ye, Q., Lin, Y. N., Xie, M. S., Yao, Y. H., Tang, S. M., Huang, Y., *et al.* (2019). Effects of etanercept on the apoptosis of ganglion cells and expression of Fas, TNF- α , caspase-8 in the retina of diabetic rats. *Int. J. Ophthalmol.*, 12(7): 1083–1088.
- Yu, T., Yang, H. S., Lu, X. J., Xia, Z. S., Ouyang, H., Shan, T. D., *et al.* (2016). Association of Bactericidal Dysfunction of Paneth Cells in Streptozocin-Induced Diabetic Mice with Insulin Deficiency. *Med. Sci. Monit.*, 22: 3062–3072.
- Zhao, M., Liao, D., & Zhao, J. (2017). Diabetes-induced mechanophysiological changes in the small intestine and colon. *World. J. Diabetes.*, 8(6):249-269.
- Zhao, S., Mysler, E. & Moots, R. J. (2018). Etanercept for the treatment of rheumatoid arthritis. *Immunotherapy*, 10(6): 433–445.
- Zhao, L., Bartnikas, T., Chu, X., Klein, J., Yun, C., Srinivasan, S., *et al.* (2019). Hyperglycemia promotes microvillus membrane expression of DMT1 in intestinal epithelial cells in a PKC α -dependent manner. *FASEB J.*, 33: 3549–3561.