

## DAFTAR PUSTAKA

- Akavipat, P., Thinkhamrop, J., Thinkhamrop, B., & Sriraj, W. (2016). Parameters Affecting Length of Stay Among Neurosurgical Patients in an Intensive Care Unit. *Acta Med. Indones.* 48 : 275–281.
- Amri, F. (2020). Correlation of Characteristics and Nutritional Status on Length Of Stay of ICU Patients in Purwakarta. *J. Appl. Food Nutr.* 1 : 17–23.
- Badenes, R., Prisco, L., Maruenda, A., & Taccone, F.S. (2017). Criteria for Intensive Care admission and monitoring after elective craniotomy. *Curr. Opin. Anaesthesiol.* 30 : 540–545.
- Bangun, C.G., Fuadi, I., & Suarjaya, I. putu P.S. (2023). Neuroanesthesia and Critical Care: Compos mentis, Delirium, Koma. Bandung : INA-SNACC.
- Basali, A., Mascha, E.J., Kalfas, L., & Schubert, A. (2000). Relation between perioperative hypertension and intracranial hemorrhage after craniotomy. *Anesthesiology.*
- Beangklang, R., Sitthinamsuwan, B., Tansirisithikul, C., & Nunta-aree, S. (2024). Hydrocephalus following Brain Tumor Surgery: Factors Correlating with Occurrence of Postoperative Hydrocephalus and Predictive Scoring Model. *Asian J. Neurosurg.*
- Beauregard, C.L., & Friedman, W.A. (2003). Routine use of postoperative ICU care for elective craniotomy: A cost-benefit analysis. *Surg. Neurol.* 60 : 483–489.
- Bethesda (2023). Childhood Central Nervous System Atypical Teratoid/Rhabdoid Tumor Treatment, in: *Cancer Information*. US : National Cancer Institute.
- Bisri, D.Y., & Bisri, T. (2016). No Title, in: *Anestesi Untuk Operasi Tumor Otak: Supratentorial Infratentorial*. pp. 41–90, Bandung : Fakultas Kedokteran Universitas Padjajaran.
- Chauhan, R., Bloria, S.D., & Luthra, A. (2019). Management of Postoperative Neurosurgical Patients. *Indian J. Neurosurg.* 08 : 179–184.
- Cinar, Z., Ozturk, B., & Aydogdu, M. (2022). The Impact of Body Mass Index on Intensive Care Unit Outcomes of Critically ill Patients. *J. Crit. Intensive Care* 13 : 97–101.
- Diabetes, D.O.F. (2010). Diagnosis and classification of diabetes mellitus. *Diabetes Care* 33.
- Fatoni, A.Z., & Kestriani, N.D. (2018). Acute Kidney Injury (AKI) pada Pasien Kritis. *Anesth. Crit. Care* 36 : 64–76.
- Fernandez de Thomas, R.J., Munakomi, S., & Jesus, O. (2023). Craniotomy. Treasure Island : Statpearls.
- Florman, J.E., Cushing, D., Keller, L.A., & Rughani, A.I. (2017). A protocol for postoperative admission of elective craniotomy patients to a non-ICU or step-down setting. *J. Neurosurg.* 127 : 1392–1397.
- Hannon, M.J., Finucane, F.M., Sherlock, M., Agha, A., & Thompson, C.J. (2012). Disorders of water homeostasis in neurosurgical patients. *J. Clin. Endocrinol. Metab.* 97 : 1423–1433.
- Hendarjana, P. (2013). Penatalaksanaan Infeksi Pada Penderita Penyakit Kritis : Pneumonia. Jakarta : PERDICI.
- Huang, Y.C., Huang, S.J., Tsauo, J.Y., & Ko, W.J. (2010). Definition, risk factors and outcome of prolonged surgical intensive care unit stay. *Anaesth. Intensive Care* 38 : 500–505.
- Joseph, A.C., MS, K., R, S., R, V., Johson, S., & Chander, U. (2016). JNC 8 versus JNC 7 Understanding the Evidences. *Int. J. Pharm. Sci. Rev. Res.* 36 : 38–43.



- Kaki, A., & Khalil, H. (2021). iMedPub Journals Predictors of Prolonged Intensive Care Unit Length of Stay among Surgical Patients : A Scoping Review Abstract 1–17.
- Kapadinhos, T., Angelopoulos, E., Vasileiadis, I., Nanas, S., Kotanidou, A., Karabinis, A., *et al.* (2017). Determinants of prolonged intensive care unit stay in patients after cardiac surgery: A prospective observational study. *J. Thorac. Dis.* 9 : 70–79.
- Kayaalti, S., & Kayaalti, O. (2019). Risk factors affecting the length of intensive care unit stay after brain tumor surgery. *Indian Anaesth. Forum* 20 : 61.
- Keown, T., Bhangu, S., & Solanki, S. (2022). Anaesthesia for Craniotomy and Brain Tumour Resection. *Anesth. Tutor. Week* 1–7.
- Kumwilaisak, K., Kyokong, O., & Indrambarya, T. (2008). Factors influencing length of stay in neurosurgical intensive care unit. *J. Med. Assoc. Thai.* 91 : 875–881.
- Lan, J., Wei, Y., Zhu, Y., Zhang, Y., Zhang, S., Mo, L., *et al.* (2023). Risk Factors for Post-Operative Pulmonary Infection in Patients with Brain Tumors: A Systematic Review and Meta-Analysis. *Surg. Infect. (Larchmt)*. 24 : 588–597.
- Lee, S.Y., Lee, S.H., Tan, J.H.H., Foo, H.S.L., Phan, P.H., Kow, A.W.C., *et al.* (2018). Factors associated with prolonged length of stay for elective hepatobiliary and neurosurgery patients: a retrospective medical record review. *BMC Health Serv. Res.* 18 : 5.
- Lee, Y.I., Ahn, J., & Ryu, J.A. (2022). Clinical Outcomes Associated with Degree of Hyponatremia in Neurocritically Ill Patients. *J. Korean Neurosurg. Soc.* 66 : 95–104.
- Manninen, P.H., Raman, S.K., Boyle, K., & El-Beheiry, H. (1999). Early postoperative complications following neurosurgical procedures. *Can. J. Anaesth.* 46 : 7–14.
- Momin, E.N., Adams, H., Shinohara, R.T., Frangakis, C., Brem, H., & Quiñones-Hinojosa, A. (2012). Postoperative mortality after surgery for brain tumors by patient insurance status in the United States. *Arch. Surg.*
- Paisansathan, C., & Ozcan, M.S. (2023). Anesthesia for craniotomy in adults. *UpToDate*.
- Phillips, K.R., Enriquez-Marulanda, A., Mackel, C., Ogbonna, J., Moore, J.M., Vega, R.A., *et al.* (2023). Predictors of extended length of stay related to craniotomy for tumor resection. *World Neurosurg.* X 19 : 100176.
- Popugaev, Konstantin A. Lubnin, A.Y. (2015). Postoperative Care in Neurooncology, in: *Neurointensive Care*. pp. 95–124, New York : Springer US.
- Pribadi, H.T. (2012). Angka Kematian Pasien Kraniotomi Di Icu Dan Hcu Rsup Dr. Kariadi Jurnal Media Medika Muda Karya Tulis Ilmiah. *J. Media Med. Muda*.
- Rajagopalan, V., Chouhan, R.S., Pandia, M.P., Lamsal, R., & Rath, G.P. (2019). Effect of intraoperative blood loss on perioperative complications and neurological outcome in adult patients undergoing elective brain tumor surgery. *J. Neurosci. Rural Pract.* 10 : 631–640.
- Rhondali, O., Genty, C., Halle, C., Gardellin, M., Ollinet, C., Oddoux, M., *et al.* (2011). Do patients still require admission to an intensive care unit after elective craniotomy for brain surgery? *J. Neurosurg. Anesthesiol.* 23 : 118–123.
- Sangtongjaraskul, S., Lertsirisothon, S., Sae-Phua, V., Kanta, S., & Kongkiattikul, L. (2023). Factors Influencing Prolonged Intensive Care Unit Length of Stay after Craniotomy for Intracranial Tumor in Children: A 10-year Analysis from A University Hospital. *Indian J. Crit. Care Med.* 27 : 205–211.
- Sastry, R.A., Pertsch, N.J., Tang, O., Shao, B., Toms, S.A., & Weil, R.J. (2020). Frailty and outcomes after craniotomy for brain tumor. *J. Clin. Neurosci.* 81 : 95–100.
- Savareh, B.A., Alibabaei, A., Ahmady, S., Mokhtari, M., Hajiesmaeili, M., & Nateghinia, S.

- (2021). Predicting patient length of stay in a neurosurgical intensive care unit of a large teaching hospital. *J. Cell. Mol. Anesth.* 6 : 132–140.
- Solheim, O., Jakola, A.S., Gulati, S., & Johannesen, T.B. (2012). Incidence and causes of perioperative mortality after primary surgery for intracranial tumors: a national, population-based study. *J. Neurosurg.* 116 : 825–834.
- Solomon Cardona, M.T., Roig Fabre, E., Pila Bandera, M., Domínguez, W., & Crombet Ramos, T. (2015). Mortality from malignant brain tumors within the first 30 days after surgery. *Rev. Cuba. Cir.* 53 : 244–255.
- Sopiyudin, D.M. (2013). *Besar Sampel dan Cara Pengambilan Sampel*. Jakarta : Salemba Medika.
- Uribe, A.A., Stoicea, N., Echeverria-Villalobos, M., Todeschini, A.B., Esparza Gutierrez, A., Folea, A.R., *et al.* (2021). Postoperative Nausea and Vomiting after Craniotomy: An Evidence-based Review of General Considerations, Risk Factors, and Management. *J. Neurosurg. Anesthesiol.* 33 : 212–220.
- Widodo, D., Andriani, R., & Haq, I.B.I. (2019). Pedoman nasional pelayanan kedokteran tumor otak. *Kom. Penanggulangan Kanker Nas.* 1–206.
- Zhang, Y., Tan, H., Jia, L., He, J., Hao, P., Li, T., *et al.* (2023). Association of preoperative glucose concentration with mortality in patients undergoing craniotomy for brain tumor. *J. Neurosurg.* 138 : 1254–1262.