

DAFTAR PUSTAKA

- Adembesa, I., Mung'ayi, V., Premji, Z., Kanya, D. A randomized control trial comparing train of four ratio > 0.9 to clinical assessment of return of neuromuscular function before endotracheal extubation on critical respiratory events in adult patients undergoing elective surgery at a tertiary hospital in Nairobi. *African Health Sciences*, East Africa. 2018.
- Agljo, LS., Lekowski, RW., Urman RD., *Essential Clinical Anesthesia Review*. Cambridge Medicine, United Kingdom. 2015.
- Baillard C, Gehan G, Reboul-Marty J, Larmignat P, Samama CM, Cupa M. Residual Curarization in the Recovery Room After General Anesthesia With Vecuronium. *Anesthesiology*. 2000 Dec;93(6):1312-8. doi: 10.1097/00000542-200012000-00003.
- Baurain, MJ., Hoton, F., D'Hollander, AA., Cantraine, FR., 1996. Is recovery of neuromuscular transmission complete after the use of neostigmine to antagonize block produced by rokuronium, vecuronium, atracurium and pancuronium. *Br. J. Anaesth* 77, 496-499.
- Boles, JM., Bion, J., Connors, A., Herridge, M., Marsh, B., Melot, C., Pearl, R., Silverman, H., Stanchina, M., Viellard-Baron, A., Welte, T. *Weaning from Mechanical Ventilation*. France. *European Respiratory Journal*. 2007;1041.
- Butterly, A., Bittner, EA., George, E., Sandberg, WS., Eikermann, M., Schmidt, U. Postoperative residual curarization from intermediate-acting neuromuscular blocking agents delays recovery room discharge. *Br J Anaesth*. 2010;105.
- Butterworth, JF., Mackey, DC., Wasnick, JD. Morgan & Mikhail's Clinical Anesthesiology. 6th ed. United States: *McGraw-Hill Education*. 2018;213.
- Brull, SJ., Murphy, GS. Residual neuromuscular Block: Lessons Unlearned. Part II: Metode to reduce the risk of residual weakness. *International Anesthesia Research Society*. Florida. 2010. Vol I. 129-131.
- Brull, SJ., Kopman, AF., Current Status of Neuromuscular Reversal and Monitoring Challenges and Opportunities. *Anesthesiol J Am Soc Anesthesiol*. 2017;126(1):173-190.

- Brown, CA., Sakles, JC., Mick, NW. The Walls Manual of Emergency Airway Management. 5th ed. *Wolters Kluwer*. Philadelphia. 2018; 642-662.
- Cook, T., Kristensen, MS. Core Topics in Airway Management. 3rd ed. *Cambridge University Press*. United Kingdom. 2021. 177-184.
- Debaene, B., Plaud, B., Dilly, MP., Donati, F. Residual Paralysis in the PACU After a Single Intubating Dose of Nondepolarizing Muscle Relaxant With an Intermediate Duration of Action. *Anesthesiology*. 2003 Aug;99(2):421-7. doi: 10.1097/00000542-200308000-00025.
- Douglas, NJ., White, DP., Pickett, CK., Weil, JV., Clifford, WZ. *Respiration during sleep in normal man*. Cardiovascular Pulmonary Research Laboratory. Colorado. 1982;37:840-844.
- Emyedu, A., Atumanya, P., Okello, E., Wabule, A., Ssemogerere, L., Mukisa, J., Sendagire, C. Prevalence and Complications of Residual Neuromuscular Blockade in the Post Anesthesia Care Units of Uganda. *Journal of Anesthesia and Clinical Research*, Uganda. 2022; 1-2.
- Errando, CL., Garutti, I., Mazzinari, G., Diaz-Cambronero, O., Bebawy, JF. Grupo Español De Estudio Del Bloqueo Neuromuscular Residual. Neuromuscular blockade in the postanesthesia care unit: observational cross-sectional study of a multicenter cohort. *Minerva Anesthesiology*. 2016;82.
- Flood, P., Rathmell, JP., Shafer, S. *Stoelting's Pharmacology & Physiology in Anesthetic Practice*. *Wolters Kluwer*, United States. 2015.
- Foldes FF, Nagashima H, Nguyen HD, Schiller WS. Pharmacokinetic and Pharmacodynamic Aspects of Neuromuscular Blockade. *Anesthesiology*. 1976 Sep;45(3):309-22. doi: 10.1097/00000542-197609000-00011.
- Fu, ES., Downs, JB., Schweiger, JW., Miguel, RV., Smith, RA. Supplemental Oxygen Impairs Detection of Hypoventilation by Pulse Oximetry. *Chest*. Florida. 2004.
- Fuchs-Buder, T., Claudius, C., Skovgaard, L. T., Eriksson, L. I., & Mirakhor, R. K. (2016). Good clinical research practice in pharmacodynamic studies of neuromuscular blocking agents II: The Stockholm revision. *Acta Anaesthesiologica Scandinavica*, 61(2), 101-121.

- Grayling, M., Sweeney, BP. Recovery from neuromuscular blockade: a survey of practice. *The Association of Anaesthesia of Great Britain and Ireland*. United Kingdom. 2007.
- Hagberg, CA., Artime, CA., Aziz, MF. Hagberg and Benumof's Airway Management. 4th ed. *Elsevier*. Philadelphia. 2018; 178-208; 844-867.
- Hahn, S. Understanding noninferiority trials. *Korean J Pediatr*. 2012. 55(11), 403-407.
- Heidegger, T. Management of Difficult Airway. *The New England Journal of medicine*. 2021; 2-10.
- Hemmerling, T. M., Schmidt, J., Boselli, E., Le, N., & Briand, M. (2013). Train-of-Four Monitoring in the Evaluation of Neuromuscular Blockade Recovery: Clinical Implications. *Journal of Clinical Monitoring and Computing*, 27(2), 107-112.
- Herrera L.D., Ostroff, R., Rogers, S.A., Sevoflurane: Approaching the Ideal Inhalation Anesthetic A Pharmacologic, Pharmacoeconomic and Clinical Review. *CNS drug Review*. 2001; 48-120.
- Hsieh, C.Y., Poon, Y.Y., Ke, T.Y., Chiang, M.H., Li, Y.Y., Tsai, P.N., Wu, S.C. Post operative Vomiting Following Laparoscopic Cholecystectomy is Association with Intraoperative Fluid Administration: A Retrospective Cohort Study. *International Journal of Enviromental Research and Public Health*. Taiwan. 2021; 1-3.
- Hunter, JM. Reversal of residual neuromuscular block: complications associated with perioperative management of muscle relaxation. *British Journal of Anaesthesia*. 2017; 119.
- JOINT TRAUMA SYSTEM CLINICAL PRACTICE GUIDELINE (JTS CPG) *Mechanical Ventilation Basics* (CPG ID: 92). (n.d.). Available at: https://jts.health.mil/assets/docs/cpgs/Mechanical_Ventilation_Basics_27_Dec_2021_ID92.pdf.
- Jubb, A., Ford, P. Extubation after Anaesthesia: A Systematic Review. *Update in Anesthesia*. United Kingdom. 2009; 30-36.

- Kajal, S., Dhankhar, M., Mukherjee, M., et al., 2016. Evaluation of Low Doses of Neostigmine for Reversal of Residual Neuromuscular Blockade. *J Anesth Crit Care*. Open Access. 4(3), 00137.
- Kim, J. S., Ahn, H. J., Kim, H. J., & Choi, J. W. (2015). Clinical Assessment of Neuromuscular Function during Anesthesia and Its Impact on Extubation Outcomes. *Journal of Clinical Anesthesia*, 27(4), 331-338.
- Lee, S., Ro, YJ., Koh, WU., Nishiyama, T., Yang, HS. The neuromuscular effect of rocuronium under sevoflurane-remifentanil or propofol-remifentanil anesthesia: a randomized clinical comparative study in an Asia population. *BMC Anesthesiology*, Seoul. 2016.
- Luo, J., Chen, S., Min, S., Peng, L. Reevaluation and update on efficacy and safety of neostigmine for reversal of neuromuscular blockade. *Dovepress*, China. 2018:14. 2239-2406.
- Mathias, LAST. Bernardis, RCG. Postoperative Residual Paralysis. *Brazilian Society of Anesthesiology*, Brazil. 2012;1.
- Miller RD, ed. *Miller's Anesthesia*. Eighth edition. Elsevier/Saunders; 2015.
- Murphy, GS., Szokol, JW., Marymont, JH., Greenberg, SB., Avram, MJ., Vender, JS. Residual Neuromuscular Blockade and Critical Respiratory Events in the Postanesthesia Care Unit: *Anesth Analg*. 2008;130-137.
- Murphy GS, Szokol JW, Avram MJ, Greenberg SB, Marymont JH, Vender JS, Gray J, Landry E, Gupta DK. Intraoperative acceleromyography monitoring reduces symptoms of muscle weakness and improves quality of recovery in the early postoperative period. *Anesthesiology*. 2011; 115:946–54.
- Murphy, GS., Brull, SJ. Quantitative Neuromuscular Monitoring and Postoperative Outcomes: A Narrative Review. *The American society of Anesthesiologists*. Chicago. 2021; 348-349.
- Naguib M. Neuromuscular Monitoring and Postoperative Respiratory Function. *Anesthesiology*. 2018 Mar;128(3):700-709. doi: 10.1097/ALN.0000000000001971.
- Renew, JR. Comparison of Visual and Electromyograph Assessments in Response to Train-of-Four Stimulation of The Ulnar Nerve. Florida. 2021;7.

- Parotto, M., Cooper, RM., Behringer, EC. Extubation of the Challenging or Difficult Airway. *Springer*. 2020;2-6.
- Plaud, B., Debaene, B., Donati, F., Marty, J. Residual Paralysis after Emergence from Anesthesia. *the American Society of Anesthesiologist*. 2010; 112: 1013-22.
- Xara, D., Santos, A., Abelha, F. Adverse respiratory events in a post- anesthesia care unit. *Archivos de Bronconeumología (English Edition)*. 2015; 51.
- Takagi, S., Kitajima, O., Yamamoto, M., Matsui, M., Suzuki, T., Potentiation by sevoflurane of rocuronium-induced neuromuscular block is greater in older than younger adult patients: a randomised controlled trial. *British Journal of Anaesthesia*.2020.
- Thilen, SR., Weigel, WA., Todd, MM., Dutton, RP., Lien, CA., Grant, SA., Szokol, JW., Eriksson, LI., Yaster, M., Gran, MD., Agarkar, M., Marbella, AM., Blanck, JF., Domino, KB. 2023 American Society of Anesthesiologist Practice Guidelines for Monitoring and Antagonism of Neuromuscular Blockade: A Report by the American Society of Anesthesiologist Task Force on Neuromuscular Blokade. *Anesthesiology*. 2022. V;138: 13-41.
- Thompson, J., Moppett, I., Wiles, M. Smith and Aitkenhead's Textbook of Anesthesia. 7th ed. *Elsevier*. Poland. 2019; 456-506.
- Walker, E., an Nowacky, AS.Understanding Equivalence and Noninferiority Testing. *J Gen Intern Med*. 2010. 26(2), 192–6.
- Wardhana, A., Kurniawaty, J., Uyun, Y. Optimised reversal without train- of-four monitoring versus reversal using quantitative train-of-four monitoring: An equivalence study. *Indian Society of Anaesthesiologists, Indonesia*. 2019;63: 361-7.
- Yang, L., Hu, N., Chang, H., Yang, D., Zuo, Y. Routine administration if neostigmine after recovery of spontaneous breathing versus neuromuscular monitor-guided administration of neostigmine in pediatric patients: a parallel, randomized, controlled study. *BMC. West China*. 2023; 24:19.
- Zhong B. How to calculate sample size in randomized controlled trial? *J Thorac Dis*. 2009;1(1):51-54.