

LIST OF CONTENTS

| | Page |
|--|------|
| TITLE PAGE | i |
| AUTHORIZATION | ii |
| PLAGIARISM-FREE STATEMENT | iii |
| ACKNOWLEDGMENT | iv |
| LIST OF CONTENTS | v |
| LIST OF TABLES | vii |
| LIST OF FIGURE | viii |
| ABSTRACT | ix |
| INTISARI | x |
| | |
| I. INTRODUCTION | 1 |
| 1.1. Background | 1 |
| 1.2. Problem formulation | 2 |
| 1.3. Objectives | 2 |
| 1.4. Advantages of research | 3 |
| 1.5. Originality | 3 |
| | |
| II. LITERATURE STUDY | 4 |
| 2.1. Rice farming | 4 |
| 2.2. Nitrogen cycling in paddy field | 5 |
| 2.3. Inceptisol | 7 |
| 2.4. Weed management | 7 |
| 2.5. Haenuki and Sasanishiki | 8 |
| 2.6. Theoretical Basic | 9 |
| 2.7. Hypothesis | 10 |
| | |
| III. MATERIAL AND METHODS | 11 |
| 3.1. Site description and preparation | 11 |
| 3.2. Plant sampling and analysis | 12 |
| 3.3. Soil Sampling and analysis | 13 |
| 1.1.1. Analysis of pH, EC, NH_4^+ , and NO_3^- | 13 |
| 1.1.2. Analysis of Soil Organic Carbon (SOC) and Total Nitrogen (TN) | 13 |
| 1.1.3. Analysis min-N or available N (Anaerobic Incubation) | 13 |
| 1.1.4. Analysis of bulk density | 14 |
| 1.1.5. Analysis of soil texture | 14 |
| 3.4. Data Calculations and Statistical Analysis | 14 |
| | |
| IV. RESULT AND DISCUSSION | 15 |
| 4.1. Soil properties | 15 |
| 4.2. NO_3^- and NH_4^+ on different soil fertility | 16 |
| 4.3. Effect of soil fertility and weeding management on rice and weeds biomass | 19 |
| 4.4. Weed diversity and percentage | 21 |



| | |
|--|----|
| 4.5. Effect of soil fertility and weeding management on nitrogen in rice and weeds | 22 |
| 4.6. Effect of soil fertility and weeding management on grain yield and its components | 27 |
| V. CONCLUSIONS | 32 |
| REFERENCES | 33 |
| APPENDIX | 40 |

LIST OF TABLES

| | Page |
|--|------|
| Table 1. Climate condition in Tsuruoka, Yamagata, Japan. | 11 |
| Table 2. Initial soil properties of high fertility (HF) and low fertility (LF) paddy fields in Tsuruoka, Yamagata, Japan. | 15 |
| Table 3. N concentration of rice and weeds among the soil fertility, weeding management, and rice variety treatment at 72 and 114 days after transplanting (DAT). | 23 |
| Table 4. N uptake of rice and weeds among the soil fertility, weeding management, and rice variety treatment at 72 and 114 days after transplanting (DAT). | 24 |
| Table 5. Grain yield and its components (panicle number, spikelet per panicle, % of filled spikelet, and grain weight). | 28 |
| Table 6. Relationship among plant biomass, N concentration, N uptake, rice grain yield, and its components. | 46 |

LIST OF FIGURE

| | Page |
|---|------|
| Figure 1. Study area map of both paddy field in Tsuruoka, Yamagata, Japan..... | 11 |
| Figure 2. Research design of both high fertility (HF) and low fertility (LF) paddy fields in Tsuruoka, Yamagata, Japan..... | 12 |
| Figure 3. Ammonium (NH_4^+) (a) and Nitrate (NO_3^-) (b) at 25, 47, 66, 95 and 108 days after transplanting (DAT)..... | 17 |
| Figure 4. Rice biomass (contribution in root, stem, leaf, and ear) and weeds biomass (contribution in various species) at 72 and 114 days after transplanting (DAT) | 19 |
| Figure 5. Weed biomass percentage of various species at 72 and 114 days after transplanting (DAT). | 21 |