

DAFTAR PUSTAKA

- Adhyaru, D. N., Bhatt, N. S., and Modi, H. A. (2015). Optimization of upstream and downstream process parameters for cellulase-poor-thermo-solvent-stable xylanase production and extraction by *Aspergillus tubingensis* FDHN1. *Bioresources and Bioprocessing*, 2(1), 1–14. <https://doi.org/10.1186/s40643-014-0029-1>
- Adnan, M., Zheng, W., Islam, W., Arif, M., Abubakar, Y. S., Wang, Z., and Lu, G. (2018). Carbon catabolite repression in filamentous Fungi. *International Journal of Molecular Sciences*, 19(1), 1–23. <https://doi.org/10.3390/ijms19010048>
- Afriyanti, S., and Setyabudi, S. (2014). *Microbial conversion of cassava stem (Manihot esculenta) cellulose into reducing sugar by Trichoderma reesei PK1J2*. <http://ejurnal.unisri.ac.id/index.php/prodfatipa/article/view/939/791>
- Aguilar, C. N., Augur, C., Favela-Torres, E., and Viniestra-González, G. (2001). Production of tannase by *Aspergillus niger* Aa-20 in submerged and solid-state fermentation: Influence of glucose and tannic acid. *Journal of Industrial Microbiology and Biotechnology*, 26(5), 296–302. <https://doi.org/10.1038/sj.jim.7000132>
- Ahmed, S., Riaz, S., and Jamil, A. (2009). Molecular cloning of fungal xylanases: An overview. *Applied Microbiology and Biotechnology*, 84(1), 19–35. <https://doi.org/10.1007/s00253-009-2079-4>
- Ajjolakewu, A. K., Peng, C., Nadiah, W., and Abdullah, W. (2017). Biocatalysis and agricultural biotechnology optimization of production conditions for xylanase production by newly isolated strain *Aspergillus niger* through solid state fermentation of oil palm empty fruit bunches. *Biocatalysis and Agricultural Biotechnology*, 11(October 2016), 239–247. <https://doi.org/10.1016/j.bcab.2017.07.009>
- Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., and Walter, P. (2002). From DNA to RNA. In *Molecular Biology of the Cell*. 4th edition. (pp. 1–19). Garland Science. <https://www.ncbi.nlm.nih.gov/books/NBK26887/>
- Álvarez-Cervantes J2, Domínguez-Hernández EM3, Mercado-Flores Y2, O. A. and D.-G. G. (2016). Mycosphere Essay 10: Properties and characteristics of microbial xylanases. *Mycosphere*, 7(December), 1600–1619. <https://doi.org/10.5943/mycosphere/si/3b/12>
- Andrade, S. D. V., Lourdes, M. De, Moraes, T. De, Terenzi, H. F., and At, J. (2004). Effect of carbon source on the biochemical properties of β -xylosidases produced by *Aspergillus versicolor*. *Process Biochemistry*, 39, 1931–1938. <https://doi.org/10.1016/j.procbio.2003.09.024>
- Andrews, S. (2010). *FastQC: A Quality Control Tool for High Throughput Sequence Data*.
- Aro, N., Ilmén, M., Saloheimo, A., and Penttilä, M. (2003). ACEI of *Trichoderma reesei* is a repressor of cellulase and xylanase expression. *Applied and Environmental Microbiology*, 69(1), 56–65. <https://doi.org/10.1128/AEM.69.1.56-65.2003>

- Barboráková, Z., Labuda, R., Häubl, G., and Tančinova, D. (2012). Effect of glucose concentration and growth conditions on the fungal biomass, pH of media and production of Fumagillin by a non-pathogenic strain *Penicillium scabrosum*. *Journal of Microbiology, Biotechnology and Food Science*, 1(4), 466–476.
- Bhardwaj N., Kumar B., Verma P. (2019). A detailed overview of xylanases: an emerging biomolecule for current and future prospective. *Bioresour. Bioprocess*, 6(36).
- Beg, Q. K., Kapoor, M., Mahajan, L., and Hoondal, G. S. (2001). Microbial xylanases and their industrial applications: A review. *Applied Microbiology and Biotechnology*, 56(3–4), 326–338. <https://doi.org/10.1007/s002530100704>
- Benassi, V. M., de Lucas, R. C., Jorge, J. A., and Moraes, M. de L. T. de. (2014). Screening of thermotolerant and thermophilic fungi aiming β -xylosidase and arabinanase production. *Brazilian Journal of Microbiology*, 45(4), 1459–1467.
- Borin, G. P., Sanchez, C. C., de Santana, E. S., Zanini, G. K., dos Santos, R. A. C., de Oliveira Pontes, A., de Souza, A. T., Dal'Mas, R. M. M. T. S., Riaño-Pachón, D. M., Goldman, G. H., and Oliveira, J. V. de C. (2017). Comparative transcriptome analysis reveals different strategies for degradation of steam-exploded sugarcane bagasse by *Aspergillus niger* and *Trichoderma reesei*. *BMC Genomics*, 18(1), 1–21. <https://doi.org/10.1186/s12864-017-3857-5>
- Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*, 34(5), 525–527. <https://doi.org/10.1038/nbt.3519>
- Brown, N. A., Ries, L. N. A., Reis, T. F., Rajendran, R., Corrêa Dos Santos, R. A., Ramage, G., Riaño-Pachón, D. M., and Goldman, G. H. (2016). RNAseq reveals hydrophobins that are involved in the adaptation of *Aspergillus nidulans* to lignocellulose. *Biotechnology for Biofuels*, 9(1), 1–17. <https://doi.org/10.1186/s13068-016-0558-2>
- Butt, M. S., Tahir-Nadeem, M., Ahmad, Z., and Sultan, M. T. (2008). Xylanases and their applications in baking industry. *Food Technology and Biotechnology*, 46(1), 22–31.
- Carrillo-Sancen, G., Carrasco-Navarro, U., Tomasini-Campocoso, A., Corzo, G., Pedraza-Escalona, M. M., and Favela-Torres, E. (2016). Effect of glucose as a carbon repressor on the extracellular proteome of *Aspergillus niger* during the production of amylases by solid state cultivation. *Process Biochemistry*, 51(12), 2001–2010. <https://doi.org/10.1016/j.procbio.2016.09.001>
- Castro, L. D. S., Antoniêto, A. C. C., Pedersoli, W. R., Silva-Rocha, R., Persinoti, G. F., and Silva, R. N. (2014). Expression pattern of cellulolytic and xylanolytic genes regulated by transcriptional factors XYR1 and CRE1 are affected by carbon source in *Trichoderma reesei*. *Gene Expression Patterns*, 14(2), 88–95. <https://doi.org/10.1016/j.gep.2014.01.003>
- Cepeda-García, C., Domínguez-Santos, R., García-Rico, R. O., García-Estrada, C., Cajiao, A., Fierro, F., and Martín, J. F. (2014). Direct involvement of the CreA transcription factor in penicillin biosynthesis and expression of the pcbAB

- gene in *Penicillium chrysogenum*. *Applied Microbiology and Biotechnology*, 98(16), 7113–7124. <https://doi.org/10.1007/s00253-014-5760-1>
- Chand, P., Aruna, A., Maqsood, A. M., and Rao, L. V. (2005). Novel mutation method for increased cellulase production. *Journal of Applied Microbiology*, 98(2), 318–323. <https://doi.org/10.1111/j.1365-2672.2004.02453.x>
- Chapla, D., Pandit, P., and Shah, A. (2012). Production of xylooligosaccharides from corncob xylan by fungal xylanase and their utilization by probiotics. *Bioresource Technology*, 115, 215–221. <https://doi.org/10.1016/j.biortech.2011.10.083>
- Chilton, I. J., Delaney, C. E., Barham-Morris, J., Fincham, D. A., Hooley, P., and Whitehead, M. P. (2008). The *Aspergillus nidulans* stress response transcription factor StzA is ascomycete-specific and shows species-specific polymorphisms in the C-terminal region. *Mycological Research*, 112(12), 1435–1446. <https://doi.org/10.1016/j.mycres.2008.06.028>
- Coradetti, S. T., Craig, J. P., Xiong, Y., Shock, T., Tian, C., and Glass, N. L. (2012). Conserved and essential transcription factors for cellulase gene expression in ascomycete fungi. *Proceedings of the National Academy of Sciences of the United States of America*, 109(19), 7397–7402. <https://doi.org/10.1073/pnas.1200785109>
- Cortés-maldonado, L., MarciaL-quino, J., Gómez-manzo, S., and Fierro, F. (2020). A method for the extraction of high quality fungal RNA suitable for RNA-seq. *Journal of Microbiological Methods*, 170(105855), 1–8. <https://doi.org/10.1016/j.mimet.2020.105855>
- Costa, M. A. L., Farinas, C. S., and Miranda, E. A. (2018). Ethanol precipitation as a downstream processing step for concentration of xylanases produced by submerged and solid-state fermentation. *Brazilian Journal of Chemical Engineering*, 35(2), 477–488. <https://doi.org/10.1590/0104-6632.20180352s20160502>
- Crick, F. (1970). Centra Dogma of Molecular Biology. In *Nature Publishing Group* (Vol. 227, pp. 561–562). <http://www.mendeley.com/research/discreteness-conductance-chnge-n-bimolecular-lipiD-membrane-presence-certin-antibiotics/>
- Dagar, S. S., Kumar, S., Mudgil, P., and Puniya, A. K. (2018). Comparative evaluation of lignocellulolytic activities of filamentous cultures of monocentric and polycentric anaerobic fungi. *Anaerobe*, 50, 76–79. <https://doi.org/10.1016/j.anaerobe.2018.02.004>
- de Gouvêa, P. F., Bernardi, A. V., Gerolamo, L. E., de Souza Santos, E., Riaño-Pachón, D. M., Uyemura, S. A., and Dinamarco, T. M. (2018). Transcriptome and secretome analysis of *Aspergillus fumigatus* in the presence of sugarcane bagasse. *BMC Genomics*, 19(1), 1–18. <https://doi.org/10.1186/s12864-018-4627-8>
- de Vries, R. P., Michelsen, B., Poulsen, C. ., Kroon, P. ., van den Heuvel, R. H. ., Faulds, C. B., Williamson, G., Hombergh, J. P. T. W. V. A. N. D. E. N., and Visser, J. (1997). The *faeA* genes from *Aspergillus niger* and *Aspergillus tubingensis* encode ferulic acid esterases involved in degradation of complex cell wall polysaccharides. *Applied and Environmental Microbiology*, 63(12),

4638–4644.

- Delmas, S., Pullan, S. T., Gaddipati, S., Kokolski, M., Malla, S., Blythe, M. J., Ibbett, R., Campbell, M., Liddell, S., Aboobaker, A., Tucker, G. A., and Archer, D. B. (2012). Uncovering the genome-wide transcriptional responses of the filamentous fungus *Aspergillus niger* to lignocellulose using RNA Sequencing. *PLoS Genetics*, 8(8), 1–13. <https://doi.org/10.1371/journal.pgen.1002875>
- Denning, E. J., and MacKerell, A. D. (2012). Intrinsic contribution of the 2'-hydroxyl to RNA conformational heterogeneity. *Journal of the American Chemical Society*, 134(5), 2800–2806. <https://doi.org/10.1021/ja211328g>. Intrinsic
- Deshpande, D., Chhugani, K., Chang, Y., Karlsberg, A., Loeffler, C., Zhang, J., Muszyńska, A., Munteanu, V., Yang, H., Rotman, J., Tao, L., Balliu, B., Tseng, E., Eskin, E., Zhao, F., Mohammadi, P., P. Łabaj, P., and Mangul, S. (2023). RNA-seq data science: From raw data to effective interpretation. *Frontiers in Genetics*, 14, 1–12. <https://doi.org/10.3389/fgene.2023.997383>
- Deutscher, J. (2008). The mechanism of CCR in bacteria. *Current Opinion in Microbiology*, 11, 87–93.
- Dey, P., and Roy, A. (2018). Molecular structure and catalytic mechanism of fungal family G acidophilic xylanases. *3 Biotech*, 8(2), 1–13. <https://doi.org/10.1007/s13205-018-1091-8>
- Dietl, A., Meir, Z., Shadkchan, Y., and Osherov, N. (2018). Riboflavin and pantothenic acid biosynthesis are crucial for iron homeostasis and virulence in the pathogenic mold *Aspergillus fumigatus*. *Virulence*, 9(1), 1036–1049. <https://doi.org/10.1080/21505594.2018.1482181>
- Dowzer, C. E. A., and Kelly, J. M. (1991). *Analysis of the creA gene, a regulator of carbon catabolite repression in Aspergillus nidulans*. 11(11). <https://doi.org/10.1128/MCB.11.11.5701>. Updated
- Ebbole, D. J. (1998). *Carbon catabolite repression of gene expression and conidiation in Neurospora crassa*. 21, 15–21.
- Eddy, S. R. (2001). Non-coding RNA genes and the modern RNA world. *Nature Reviews Genetics*, 2(12), 919–929. <https://doi.org/10.1038/35103511>
- Farani de Souza, D., Giatti Marques de Souza, C., and Peralta, R. M. (2001). Effect of easily metabolizable sugars in the production of xylanase by *Aspergillus tamaritii* in solid-state fermentation. *Process Biochemistry*, 36(8–9), 835–838. [https://doi.org/10.1016/S0032-9592\(00\)00295-8](https://doi.org/10.1016/S0032-9592(00)00295-8)
- Flores, M. E., Perea, M., Rodriguez, O., Malvaez, A., and Huitron, C. (1996). Physiological studies on induction and catabolite repression of β -xylosidase and endoxylanase in *Streptomyces* CH-M-1035. *Journal of Biotechnology*, 49, 179–187.
- Fu, L. H., Jiang, N., Li, C. X., Luo, X. M., Zhao, S., and Feng, J. X. (2019). Purification and characterization of an endo-xylanase from *Trichoderma sp.*, with xylobiose as the main product from xylan hydrolysis. *World Journal of Microbiology and Biotechnology*, 35(11). <https://doi.org/10.1007/s11274-019-2747-1>
- Furukawa, T., Shida, Y., Kitagami, N., Mori, K., Kato, M., Kobayashi, T., Okada,

- H., Ogasawara, W., and Morikawa, Y. (2009). Identification of specific binding sites for XYR1, a transcriptional activator of cellulolytic and xylanolytic genes in *Trichoderma reesei*. *Fungal Genetics and Biology*, 46(8), 564–574. <https://doi.org/10.1016/j.fgb.2009.04.001>
- Ghoshal, G., Banerjee, U. C., and Shivhare, U. S. (2016). Utilization of Agrowaste And Xylanase Production in Solid State Fermentation. *Journal of Biochemical Technology*, 6(3), 1013–1024.
- Grimmler, C., Held, C., Liebl, W., and Ehrenreich, A. (2010). Transcriptional analysis of catabolite repression in *Clostridium acetobutylicum* growing on mixtures of D-glucose and D-xylose. *Journal of Biotechnology*, 150, 315–323. <https://doi.org/10.1016/j.jbiotec.2010.09.938>
- Gupta, U., and Kar, R. (2009). Xylanase production by a thermo-tolerant *Bacillus* species under solid-state and submerged fermentation. *Brazilian Archives of Biology and Technology*, 52(6), 1363–1371. <https://doi.org/10.1590/S1516-89132009000600007>
- Hidayati, F. L. N., Suroto, D. A., Sardjono, Cahyanto, M. N., and Widada, J. (2022). Whole-genome sequence data of cellulase-producing fungi *Trichoderma asperellum* PK1J2, isolated from palm empty fruit bunch in Riau, Indonesia. *Data in Brief*, 45, 108607. <https://doi.org/10.1016/j.dib.2022.108607>
- Hidayati, Fela L. N. (2024). Pendekatan genotranskriptomik untuk memahami regulasi represi katabolit karbon pada produksi selulase oleh jamur lokal Indonesia [Universitas Gadjah Mada]. In *Fakultas Teknologi Pertanian*. <https://etd.repository.ugm.ac.id/>
- Hidayatullah, I. M., Setiadi, T., Tri, M., and Penia, A. (2020). *Bioresource technology xylanase inhibition by the derivatives of lignocellulosic material*. 300(December 2019).
- Huang, D. W., Sherman, B. T., Tan, Q., Kir, J., Liu, D., Bryant, D., Guo, Y., Stephens, R., Baseler, M. W., Lane, H. C., and Lempicki, R. A. (2007). DAVID Bioinformatics Resources: Expanded annotation database and novel algorithms to better extract biology from large gene lists. *Nucleic Acids Research*, 35, 169–175. <https://doi.org/10.1093/nar/gkm415>
- Ichinose, S., Tanaka, M., and Shintani, T. (2014). Improved α -amylase production by *Aspergillus oryzae* after a double deletion of genes involved in carbon catabolite repression. *Applied Microbiology and Biotechnology*, 89, 335–343. <https://doi.org/10.1007/s00253-013-5353-4>
- Ichinose, S., Tanaka, M., Shintani, T., and Gomi, K. (2018). Increased production of biomass-degrading enzymes by double deletion of *creA* and *creB* genes involved in carbon catabolite repression in *Aspergillus oryzae*. *Journal of Bioscience and Bioengineering*, 125(2), 141–147. <https://doi.org/10.1016/j.jbiosc.2017.08.019>
- Ilmén, M., Thrane, C., and Penttilä, M. (1996). The glucose repressor gene *cre1* of *Trichoderma*: Isolation and expression of a full-length and a truncated mutant form. *Molecular and General Genetics MGG*, 251(4), 451–460. <https://doi.org/10.1007/bf02172374>
- Iqbal, M., Rianse, K., Millati, R., and Indrati, R. (2020). Enhanced Fermentable Sugars Production from Sago Waste by *Trichoderma reesei* Pk1J2 and

- Aspergillus niger* FNCC 6114 Fermentation. *International Journal of Science and Research (IJSR)*, 9(6), 1200–1204. <https://doi.org/10.21275/SR20617111902>
- Irfan, M., Nadeem, M., and Syed, Q. (2014). One-factor-at-a-time (OFAT) optimization of xylanase production from *Trichoderma viride* -IR05 in solid-state fermentation. *Journal of Radiation Research and Applied Sciences*, 7(3), 317–326. <https://doi.org/10.1016/j.jrras.2014.04.004>
- Isikgor, F. H., and Becer, C. R. (2015). Lignocellulosic biomass: a sustainable platform for the production of bio-based chemicals and polymers. *Polym. Chem.*, 6(25), 4497–4559. <https://doi.org/10.1039/C5PY00263J>
- Ja'afaru, M. I. (2013). Screening of fungi isolated from environmental samples for xylanase and cellulase production. *ISRN Microbiology*, 2013, 1–7. <https://doi.org/10.1155/2013/283423>
- Jekosch, K., and Kuck, U. (2000). Loss of glucose repression in an *Acremonium chrysogenum* β -lactam producer strain and its restoration by multiple copies of the cre1 gene. *Applied Microbiology and Biotechnology*, 54(4), 556–563. <https://doi.org/10.1007/s002530000422>
- Jiang, X., Geng, A., He, N., Li, Q., and Ioeng, J. B. I. B. (2011). New isolate of *Trichoderma viride* strain for enhanced cellulolytic enzyme complex production. *JBIOSC*, 111(2), 121–127. <https://doi.org/10.1016/j.jbiosc.2010.09.004>
- Juturu, V., and Wu, J. C. (2012). Microbial xylanases: Engineering, production and industrial applications. *Biotechnology Advances*, 30(6), 1219–1227. <https://doi.org/10.1016/j.biotechadv.2011.11.006>
- Juturu, V., and Wu, J. C. hua. (2014). Microbial exo-xylanases: a mini review. *Applied Biochemistry and Biotechnology*, 174(1), 81–92. <https://doi.org/10.1007/s12010-014-1042-8>
- Kelly, J. M. (2004). *19 The Regulation of carbon metabolism in filamentous fungi I . Introduction 111 . genetic analysis of carbon catabolite repression in Aspergillus nidulans 11 . Systems regulated by carbon.*
- Khanahmadi, M., Arezi, I., Amiri, M., and Miranzadeh, M. (2018). Bioprocessing of agro-industrial residues for optimization of xylanase production by solid-state fermentation in flask and tray bioreactor. *Biocatalysis and Agricultural Biotechnology*, 13(January), 272–282. <https://doi.org/10.1016/j.bcab.2018.01.005>
- Kiss, T., and Kiss, L. (2000). Purification and characterization of an extracellular-xylosidase from *Aspergillus carbonarius*. *World Journal of Microbiology and Biotechnology*, 16, 465–470.
- Knob, A., and Carmona, E. C. (2010). β -Xylosidases from filamentous fungi : An overview. *World Journal of Microbiology and Biotechnology*, 26, 389–407. <https://doi.org/10.1007/s11274-009-0190-4>
- Kukurba, K. R., and Montgomery, S. B. (2015). RNA sequencing and analysis. *Cold Spring Harb Protoc*, 11, 951–969. <https://doi.org/10.1201/b11351-11>
- Kumar, D., Kumar, S. S., Kumar, J., Kumar, O., Mishra, S. V., Kumar, R., and Malyan, S. K. (2017). Xylanases and their industrial applications: A review. *Biochemical and Cellular Archives*, 17(1), 353–360.

- Kundu, A., Rani, R., Sweet, Á., and Optimization, Á. (2013). Production of intracellular β -xylosidase from the submerged fermentation of citrus wastes by *Penicillium janthinellum* MTCC. *Biotech*, 3, 241–246. <https://doi.org/10.1007/s13205-012-0091-3>
- Larcombe, D. E., Bohovych, I. M., Pradhan, A., Ma, Q., Hickey, E., Leaves, I., Cameron, G., Avelar, G. M., de Assis, L. J., Childers, D. S., Bain, J. M., Lagree, K., Mitchell, A. P., Netea, M. G., Erwig, L. P., Gow, N. A. R., and Brown, A. J. P. (2023). Glucose-enhanced oxidative stress resistance —A protective anticipatory response that enhances the fitness of *Candida albicans* during systemic infection. *PLoS Pathogens*, 19(7), 1–26. <https://doi.org/10.1371/journal.ppat.1011505>
- Lee, S. H., Lim, V., and Lee, C. K. (2018). Newly isolate highly potential xylanase producer strain from various environmental sources. *Biocatalysis and Agricultural Biotechnology*, 16(August), 669–676. <https://doi.org/10.1016/j.bcab.2018.09.024>
- Li, C., and Yan, T. (2014). Use of *Aspergillus niger* β -glucosidase II gene (*bgl II*) promoter elements to construct an efficient expression vector. *Journal of the Taiwan Institute of Chemical Engineers*, 45(3), 749–754. <https://doi.org/10.1016/j.jtice.2013.09.022>
- Li, J., Lin, L., Li, H., Tian, C., and Ma, Y. (2014). Transcriptional comparison of the filamentous fungus *Neurospora crassa* growing on three major monosaccharides D-glucose, D-xylose and L-arabionose. *Biotechnology for Biofuels*, 7(31), 1–15. <https://doi.org/10.1186/1754-6834-7-31>
- Liu, S., Hu, W., Wang, Z., and Chen, T. (2020). Production of riboflavin and related cofactors by biotechnological processes. *Microbial Cell Factories*, 1–16. <https://doi.org/10.1186/s12934-020-01302-7>
- López, Y., García, A., Karimi, K., Taherzadeh, M. J., and Martín, C. (2010). Chemical characterisation and dilute-acid hydrolysis of rice hulls from an Artisan Mill. *BioResources*, 5(4), 2268–2277. <https://doi.org/10.15376/biores.5.4.2268-2277>
- Luong, T. T. H., Poëaim, S., and Tangthirasunun, N. (2023). Isolation and characterization of xylanase from a novel strain, *Penicillium menonorum* SP10. *Mycobiology*, 51(4), 239–245. <https://doi.org/10.1080/12298093.2023.2247221>
- Mach-Aigner, A. R., Pucher, M. E., and Mach, R. L. (2010). D-xylose as a repressor or inducer of xylanase expression in *Hypocrea jecorina* (*Trichoderma reesei*). *Applied and Environmental Microbiology*, 76(6), 1770–1776. <https://doi.org/10.1128/AEM.02746-09>
- Mach, R. L., Strauss, J., Zeilinger, S., Schindler, M., and Kubicek, C. P. (1996). Carbon catabolite repression of xylanase I (*xynI*) gene expression in *Trichoderma reesei*. *Molecular Microbiology*, 21(6), 1273–1281. <https://doi.org/10.1046/j.1365-2958.1996.00094.x>
- Maciel, G. M., Vandenbergh, L. P. D. S., Haminiuk, C. W. I., Fendrich, R. C., Bianca, B. E. D., Da Silva Brandalize, T. Q., Pandey, A., and Soccol, C. R. (2008). Xylanase production by *Aspergillus niger* LPB 326 in solid-state fermentation using statistical experimental designs. *Food Technology and*

- Biotechnology*, 46(2), 183–189.
- Mandels, M., Parrish, F. W., and Reese, E. T. (1962). Sophorose as an inducer of cellulase in *Trichoderma viride*. *Journal of Bacteriology*, 83, 400–408. <https://doi.org/10.1128/jb.83.2.400-408.1962>
- Mangan, D., Cornaggia, C., Liadova, A., McCormack, N., Ivory, R., McKie, V. A., Ormerod, A., and McCleary, B. V. (2017). Novel substrates for the automated and manual assay of endo-1,4- β -xylanase. *Carbohydrate Research*, 445, 14–22. <https://doi.org/10.1016/j.carres.2017.02.009>
- Marques, N. P., de Cassia Pereira, J., Gomes, E., da Silva, R., Araújo, A. R., Ferreira, H., Rodrigues, A., Dussán, K. J., and Bocchini, D. A. (2018). Cellulases and xylanases production by endophytic fungi by solid state fermentation using lignocellulosic substrates and enzymatic saccharification of pretreated sugarcane bagasse. *Industrial Crops and Products*, 122, 66–75. <https://doi.org/10.1016/j.indcrop.2018.05.022>
- McCleary, B. V., and McGeough, P. (2015). A comparison of polysaccharide substrates and reducing sugar methods for the measurement of endo-1,4- β -xylanase. *Applied Biochemistry and Biotechnology*, 177(5), 1152–1163. <https://doi.org/10.1007/s12010-015-1803-z>
- Megazyme, P. A. (2017). *Assay procedure XylX6 method*.
- Metreveli, E., Khardziani, T., and Elisashvili, V. (2021). The carbon source controls the secretion and yield of polysaccharide-hydrolyzing enzymes of *Basidiomycetes*. *Biomolecules*, 11(1341), 1–10. <https://doi.org/10.3390/biom11091341>
- Midorikawa, G. E. O., Correa, C. L., Noronha, E. F., Filho, E. X. F., Togawa, R. C., Costa, M. M. do C., Silva-junior, O. B., Grynberg, P., and Neil, G. M. R. (2018). Analysis of the transcriptome in *Aspergillus tamaris* during enzymatic degradation of sugarcane bagasse. *Frontiers in Bioengineering and Biotechnology*, 6(123), 1–17. <https://doi.org/10.3389/fbioe.2018.00123>
- Mikus, L., Dodok, L., Kovacova, M., Staruch, L., and Koman, V. (2012). Bakery enzymes in cereal technologies. *Potravinarstvo*, 6(3), 10–15. <https://doi.org/10.5219/193>
- Mirhendi, H., Zarei, F., Motamedi, M., and Nouripour-Sisakht, S. (2016). *Aspergillus tubingensis* and *Aspergillus niger* as the dominant black *Aspergillus*, use of simple PCR-RFLP for preliminary differentiation. *Journal de Mycologie Medicale*, 26(1), 9–16. <https://doi.org/10.1016/j.mycmed.2015.12.004>
- Monteiro, L. M. oliveira, Vici, A. claudia, Pinheiro, M. P., Heinen, P. R., Oliveira, A. H. C. de, Ward, R. J., Prade, R. A., Buckeridge, M. S., & Polizeli, M. de L. T. de M. (2020). A Highly glucose tolerant β -Glucosidase from *Malbranchea pulchella* (Mp Bg3) enables cellulose saccharification. *Scientific Reports*, 10(6998), 1–12. <https://doi.org/10.1038/s41598-020-63972-y>
- Mulder, H. J., Saloheimo, M., Penttilä, M., and Madrid, S. M. (2004). The transcription factor HACA mediates the unfolded protein response in *Aspergillus niger*, and up-regulates its own transcription. *Molecular Genetics and Genomics*, 271(2), 130–140. <https://doi.org/10.1007/s00438-003-0965-5>
- Naidu, D. S., Hlangothi, S. P., and John, M. J. (2018). Bio-based products from

- xylan: A review. *Carbohydrate Polymers*, 179, 28–41. <https://doi.org/10.1016/j.carbpol.2017.09.064>
- Nurnawati, E., Margino, S., Martani, E., and Sarto. (2014). Isolation, screening and identification xylanolytic local fungi that potentially as pulp bleaching agents. *Jurnal Manusia Dan Lingkungan*, 21(3), 317–322.
- Pandey, A. K., Edgard, G., and Negi, S. (2016). Optimization of concomitant production of cellulase and xylanase from *Rhizopus oryzae* SN5 through EVOP-factorial design technique and application in Sorghum Stover based bioethanol production. *Renewable Energy*, 98, 51–56. <https://doi.org/10.1016/j.renene.2016.05.071>
- Piaga, F., Fernández-Espinar, M. T., Vallés, S., and Ramón, D. (2006). Xylanase production in *Aspergillus nidulans*: induction and carbon catabolite repression. *FEMS Microbiology Letters*, 115(2–3), 319–323. <https://doi.org/10.1111/j.1574-6968.1994.tb06657.x>
- Pinaga, F., Fernandez-Espinar, M. T., Valles, S., and Ramon, D. (1994). Xylanase production in *Aspergillus nidulans*: induction and carbon catabolite repression. *FEMS Microbiology Letters*, 115, 319–323. <https://doi.org/10.1111/j.1574-6968.1994.tb06657.x>
- Ping, L., Wang, M., Yuan, X., Cui, F., Huang, D., Sun, W., Zou, B., Huo, S., and Wang, H. (2018). Production and characterization of a novel acidophilic and thermostable xylanase from *Thermoascus aurantiacus*. *International Journal of Biological Macromolecules*, 109, 1270–1279. <https://doi.org/10.1016/j.ijbiomac.2017.11.130>
- Pócsi, I., Prade, R. A., and Penninckx, M. J. (2004). Glutathione, altruistic metabolite in fungi. In *Advances in Microbial Physiology* (Vol. 49, Issue 04). [https://doi.org/10.1016/S0065-2911\(04\)49001-8](https://doi.org/10.1016/S0065-2911(04)49001-8)
- Polizeli, M. L. T. M., Rizzatti, A. C. S., Monti, R., Terenzi, H. F., Jorge, J. A., and Amorim, D. S. (2005). Xylanases from fungi: Properties and industrial applications. *Applied Microbiology and Biotechnology*, 67(5), 577–591. <https://doi.org/10.1007/s00253-005-1904-7>
- Portnoy, T., Margeot, A., Linke, R., Atanasova, L., Fekete, E., Sándor, E., Hartl, L., Karaffa, L., Druzhinina, I. S., Seiboth, B., Le Crom, S., and Kubicek, C. P. (2011). The CRE1 carbon catabolite repressor of the fungus *Trichoderma reesei*: A master regulator of carbon assimilation. *BMC Genomics*, 12, 1–12. <https://doi.org/10.1186/1471-2164-12-269>
- Rauscher, R., Würleitner, E., Wacenovský, C., Aro, N., Stricker, A. R., Zeilinger, S., Kubicek, C. P., Penttilä, M., and Mach, R. L. (2006). Transcriptional regulation of *xynI*, encoding xylanase I, in *Hypocrea jecorina*. *Eukaryotic Cell*, 5(3), 447–456. <https://doi.org/10.1128/EC.5.3.447-456.2006>
- Ravichandra, K., Yaswanth, V. V. N., Nikhila, B., Ahmad, J., Srinivasa Rao, P., Uma, A., Ravindrababu, V., and Prakasham, R. S. (2016). Xylanase production by isolated fungal strain, *Aspergillus fumigatus* RSP-8 (MTCC 12039): Impact of Agro-industrial Material as Substrate. *Sugar Tech*, 18(1), 29–38. <https://doi.org/10.1007/s12355-014-0357-7>
- Ries, L., Pullan, S. T., Delmas, S., Malla, S., Blythe, M. J., and Archer, D. B. (2013). Genome-wide transcriptional response of *Trichoderma reesei* to

- lignocellulose using RNA sequencing and comparison with *Aspergillus niger*. *BMC Genomics*, *14*(541), 1–12.
- Roy, P., Lockington, R. A., and Kelly, J. M. (2008). CreA-mediated repression in *Aspergillus nidulans* does not require transcriptional auto-regulation, regulated intracellular localisation or degradation of CreA. *Fungal Genetics and Biology*, *45*(5), 657–670. <https://doi.org/10.1016/j.fgb.2007.10.016>
- Said, S.D., Hasan, M., and Romachandran, K.B. (2013). Effect of pH on growth, spore production and spore viability of biocontrol agent *Trichoderma harzianum* in submerged fermentation. *ASEAN Journal of Chemical Engineering*, *3*(1), 14-18.
- Safitri, I. O. (2022). Draft genome sequence of *Trichoderma asperellum* MLT1J1 , isolated from coconut husk in Maluku , Indonesia. *Korean Journal of Microbiology*, *58*(3), 1–3.
- Sakthiselvan, P., Naveena, B., and Partha, N. (2014). Molecular characterization of a Xylanase-producing fungus isolated from fouled soil. *45*(4), 1293–1302.
- Sanghvi, G. V., Koyani, R. D., and Rajput, K. S. (2010). Thermostable xylanase production and partial purification by solid-state fermentation using agricultural waste wheat straw. *1203*. <https://doi.org/10.1080/21501203.2010.484029>
- Scotti, C. T., Vergoignan, C., Feron, G., and Durand, A. (2001). Glucosamine measurement as indirect method for biomass estimation of *Cunninghamella elegans* grown in solid state cultivation conditions. *Biochemical Engineering Journal*, *7*(1), 1–5. [https://doi.org/10.1016/S1369-703X\(00\)00090-5](https://doi.org/10.1016/S1369-703X(00)00090-5)
- Shariq, M., and Sohail, M. (2018). Application of *Candida tropicalis* MK-160 for the production of xylanase and ethanol. *Journal of King Saud University - Science*. <https://doi.org/10.1016/j.jksus.2018.04.009>
- Shida, Y., Furukawa, T., and Ogasawara, W. (2016). Deciphering the molecular mechanisms behind cellulase production in *Trichoderma reesei*, the hypercellulolytic filamentous fungus. *Bioscience, Biotechnology and Biochemistry*, *80*(9), 1712–1729. <https://doi.org/10.1080/09168451.2016.1171701>
- Shimokawa, T., Yamaguchi, M., Shibuya, H., Ishihara, M., Ikeda, T., Magara, K., and Nojiri, M. (2012). Screening of lignocellulolytic enzyme producers : enzyme system from *Aspergillus tubingensis* for hydrolysis of sugi pulp. *Bulletin of FFPRI*, *11*(2), 57–63.
- Singh, N., and Gaur, S. (2021). GRAS Fungi: A new horizon in safer food product. In *Fungi in Sustainable Food Production* (pp. 27–37). https://doi.org/10.1007/978-3-030-64406-2_3.
- Singh, N., Sithole, B., & Govinden, R. (2023). Screening for cellulases and preliminary optimisation of glucose tolerant β -glucosidase production and characterisation. *Mycology*, *14*(2), 91–107. <https://doi.org/10.1080/21501203.2022.2155261>
- Solís-Pereira, S., Favela-Torres, E., Viniegra-González, G., and Gutiérrez-Rojas, M. (1993). Effects of different carbon sources on the synthesis of pectinase by *Aspergillus niger* in submerged and solid state fermentations. *Applied Microbiology and Biotechnology*, *39*(1), 36–41. <https://doi.org/10.1007/BF00166845>

- Strauss, J., Horvath, H. K., Basem, M., Kindermann, J., Mach, R. L., and Kubicek, C. P. (1999). *The function of CreA , the carbon catabolite repressor of Aspergillus nidulans , is regulated at the transcriptional and post-transcriptional level.* 32, 169–178.
- Strauss, J., Mach, R. L., Zeilinger, S., Hartler, G., Stöffler, G., Wolschek, M., and Kubicek, C. P. (1995). CreI, the carbon catabolite repressor protein from *Trichoderma reesei*. *FEBS Letters*, 376(1–2), 103–107. [https://doi.org/10.1016/0014-5793\(95\)01255-5](https://doi.org/10.1016/0014-5793(95)01255-5)
- Stricker, A. R., Grosstessner-Hain, K., Würleitner, E., and Mach, R. L. (2006). Xyr1 (Xylanase Regulator 1) regulates both the hydrolytic enzyme system and D-xylose metabolism in *Hypocrea jecorina*. *Eukaryotic Cell*, 5(12), 2128–2137. <https://doi.org/10.1128/EC.00211-06>
- Stricker, A. R., Trefflinger, P., Aro, N., Penttilä, M., and Mach, R. L. (2008). Role of Ace2 (Activator of Cellulases 2) within the *xyn2* transcriptosome of *Hypocrea jecorina*. *Fungal Genetics and Biology*, 45(4), 436–445. <https://doi.org/10.1016/j.fgb.2007.08.005>
- Susca, A., Stea, G., and Perrone, G. (2007). PCR identification of *Aspergillus niger* and *Aspergillus tubingensis* based on calmodulin gene. *Food Additive and Contaminants*, 24(10), 1154–1160.
- Tanaka, T., Terauchi, Y., and Yoshimi, A. (2022). *Aspergillus hydrophobins : Physicochemical properties , biochemical properties , and functions in solid polymer degradation.*
- Todd, R. B., Davis, M. A., and Hynes, M. J. (2007). Genetic manipulation of *Aspergillus nidulans*: Heterokaryons and diploids for dominance, complementation and haploidization analyses. *Nature Protocols*, 2(4), 822–830. <https://doi.org/10.1038/nprot.2007.113>
- Vautard, G., Cotton, P., and Fèvre, M. (1999). The glucose repressor CRE1 from *Sclerotinia sclerotiorum* is functionally related to CREA from *Aspergillus nidulans* but not to the Mig proteins from *Saccharomyces cerevisiae*. *FEBS Letters*, 453(1–2), 54–58. [https://doi.org/10.1016/S0014-5793\(99\)00691-2](https://doi.org/10.1016/S0014-5793(99)00691-2)
- Wang, Y., Yan, H., Neng, J., Gao, J., Yang, B., and Liu, Y. (2020). The Influence of NaCl and Glucose Content on Growth and Ochratoxin A Production by *Aspergillus ochraceus*, *Aspergillus carbonarius* and *Penicillium nordicum*. *Toxins*, 12(515), 1–15. <https://doi.org/10.3390/toxins12080515>
- Wang, D., and Farhana, A. (2023). Biochemistry , RNA Structure. In *Biochemistry, RNA Structure.* StatPearls Publishing. <https://www.ncbi.nlm.nih.gov/books/NBK558999/>
- Wang, Y., Yan, H., Neng, J., Gao, J., Yang, B., and Liu, Y. (2020). The Influence of NaCl and glucose content on growth and ochratoxin a production by *Aspergillus ochraceus*, *Aspergillus carbonarius* and *Penicillium nordicum*. *Toxins*, 12(515), 1–15. <https://doi.org/10.3390/toxins12080515>
- Wang, Z., Gerstein, M., and Snyder, M. (2009). RNA-Seq: a revolutionary tool for transcriptomics *Zhong. Nature Review Genetics*, 10(1), 57–63. <https://doi.org/10.1038/nrg2484>.RNA-Seq
- Wei, X., Zheng, K., Chen, M., Liu, G., Li, J., Lei, Y., Qin, Y., and Qu, Y. (2011). Transcription analysis of lignocellulolytic enzymes of *Penicillium decumbens*

- 114-2 and its catabolite-repression-resistant mutant. *Comptes Rendus Biologies*, 334, 806–811. <https://doi.org/10.1016/j.crv.2011.06.002>
- Wong, K. E. N. K. Y., Tan, L. U. L., and Saddler, J. N. (1988). *Multiplicity of 3-1, 4-Xylanase in Microorganisms : Functions and Applications*. 52(3), 305–317.
- Young, M. D., Wakefield, M. J., Smyth, G. K., and Oshlack, A. (2010). Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biology*, 11(2). <https://doi.org/10.1186/gb-2010-11-2-r14>
- Zhang, X., Ma, B., Liu, J., Chen, X., Li, S., Su, E., Gao, L., & Li, H. (2020). β -glucosidase genes differentially expressed during composting. *Biotechnology for Biofuels*, 13(174), 1–11. <https://doi.org/10.1186/s13068-020-01813-w>
- Zeilinger, S., Mach, R. L., Schindler, M., Herzog, P., and Kubicek, C. P. (1996). Different inducibility of expression of the two xylanase genes *xyn1* and *xyn2* in *Trichoderma reesei*. *Journal of Biological Chemistry*, 271(41), 25624–25629. <https://doi.org/10.1074/jbc.271.41.25624>