

DAFTAR PUSTAKA

- Arnama, I.N. 2018. Pertumbuhan dan produksi dua varietas padi sawah (*Oryza sativa* L.) dengan variasi jumlah bibit per rumpun. *Skripsi*. Universitas Hasanuddin, Makassar.
- Azizah, F., Suwarsito, S., & Sarjanti, E. (2021). Pengaruh Pola Curah Hujan Terhadap Produktivitas Padi di Kecamatan Bukateja Kabupaten Purbalingga. *Sainteks*, 18(1), 1. <https://doi.org/10.30595/sainteks.v18i1.10567>
- Azeez, M. A., Adubi, A. O., & Durodola, F. A. (2018). Landraces and Crop Genetic Improvement. *Rediscovery of Landraces as a Resource for the Future*. doi:10.5772/intechopen.75944.
- Bouman, B.A.M., Lampayan, R. M., & Tuong, T. P. 2007. Water Management in Irrigated Rice. Internasional Rice Research Institute. Los Banos Phillipiness.
- Berntson, G.M., Lynch, J.P., Snapp, S., 1997. Fractal geometry and plant root systems: current perspectives and future applications. In: Baveye, P., Parlange, J.Y., Stewart, B.A. (Eds.), *Fractals in Soil Science*. Lewis Publishers, New York, pp. 113–152.
- BPS 2021. Produksi Padi tahun 2021. diunduh dari <https://www.bps.go.id/pressrelease/2022/03/01/1909/produksi-padi-tahun-2021-turun-0-43-persen--angka-tetap>. diakses 8 Juni 2023.
- Champoux M., Wang G., Sarkarung S., Mackill D., O'Toole J., Huang N., McCouch S. Locating genes associated with root morphology and drought avoidance in rice via linkage to molecular markers. *Theor. Appl. Genet.* 1995;90:969–981. doi: 10.1007/BF00222910.
- Chaves, M. M., Maroco, J. P., & Pereira, J. S. (2003). Understanding plant responses to drought - From genes to the whole plant. *Functional Plant Biology*, 30(3), 239–264. <https://doi.org/10.1071/FP02076>
- Cheng, C., Motohashi, R., Tsuchimoto, S., Fukuta, Y., Ohtsubo, H., & Ohtsubo, E. (2003). Polyphyletic origin of cultivated rice: Based on the interspersion pattern of SINEs. *Molecular Biology and Evolution*, 20(1), 67–75. <https://doi.org/10.1093/molbev/msg004>
- Clark, R. T., MacCurdy, R. B., Jung, J. K., Shaff, J. E., McCouch, S. R., Aneshansley, D. J., & Kochian, L. V. (2011). Three-dimensional root phenotyping with a novel imaging and software platform. *Plant Physiology*, 156(2), 455–465. <https://doi.org/10.1104/pp.110.169102>.
- Dannowski, M., Block, A., 2005. Fractal geometry and root system structures of heterogeneous plant communities. *Plant Soil* 272 (1–2), 61.

- De Datta S.K. 1981. Principles and Practices of Rice Production. *A Wiley Interscience Publication*. New York : John Wiley & Sons.
- Eghball, B., Mielke, L. N., Calvo, G. A., & Wilhelm, W. W. (1993). Fractal Description of Soil Fragmentation for Various Tillage Methods and Crop Sequences. *Soil Science Society of America Journal*, 57(5), 1337–1341. <https://doi.org/10.2136/sssaj1993.03615995005700050029x>.
- Farooq M, A. Wahid, N. Kobayashi, D. Fujita, dan Basra, 2009. Plant Drought Stress: Effects, Mechanisms and Management. *Agron Sustain. Dev.*, 29(1): 185–212.
- Fitter, A.H., 1994. Architecture and biomass allocation as components of the plastic response of root systems to soil heterogeneity. In: Caldwell, M.M., Percy, R.W. (Eds.), *Exploitation of Environmental Heterogeneity by Plants*. Academic Press, San Diego, CA, pp. 305–323.
- Fitter, A.H., Stickland, T.R., 1992. Fractal characterization of root-system architecture. *Funct. Ecol.* 6, 632–635.
- Fuadi, N.A, Purwanto, M.Y, Tarigan, S.G. 2016. Kajian kebutuhan air dan produktivitas air padi sawah dengan sistem pemberian air secara SRI dan konvensional menggunakan irigasi pipa. *Jurnal Irigasi*. Vol 11 No 1.
- Ghozali, G. M dan Supriyanta. 2022. Deteksi Molekuler dan Validasi Kemampuan Padi Lokal (*Oryza sativa* L.) Membentuk Perakaran Dalam pada Kondisi Cekaman Kekeringan. *Skripsi*. Fakultas Pertanian Universitas Gadjah Mada.
- Gu, D., Zhen, F., Hannaway, D. B., Zhu, Y., Liu, L., Cao, W., & Tang, L. 2017. Quantitative classification of rice (*Oryza sativa* L.) root length and diameter using image analysis. *PLoS ONE*, 12(1), 1–14.
- Grist. D.H. 1959. Rice. Longmans, London. 472 p.
- Han, J., N. Shin, J. Moon, J. H. Chin, S. Yoo. 2016. A simple phenotyping method for deep-rooting rice grown in pots. *Plant Biotechnology* 43: 444-449.
- Handayani F., S. Sumarmiyanti, dan N.R Ahmadi. 2017. Morphological Variation of 20 Local Rice Cultivars of East Kalimantan. *Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia*.
- Harmansis, A. 2019. Warta, Penelitian dan Pengembangan Pertanian. Badan Penelitian dan Pengembangan Pertanian. Kementerian Pertanian. Vol 41 No 1.
- Hasanah N, E.S Bayu, E.H Kardhinata. 2020. Pengaruh Cekaman Kekeringan Terhadap Morfologi Akar Beberapa Genotipe Padi Beras Merah (*Oryza sativa*) pada Fase Vegetatif. *Jurnal Online Agroteknologi* 8 (1) : 73-80 DOI: 10.32734/jaet.
- Hikmah, N. & Dharmono. (2018). Keanekaragaman spesies famili poaceae di

Hutan Pantai Tabanio, Kecamatan Takisung, Kabupaten Tanah Laut.
Prosiding Seminar Nasional Lingkungan Lahan Basah, 3 (1), 249-253.

- Hasanah, I. 2007. Bercocok Tanam Padi. Azka Mulia Media. Jakarta. 68 hal.
- Izumi, Y., Kono, Y., Yamauchi, A., Iijima, M., 1995. Analysis of timecourse changes in root-system morphology of rice in excised root culture. *Jpn. J. Crop Sci.* 64,636–643.
- Izumi, Y., Kono, Y., Yamauchi, A., Iijima, M., 1997. Quantitative analysis of the architecture of seminal root system of rice (*Oryza sativa* L.) grown under different soil moisture conditions. *Jpn. J. Crop Sci.* 66, 418–426.
- Jain, A., Roorkiwal, M., Kale, S., Garg, V., Yadala, R., & Varshney, R. K. 2019. InDel markers: An extended marker resource for molecular breeding in chickpea. *PLoS ONE*, 14(3), 1–13. <https://doi.org/10.1371/journal.pone.0213999>
- Jehan, T., Lakhanpaul, S. 2006. Single Nucleotide Polymorphism (SNP)-Methods and Applications in Plant Genetics: A review. *Indian Journal of Biotechnology.* 5(4): 435-459.
- Jeki, Jeki. 2016. Indeks Sensitifitas Stres Beberapa Varietas Padi Gogo Pada Cekaman Kekeringan. *Agrotekbis: Jurnal Ilmu Pertanian* (e-journal) 4.4: 369-373.
- Kawata S, Yamazaki K, Ishihara K, Shibatama H, Lai KL. Studies on root system formation in rice plants in a paddy. *Japanese Journal of Crop Science.* 1963; 32:163-180. DOI: 10.1626/jcs.32.163
- Kano-Nakata, M., Inukai, Y., Wade, L. J., Siopongco, J. D., & Yamauchi, A. (2011). Root development, water uptake, and shoot dry matter production under water deficit conditions in two cssls of rice: Functional roles of root plasticity. *Plant Production Science*, 14(4), 307–317. <https://doi.org/10.1626/pps.14.307>.
- Kim, Y., Y. S. Chung, E. Lee, P. Tripathi, S. Heo, K. Kim. 2020. Root response to drought stress in rice (*Oryza sativa* L.). *International Journal of Molecular Sciences* 21: 1-22.
- Kumar, A., Ramegowda, S. Basu, A. Pereira. 2016. Mechanism of drought tolerance in rice. *Burleigh Dodds Science.*1-33.
- Kohyama K, Sodhi NS, Suzuki K, Sasaki T. 2016. Texture evaluation of cooked rice prepared from Japanese cultivars using two-bite instrumental test and electromyography. *Journal of Texture Studies.* 47: 188–198.
- Lynch, J.P., van Beem, J., 1993. Growth and architecture of seedling roots of common bean genotypes. *Crop Sci.* 33, 1253–1257.

- Makarim AK dan Suhartatik. 2009. Morfologi dan Fisiologi Tanaman Padi. Balai Besar Penelitian Tanaman Padi. Subang, Jawa Barat.
- Mandelbrot, B.B., 1983. The Fractal Geometry of Nature. WH Freeman, New York.
- Manivannan P, Jaleel CA, Kishorekumar A, Sankar B, Somasundaram R, Sridharan R, Panneerselvam R. 2007. Drought stress induced changes in the biochemical parameters and photosynthetic pigments of cotton (*Gossypium hirsutum* L.). *Indian J. Appl. Pure Biol.* 52:369-372.
- Maulana Z. 2017. Keragaman Plasma Nutfah Padi Lokal Sulawesi Selatan. CV. Sah Media. Makassar.
- Masdar. 2005. Interaksi jarak tanam dan jumlah bibit per titik tanam pada sistem intensifikasi padi terhadap pertumbuhan vegetatif tanaman. *Akta Agrosia Ed. Khusus*, (1):92-98.
- Masi, C.E.A., Maranville, J.W., 1998. Evaluation of sorghum root branching using fractals. *J. Agric. Sci.* 131, 259–265.
- Mawardi, C.N. Ichsan dan Syamsuddin. 2016. Pertumbuhan dan Hasil Beberapa Varietas Tanaman Padi (*Oryza sativa* L.) pada Tingkat Kondisi Kekeringan. *J. Ilmiah.* 1(1):176 –187.
- Matsuo T, Futsuhara Y, Kikuchi F, Yamaguchi H. 1997. Science of the Rice Plant, Food and Agriculture. Policy Research Center, Tokyo.
- Nadeem, M. A., M. A. Nawaz, M. Q. Shahid, Y. Dogan, G. Comertpay, M. Yildiz, R. Hatipoglu, F. Ahmad, A. Alsaleh, N. Labhane, H. Ozkan, G. Chung, F. S. Baloch. 2018. DNA molecular markers in plant breeding: status and recent advancements in genomic selection and genome editing. *Biotechnology & Biotechnological Equipment.* 32(2): 261-285.
- Nadzir Z.A, N. Simarmata, Aliffia. 2020. Pengembangan Algoritma Identifikasi Sawah Padi Berdasarkan Spektra Fase Padi (Studi Kasus : Lampung Selatan). *Jurnal Sains Informasi Geografi (SIG)* 3(1):23-36
- Nagai, I. 1962. Japonica Rice its breeding and culture. Yohendo Ltd. Tokyo.
- Nuraida, D. 2012. Pemuliaan tanaman cepat dan tepat melalui pendekatan marka molekuler. *El-Hayah.* 2(2): 97-103.
- Obara, M., W. Tamura, T. Ebitani, M. Yano, T. Sato, T. Yamaya. 2010. Fine-mapping of *qRL6.1*, a major QTL for root length of rice seedlings grown under a wide range of NH_4^+ concentrations in hydroponic conditions. *Theoretical and Applied Genetics.* 121(3):535-547.
- Paski, J.A.I., G.I.S.L Faski, M.F Handoyo, dan D.A.S Pertiwi. 2018. Analisis Neraca Air Lahan untuk Tanaman Padi dan Jagung di Kota Bengkulu. *Jurnal Ilmu Lingkungan* 15(2):83-89

- Pathak, H., Tewari, A., Sankhyan, S., Dubey, D., Mina, U., Singh, V. K., Jain, N., & Bhatia, A. (2011). Direct - seeded rice: Potential, performance and problems – A review. *Current Advances in Agricultural Sciences*. 3(2), 77–88.
- Richard, C.A., Hickey, L.T., Fletcher, S., Jennings, R., Chenu, R. & Christopher, J.T. 2015. High-throughput phenotyping of seminal root traits in wheat. *J. Plant Methods*. 11.13.
- Rohlf, F. J. (1998). RAPD protocol using NTSYSpc Version 2.0: User Guide. Applied Biostatistics Inc. Department of Ecology and Evolution. State University of New York. New York, United State of America, 43pp.
- Rollins, J. A., Habte, E., Templer, S. E., Colby, T., Schmidt, J., and Von Korff, M. (2013). Leaf proteome alterations in the context of physiological and morphological responses to drought and heat stress in barley (*Hordeum vulgare* L.). *J. Exp. Bot.* 64, 3201–3212. doi: 10.1093/jxb/ert158
- Rosadi, F. N. 2013. Studi Morfologi dan Fisiologi Padi (*Oryza sativa* L.) Toleran Kekeringan. Bogor: Tesis. Agronomi dan Hortikultura Institut Pertanian Bogor.
- Sasaki O, Yamazaki K, Kawata S. The relationship between the diameters and the structures of lateral roots in rice plants. *Japanese Journal of Crop Science*. 1984; 53:169-175. DOI: 10.1626/jcs.50.476
- Shashidhar, H. E, Henry. H, Hardy, B. 2012. Methodologies for Root Drought Studies in Rice. Los Baños, Filipina. International Rice Research Institute. 65p.
- Singh S, Prasad S, Yadav V, Kumar A and B Jaiswal, Kumar A, Khan N A and Dwivedi D K 2018 Effect of Drought Stress on Yield and Yield Components of Rice (*Oryza sativa* L.) Genotypes *International Journal of Current Microbiology and Applied Sciences*. (7) p2752 – 2759
- Sitorus, H.L. 2014. Respon Beberapa Kultivar Padi Gogo pada Ultisol terhadap Pemberian Alumunium dengan Konsentrasi Berbeda. *Skripsi*. Fakultas Pertanian Universitas Bengkulu. Bengkulu.
- Sujinah, dan A. Jamil. 2016. Mekanisme Respon Tanaman Padi terhadap Cekaman Kekeringan dan Varietas Toleran. *Iptek Tanaman Pangan* 11(1): 1-8.
- Sridevi dan Cheallamuthu, 2015. Beras SNI 6128:2015. Jakarta. Badan Standar Nasional.
- Swapna, M., M.S. Srivastava. 2012. Molecular Marker Applications for Improving Sugar Content in Sugarcane. *Springer*, London.
- Tatsumi, J., 1995. Fractal geometry in root systems: quantitative evaluation of distribution pattern. *Jpn. J. Crop Sci*. 64, 50–57.

- Tatsumi, J., 2001. Fractal geometry of root system morphology: fractal dimension and multifractals. In: Proc. 6th Symp. Int. Soc. Root Res., Nagoya, Japan, November 11–15, pp. 24–25.
- Tatsumi, J., Yamauchi, A., Kono, Y., 1989. Fractal analysis of plant root systems. *Ann.Bot.* 64, 499–503.
- Uga, Y., K. Okuno, M. Yano. 2011. *Dro1*, a major QTL involved in deep rooting of rice under upland field conditions. *Journal of Experimental Botany* 62(8): 2485-2494.
- Uga, Y., E. Hanzawa, S. Nagai, K. Sasaki, M. Yano. 2012. Identification of *qSOR1*, a major rice QTL involved in soil-surface rooting in paddy fields. *Theoretical and Applied Genetics*. 124: 75-86.
- Uga, Y., K. Sugimoto, S. Ogawa, J. Rane, M. Ishitani, N. Hara, Y. Kitomi, Y. Inukai, K. Ono, N. Kanno, H. Inoue, H. Takehisa, R. Motoyama, Y. Nagamura, J. Wu, T. Matsumoto, T. Takai, K. Okuno, M. Yano. 2013a. Control of root architecture by DEEPER ROOTING 1 increases rice yield under drought conditions. *Nature Genetics* 45(9):1097-1102.
- Walk, T.C., Van Erp, E., Lynch, J.P., 2004. Modelling applicability of fractal analysis to efficiency of soil exploration by roots. *Ann. Bot.* 94, 119–128.
- Wang, X., N. Samo, L. Li, M. Wang, M. Qadir, K. Jiang, J. Qin, F. Rasul, G. Yang, Y. Hu. 2019. Root distribution and its impacts on the drought tolerance capacity of hybrid rice in the Sinchuan Basin Area of China. *MDPI Agronomy* 9(13): 1-13.
- [USDA] United State Departement of Agriculture. 2024. Natural resources conservation service. <https://plants.usda.gov/home/classification/24211> (18 Juli 2024).
- Utama, M.Z. 2015. Budidaya Padi pada Lahan Marjinal Kiat Meningkatkan Produksi Padi. CV Andi Offset : Yogyakarta.
- Yin, N.-W.; Li, J.-N.; Liu, X.; Lian, J.-P.; Fu, C.; Li, W.; Jiang, J.-Y.; Xue, Y.-F.; Wang, J.; Chai, Y.-R. Lignification Response and the Difference between Stem and Root of *Brassica napus* under Heat and Drought Compound Stress. *Acta Agron. Sin.* 2017, 43, 1689.
- Yoshida. 1981. Fundamentals of Rice Crop Science. Internasional Rice Research Institute. Los Banos Phillipiness.
- Xiao, S.; Liu, L.; Zhang, Y.; Sun, H.; Zhang, K.; Bai, Z.; Dong, H.; Li, C. Fine root and root hair morphology of cotton under drought stress revealed with RhizoPot. *J. Agron. Crop Sci.* 2020, 206, 679–693.