

## DAFTAR PUSTAKA

- Abdurrasyid, Z., Astawan, M., Wresdiyati, T., Nurtama, B., & Sirait, Y.I.S. (2021). Mutu fisikokimia dan sensori minuman serbuk tempe. *PANGAN*, 30(2), 117–128.
- Abdurrasyid, Z., Astawan, M., Lioe, H.N., & Wresdiyati, T. (2023). Physicochemical and antioxidant properties of germinated soybean tempe after two days additional fermentation time. *Biointerface Research in Applied Chemistry*, 13(3), 1-14. <https://doi.org/10.33263/BRIAC133.238>.
- Adawiyah, D.R., Tjiptoputri, O.M., & Lince. (2020). Profil sensori sediaan pemanis dengan metode Rate-All-That-Apply (RATA). *Jurnal Mutu Pangan*, 7(1), 38–45. <https://doi.org/10.29244/jmpi.2020.7.1.38>.
- Adnan, A.A. (2006). *Karakterisasi fisiko kimia dan mekanis kelobot jagung sebagai bahan kemasan*. [Institut Pertanian Bogor].
- Agranoff, J. ed. (2001). *The complete handbook of tempe*. Second ed. Singapore. American Soybean Association.
- Alaydrus, A.Z.A. (2022). Analisa kuantitas sistem proses pembuatan es krim berbahan tambahan tepung tempe kedelai. *Journal of Scientech Research and Development*, 4(2), 179–191. <http://idm.or.id/JSCR>.
- Alfanesa, R., Rahayuni, T., & Hartani, L. (2021). Pengaruh jenis kemasan terhadap sifat organoleptik dan kimiawi tempe biji karet. *Jurnal Sains Pertanian Equator*, 10(2), 1–17.
- Aliyah, N. (2024). Pembuatan tempe kacang tunggak (*Vigna unguiculanta*) dengan variasi lama fermentasi menggunakan inokulum tempe. *Journal of Comprehensive Science*, 3(1), 62–78.
- Aminah, S., & Hersoelistyorini, W. (2012). Karakteristik kimia tepung kecambah serelia dan kacang-kacangan dengan variasi blanching. *Jurnal Universitas Muhammadiyah Semarang*, 1(1), 1–9.
- Andriani, M., Katri, B., & Nurhartadi, E. (2014). Studies on physicochemical and sensory characteristics of overripe tempeh flour as food seasoning. *Academic Research International*, 5(5), 36–45.
- Ang, H.G., Nga, B.H., & Lim, K.K. ed. (1989). *7<sup>th</sup> World Congress of Food Science and Technology*. Singapore. Singapore Institute of Food Science and Technology.
- Anggraini, R., & Permatasari, N.D. (2018). Pengaruh lubang perforasi dan jenis plastik kemasan terhadap kualitas sawi hijau (*Brassica juncea* L.). *Jurnal Penelitian Pascapanen Pertanian*, 14(3), 154–162. <https://doi.org/10.21082/jpasca.v14n3.2017.154-162>.
- AOAC. (1995). *Official methods of analysis (16<sup>th</sup> ed.)*. Washington, DC. Association of Official Analytical Chemists.

- Ares, G., Bruzzone, F., Vidal, L., Cadena, R.S., Giménez, A., Pineau, B., Hunter, D.C., Paisley, A.G., & Jaeger, S.R. (2014). Evaluation of a rating-based variant of check-all-that-apply questions: Rate-all-that-apply (RATA). *Food Quality and Preference*, 36, 87–95. <https://doi.org/10.1016/j.foodqual.2014.03.006>.
- Astawan, M., Cahyani, A., & Wresdiyati, T. (2023). Antioxidant activity and isoflavone content of overripe Indonesian tempe. *Food Research*, 7(1), 42–50.
- Astawan, M., Mardhiyyah, Y.S., & Wijaya, C.H. (2018). Potential of bioactive components in tempe for the treatment of obesity. *Jurnal Gizi Dan Pangan*, 13(2), 79–86. <https://doi.org/10.25182/jgp.2018.13.2.79-86>.
- Astawan, M., Wresdiyati, T., & Ichsan, M. (2016). Karakteristik fisikokimia tepung tempe kecambah kedelai. *Jurnal Pangan dan Gizi*, 11(1), 35–42. <http://journal.ipb.ac.id/index.php/jgizipangan/article/download/13167/9919>.
- Astuti, N.P. (2009). *Sifat organoleptik tempe kedelai yang dibungkus plastik, daun pisang dan daun jati*. [Universitas Muhammadiyah Surakarta].
- Aýun, Q., Suryani, S., & Kurnia, C. (2022). Identifikasi kapang pada tempe bungkus daun pisang dan plastik asal pengrajin tempe Jatiasih, Bekasi. *Bioed: Jurnal Pendidikan Biologi*, 10(2), 45–51.
- Baião, L.F., Rocha, C., Lima, R.C., Valente, L.M.P., & Cunha, L.M. (2022). Development of a Rate-All-That-Apply (RATA) ballot for sensory profiling of sea urchin (*Paracentrotus lividus*) gonads. *Food Research International*, 153, 1–10. <https://doi.org/10.1016/j.foodres.2022.110976>.
- Bavia, A.C.F., da Silva, C.E., Ferreira, M.P., Santos Leite, R., Mandarino, J.M.G., & Carrão-Panizzi, M.C. (2012). Chemical composition of tempeh from soybean cultivars specially developed for human consumption. *Ciencia e Tecnologia de Alimentos*, 32(3), 613–620. <https://doi.org/10.1590/S0101-20612012005000085>.
- Belgis, M., Arifin, T.Z., Prameswari, D., & Taruna, I. (2023). Sensory profile on robusta coffee by Rate-All-That-Apply (RATA). *Pelita Perkebunan*, 39(1), 32–42. <https://doi.org/10.22302/iccri.jur.pelitaperkebunan.v39i1.546>.
- Belitz, H.D., Grosch, W., & Schieberle, P. (2009). 5 Aroma compounds. In *Food Chemistry*. 340–402.
- Boekel, M.A.J.S.V. (2006). Formation of flavour compounds in the Maillard reaction. *Biotechnology Advances*, 24(2), 230–233. <https://doi.org/10.1016/j.biotechadv.2005.11.004>.
- Brooker, D.B., Bakker-Arkema, F.W., & Hall, C.W. (1980). *Drying cereal grains*. Westport, Connecticut. The AVI Publishing Company, Inc.
- BSN (Badan Standarisasi Nasional). (1995). SNI: 01-3709-1995. Rempah-rempah bubuk.

- BSN (Badan Standarisasi Nasional). (1998). SNI: 01-4476-1998. Tepung bumbu.
- Chen, L., Zeng, W., Rong, Y., & Lou, B. (2021). Characterisation of taste-active compositions, umami attributes and aroma compounds in Chinese shrimp. *International Journal of Food Science and Technology*, 56(12), 6311–6321. <https://doi.org/10.1111/ijfs.15304>.
- Danso-Boateng, E. (2013). Effect of drying methods on nutrient quality of Basil (*Ocimum viride*) leaves cultivated in Ghana. *International Food Research Journal*, 20(4), 1569–1573.
- Dewi, N.M.I.K., Suparthana, I.P., & Pratiwi, I.D.P.K. (2021). Evaluasi profil sensori abon ikan jenis pelagis besar menggunakan metode Rate-All-That-Apply (RATA). *ITEPA: Jurnal Ilmu Dan Teknologi Pangan*, 10(1), 324–336. <https://doi.org/10.24843/itepa.2021.v10.i03.p01>.
- Dwitama, F.S. (2018). *Profiling karakteristik sensoris kopi hijau dampit berdasarkan perbedaan metode pengolahan pasca panen dan teknik penyeduhan menggunakan metode Rate-All-That-Apply (RATA)*. [Universitas Brawijaya].
- Efendi, Z., Surawan, F.E.D., & Winarto. (2015). Efek blanching dan metode pengeringan terhadap sifat fisikokimia tepung ubi jalar orange (*Ipomoea batatas* L.). *Jurnal Agroindustri*, 5(2), 109-117.
- Efriwati & Nuraida, L. (2013). Effect of two production methods on macro nutrient and isoflavone-aglycone composition in tempeh produced by household industries. *Health Science Indonesian*, 4(2), 69–73.
- Ellent, S.S.C., Dewi, L., & Tapilouw, M.C. (2022). Karakteristik mutu tempe kedelai (*Glycine max* L.) yang dikemas dengan klobot. *AGRITEKNO: Jurnal Teknologi Pertanian*, 11(1), 32–40. <https://doi.org/10.30598/jagritekno.2022.11.1.32>.
- Fadel, H.H.M., Lotfy, S.N., Asker, M.M.S., Mahmoud, M.G., & Al-Okbi, S.Y. (2018). Nutty-like flavor production by *Corynebacterium glutamicum* 1220T from enzymatic soybean hydrolysate. Effect of encapsulation and storage on the nutty flavoring quality. *Journal of Advanced Research*, 10, 31–38. <https://doi.org/10.1016/j.jare.2018.01.003>.
- Feng, X.M., Larsen, T.O., & Schnurer, J. (2007). Production of volatile compounds by *Rhizopus oligosporus* during soybean and barley tempeh fermentation. *International Journal of Food Microbiology*, 113, 133–141. <https://doi.org/10.1016/j.ijfoodmicro.2006.06.025>.
- Fibrianto, K., & Dwihindarti, M. (2016). Profiling atribut jamu kunyit asam dan jamu sinom dengan metode RATA (Rate-All-That-Apply) pada beberapa kota di Jawa Timur. *J.Rekapangan*, 10(1), 15–22.
- Fuke, S., & Ueda, Y. (1996). Interactions between umami and other flavor characteristics. *Trends in Food Science and Technology*, 7(12), 407–411. [https://doi.org/10.1016/S0924-2244\(96\)10042-X](https://doi.org/10.1016/S0924-2244(96)10042-X).

- Fung, D.Y.C., & Crozier-Dodson, A. (2008). *Tempeh, a mold-modified indigenous fermented food*. Boca Raton, London. CRC Press.
- Gao, J., Fang, D., Muinde, B., Chen, X., Wu, X., Du, J., Yang, Q., Chen, H., Zheng, H., An, X., Zhao, L., & Hu, Q. (2021). Analysis of umami taste substances of morel mushroom (*Morchella sextelata*) hydrolysates derived from different enzymatic systems. *Food Chemistry*, 362, 1–11.
- Gunawan-Puteri, M.D.P.T., Hassanein, T.R., Prabawati, E.K., Wijaya, C.H., & Mutukumira, A.N. (2015). Sensory characteristics of seasoning powders from overripe tempeh, a solid state fermented soybean. *Procedia Chemistry*, 14, 263–269. <https://doi.org/10.1016/j.proche.2015.03.037>.
- Habibi, A. (2011). *Karakteristik sensori dan kimia tempe overripe (tempe "bosok") sebagai food seasoning (bumbu penyedap masakan)*. [Universitas Sebelas Maret].
- Hachmeister, K.A., & Fung, D.Y.C. (1993). Tempeh: A Mold-modified indigenous fermented food made from soybeans and/or cereal grains. *Critical Reviews in Microbiology*, 19(3), 137–188. <https://doi.org/10.3109/10408419309113527>.
- Haliza, W., Purwani, E. Y., & Thahir, R. (2007). Pemanfaatan kacang-kacangan lokal sebagai substitusi bahan baku tempe dan tahu. *Buletin Teknologi Pascapanen Pertanian*, 3, 1–8.
- Han, B.Z., Ma, Y., Rombouts, F.M., & Nout, M.J.R. (2003). Effects of temperature and relative humidity on growth and enzyme production by *Actinomucor elegans* and *Rhizopus oligosporus* during sufu pehtze preparation. *Food Chemistry*, 81, 27–34.
- Handoyo, T., & Morita, N. (2006). Structural and functional properties of fermented soybean (Tempeh) by using *rhizopus oligosporus*. *International Journal of Food Properties*, 9(2), 347–355. <https://doi.org/10.1080/10942910500224746>.
- Haron, H., & Raob, N. (2014). Changes in macronutrient, total phenolic and anti-nutrient contents during preparation of tempeh. *Journal of Nutrition & Food Sciences*, 4(2), 1–5. <https://doi.org/10.4172/2155-9600.1000265>.
- Hasni, D., Irfan, & Saputri, R. (2021). Pengaruh formulasi bahan baku dan CMC (Carboxy Methyl Cellulose) terhadap mutu dan penerimaan konsumen susu nabati. *Jurnal Teknologi dan Industri Pertanian Indonesia*, 13(2), 78–85. <https://doi.org/10.17969/jtipi.v13i2.21268>.
- Hastuti, S., Harijono, H., Murtini, E.S., & Fibrianto, K. (2018). Parametric and non-parametric approach for sensory RATA (Rate-All-That-Apply) method of ledre profile attributes. *IOP Conference Series: Earth and Environmental Science*, 131, 1–5. <https://doi.org/10.1088/1755-1315/131/1/012015>.
- Ikasari, L., & Mitchell, D.A. (1998). Mimicking gas and temperature changes during enzyme production by *Rhizopus oligosporus* in solid

- state fermentation. *Biotechnology Letters*, 20(4), 349–353. <https://doi.org/10.1023/A:1005319128217>.
- Irvan, Nabilah, Y., Prananta, L., Setiyaningsih, I., & Widi, P. (2021). Study of the effect of wrapping type and yeast concentration on the making of peanuts tempeh. *IOP Conference Series: Materials Science and Engineering*, 1122, 1–10. <https://doi.org/10.1088/1757-899x/1122/1/012091>.
- Jayanti, R.C.T. (2022). *Analisis profil sensori produk cokelat ready to drink dengan metode Rate-All-That-Apply (RATA) berdasarkan kesukaan konsumen*. [Universitas Sebelas Maret].
- Kaczmarek, K.T., Chandra-Hioe, M.V., Frank, D., & Arcot, J. (2018). Aroma characteristics of lupin and soybean after germination and effect of fermentation on lupin aroma. *LWT - Food Science and Technology*, 87, 225–233. <https://doi.org/10.1016/j.lwt.2017.08.080>.
- Kadaryati, S., Arinanti, M., & Afriani, Y. (2021). Formulasi dan uji sensori produk bumbu penyedap berbasis jamur tiram (*Pleurotus ostreatus*). *AgriTECH*, 41(3), 285–293.
- Kainuma, K., Odat, T., & Cuzuki, S. (1967). Study of starch phosphates monoesters. *Journal of Technology Society Starch*, 14, 24–28.
- Kato, H., Rhue, M.R., & Nishimura, T. (1989). *Role of free amino acids and peptides in food taste*. In Flavor chemistry trends and developments (pp. 158–174). American Chemical Society. <https://doi.org/10.1021/bk-1989-0388.ch013>.
- Kementerian Kesehatan. (2018). Data Komposisi Pangan Indonesia: Kacang kedelai, segar.
- Kementerian Kesehatan. (2018). Data Komposisi Pangan Indonesia: Tempe kedelai murni, mentah.
- Kibar, E., Gonenc, I., & Us, F. (2010). Gelatinization of waxy, normal and high amylose corn starches. *Research*, 35(4), 237–244.
- Kim, S.Y., Kim, B.M., Kim, J.B., Shanmugavelan, P., Kim, H.W., Kim, S.Y., Kim, S.N., Cho, Y.S., Choi, H.S., & Park, K.M. (2014). Effect of steaming, blanching, and high temperature/high pressure processing on the amino acid contents of commonly consumed Korean vegetables and pulses. *Preventive Nutrition and Food Science*, 19(3), 220–226. <https://doi.org/10.3746/pnf.2014.19.3.220>.
- Kovac, B., & Raspor, P. (1997). The use of the mould *rhizopus oligosporus* in food production. *Food technol. Biotechnol*, 35(1), 69–73.
- Kuligowski, M., Pawłowska, K., Jasińska-Kuligowska, I., & Nowak, J. (2016). Isoflavone composition, polyphenols content and antioxidative activity of soybean seeds during tempeh fermentation. *CYTA - Journal of Food*, 15(1), 27–33. <https://doi.org/10.1080/19476337.2016.1197316>.
- Kun, J., Meng, Q., Wei, C.C., Xie, G., Yan, J., Ho, C.T., & Tong, H. (2022). Characterization of the key compounds responsible for the fermented



- soybean-like cup aroma of raw Pu-erh tea using instrumental and sensory methods. *Lwt.* 162, 1–8. <https://doi.org/10.1016/j.lwt.2022.113458>.
- Leksrisonpong, P., Gerard, P., Lopetcharat, K., & Drake, M. (2012). Bitter taste inhibiting agents for whey protein hydrolysate and whey protein hydrolysate beverages. *Journal of Food Science*, 00(0), 1–6. <https://doi.org/10.1111/j.1750-3841.2012.02800.x>.
- Mahdi, S. A., Astawan, M., Wulandari, N., & Muhandri, T. (2023). Sensory profiling of tempe functional drink powder using rate-all-that-apply method. *Food Research*, 7(2), 19–26.
- Mardiah. (1992). *Mempelajari sifat fungsional dan nilai gizi tepung tempe serta pengembangan produk olahannya sebagai makanan tambahan bagi anak*. [Institut Pertanian Bogor].
- Martha, E. A., Jariyah, & Rosida. (2022). Pengaruh penambahan guar gum dan minyak sawit merah terhadap karakteristik fisikokimia mellorine sari tempe. *AGRITEPA: Jurnal Ilmu Dan Teknologi Pertanian*, 9(1), 53–68.
- Mastika, A.C., Praptiningsih, Y., & Fauzi, M. (2011). Karakteristik sambal tumpang dari campuran tempe segar dan tempe terfermentasi lanjut selama penyimpanan suhu rendah. *Berkala Ilmiah Pertanian*, 10, 1–5.
- Mastuti, T. S., & Handayani, R. (2014). Senyawa kimia penyusun ekstrak ethyl asetat dari daun pisang batu dan ambon hasil distilasi air. *Prosiding SNST Ke-5 Tahun 2014 Fakultas Teknik Universitas Wahid Hasyim Semarang*, 60–64.
- Mathatheeranan, P., Wongprasert, T., Wang, Y., Fang, M., Lu, T.J., & Suppavorasatit, I. (2024). Characterization of potent odorants in northern Thai fermented black and yellow soybeans (thua nao) using solvent-assisted flavor evaporation, aroma extract dilution analysis techniques and omission studies of aroma reconstitution models. *Journal of Food Composition and Analysis*, 128, 1–12. <https://doi.org/10.1016/j.jfca.2024.106031>.
- Mau, J.L. (2005). The umami taste of edible and medicinal mushrooms. *International Journal of Medicinal Mushrooms*, 7(1–2), 119–125. <https://doi.org/10.1615/IntJMedMushr.v7.i2.120>.
- Metty, Inayah, Widiyanti, F.L., & Shayida, H. (2022). Analisis kesukaan tepung tempe semangit berbahan dasar kacang kedelai import dan lokal sebagai bumbu penyedap. *Prosiding Seminar Nasional Universitas Respati Yogyakarta*, 4(1), 49–53.
- Mital, B.K., & Steinkraus, K.H. (1975). Utilization of oligosaccharides during fermentation by lactic acid bacteria. *Journal of Food Science*, 40, 114–118.
- Mufidah, I., Azizah, N., & Darni, J. (2018). Analisis perbedaan jenis pembungkus terhadap kadar proksimat dan daya terima tempe biji

- lamtoro (*Leucaena leucocephala*). *Darussalam Nutrition Journal*, 2(2), 21–31.
- Mujiyanto. (2013). Analisis faktor yang mempengaruhi proses produksi tempe produk UMKM di Kabupaten Sidoarjo. *REKA Agroindustri*, 1(1), 1–8. <http://ejournal.uwks.ac.id/detailjurnal.aspx?v=201310540413349173&x=1>.
- Mukherjee, S., & Chattopadhyay, P.K. (2007). Whirling bed blanching of potato cubes and its effects on product quality. *Journal of Food Engineering*, 78, 52–60. <https://doi.org/10.1016/j.jfoodeng.2005.09.001>.
- Mulyani, S. (2013). *Karakterisasi tepung tempe dari empat varietas kedelai impor dan aplikasinya menjadi minuman*. [Institut Pertanian Bogor].
- Mulyowidarso, R., Fleet, G., & Buckle, K. (1991). Changes in the concentration of organic acids during the soaking of soybeans for tempe production. *International Journal of Food Scien*, 26, 607–614.
- Munarko, H., Jariyah, J., & Kurnianto, M.A. (2023). Profiling atribut sensori kukis nastar menggunakan metode Rate-All-That-Apply (RATA). *Gorontalo Agriculture Technology Journal*, 6(2), 55–64. <https://doi.org/10.32662/gatj.v0i0.2711>.
- Murdijati-Gardjito, Indrati, R., Yuniarti, Z., & Hendrasty, H.K. (2019). *Gastronomi Indonesia Jilid 2*. Yogyakarta. Global Pustaka Utama.
- Nadilla. (2018). *Pengaruh penggunaan daun jati (Tectona grandis) terhadap lama fermentasi dan uji organoleptik tempe sebagai penunjang mata kuliah bioteknologi*. [Universitas Islam Negeri Ar-Raniry].
- Nedele, A.K., Gross, S., Rigling, M., & Zhang, Y. (2021). Reduction of green off-flavor compounds: Comparison of key odorants during fermentation of soy drink with *Lycoperdon pyriforme*. *Food Chemistry*, 334, 1–10. <https://doi.org/10.1016/j.foodchem.2020.127591>.
- Nout, M.J.R. & Rombouts, F.M. (1990). A review: Recent developments in tempeh research. *Journal of Applied Bacteriology*, 69, 609–633.
- Nuraini, V., Puyanda, I.R., Kunciaty, W.A.S., & Margareta, L.A. (2022). Perubahan kimia dan mikrobiologi tempe busuk selama fermentasi. *Jurnal Agroteknologi*, 15(02), 127–137. <https://doi.org/10.19184/j-agt.v15i02.25729>.
- Nurmadhani, B.S. (2017). *Pengaruh jenis bahan pengemas terhadap karakteristik sensoris dan kenampakan tempe kedelai lokal*. [Universitas Gadjah Mada].
- Nurrahman, Astuti, M., Suparmo, & Soesatyo, M.H.N.E. (2012). Pertumbuhan jamur, sifat organoleptik dan aktivitas antioksidan tempe kedelai hitam yang diproduksi dengan berbagai jenis inokulum. *AgriTECH*, 32(1), 60–65. <https://doi.org/10.22146/agritech.9657>.
- Nowak, J., & Szebiotko, K. (1992). Some biochemical changes during soybean and pea tempeh fermentation. *Food Microbiology*, 9, 37–43. [https://doi.org/10.1016/0740-0020\(92\)80060-H](https://doi.org/10.1016/0740-0020(92)80060-H).

- Oda, Y., Yajima, Y., Kinoshita, M., & Ohnishi, M. (2003). Differences of *Rhizopus oryzae* strains in organic acid synthesis and fatty acid composition. *Food Microbiology*, 20, 371–375. [https://doi.org/10.1016/S0740-0020\(02\)00131-4](https://doi.org/10.1016/S0740-0020(02)00131-4).
- Olanipekun, B.F., Otunola, E.T., Adelakun, O.E., & Oyelade, O.J. (2009). Effect of fermentation with *Rhizopus oligosporus* on some physico-chemical properties of starch extracts from soybean flour. *Food and Chemical Toxicology*, 47(7), 1401–1405. <https://doi.org/10.1016/j.fct.2009.02.023>.
- Owens, J. D. (2014). *Indigenous fermented foods of southeast Asia*. In *Indigenous Fermented Foods of Southeast Asia*. <https://doi.org/10.1201/b17835>.
- Pamungkas, E.M.P., Dewi, L., & Tapilouw, M.C. (2022). Penambahan angkak (*monascus purpureus*) pada tempe dalam peningkatan antioksidan. *Teknologi Pangan: Media Informasi dan Komunikasi Ilmiah Teknologi Pertanian*, 13(2), 144–155. <https://doi.org/10.35891/tp.v13i2.2923>.
- Pangastuti, A., Alfisah, R.K., Istiana, N.I., Sari, S.L.A., Setyaningsih, R., Susilowati, A., & Purwoko, T. (2019). Metagenomic analysis of microbial community in over-fermented tempeh. *Biodiversitas*, 20(4), 1106–1114. <https://doi.org/10.13057/biodiv/d200423>.
- Peng, X., Li, X., Shi, X., & Guo, S. (2014). Evaluation of the aroma quality of Chinese traditional soy paste during storage based on principal component analysis. *Food Chemistry*, 151, 532–538. <https://doi.org/10.1016/j.foodchem.2013.11.095>.
- Peng, X., Wang, Y., Xing, J., Wang, R., Shi, X., & Guo, S. (2017). Food Hydrocolloids Characterization of particles in soymilks prepared by blanching soybeans and traditional method: A comparative study focusing on lipid-protein interaction. *Food Hydrocolloids*, 63, 1–7.
- Pradana. (2019). *Korelasi kadar protein dan lemak terhadap cita rasa tempe kedelai (Glycine max L.) goreng yang beredar di kota Malang*. [Universitas Brawijaya].
- Pramita, A.R.I. (2012). *Peningkatan kestabilan larutan dan penambahan cocoa bubuk pada pembuatan formula tepung tempe*. [Universitas Hasanuddin].
- Prativi, M.B.N., Astuti, D.I., Putri, S.P., Laviña, W.A., Fukusaki, E., & Aditiawati, P., (2023). Metabolite changes in indonesian tempe production from raw soybeans to over-fermented tempe. *Metabolites*, 13(2), 1–16. <https://doi.org/10.3390/metabo13020300>.
- Prinyawiatkul, W., Beuchat, L.R., McWatters, K.H., & Phillips, R.D. (1996). Changes in fatty acid, simple sugar, and oligosaccharide content of cowpea (*Vigna unguiculata*) flour as a result of soaking, boiling, and fermentation with *Rhizopus microsporus* var. *oligosporus*. *Food*



- Chemistry*, 57(3), 405–413. [https://doi.org/10.1016/0308-8146\(95\)00242-1](https://doi.org/10.1016/0308-8146(95)00242-1).
- Pudjiastuti, W., Listyarini, A., & Rizki, M.I. (2013). Pengaruh laju transmisi uap air polymer blend polibutilen suksinat (PBS) dan linear low density polyethylene (LLDPE) terhadap umur simpan sup krim instan rasi. *Jurnal Kimia Kemasan*, 35(1), 1–5.
- Purwaningsih & Wanita, Y.P. (2013). *Kacang tanah sebagai alternatif pengganti bahan baku pada usaha mikro kecil menengah tempe di Gunungkidul*. Balai Pengkajian Teknologi Pertanian Yogyakarta.
- Putri, F.L. & Kartikawati, D. (2022). Optimasi konsentrasi ragi dan jenis pembungkus dalam pembuatan tempe kacang tunggak (*Vigna unguiculata* (L.) Walp.). *Jurnal Agrifoodtech*, 1(2), 103–118.
- Putri, B.D., Widyastuti, S., & Werdiningsih, W. (2018). Tempe kacang komak dengan beberapa pembungkus yang berbeda selama fermentasi. *Pro Food (Jurnal Ilmu Dan Teknologi Pangan)*, 4(2), 343–350. <https://doi.org/10.29303/profood.v4i2.86>.
- Rachmawati, M., Soetjipto, H., & Kristijanto, A. (2019). Profil asam lemak minyak tempe busuk. *Jurnal Kimia*, 13(1), 82–87.
- Radiati, A., & Sumarto. (2016). Analisis sifat fisik, sifat organoleptik, dan kandungan gizi pada produk tempe dari kacang non-kedelai. *Jurnal Aplikasi Teknologi Pangan*, 5(1), 16–22. <https://doi.org/10.17728/jatp.v5i1.32>.
- Rahayu, E.S., Indrati, R., Utami, T., Harmayani, E., & Cahyanto, M.N. (1993). *Bahan pangan hasil fermentasi*. Yogyakarta. PAU Pangan & Gizi.
- Rahayu, W.P., Pambayun, R., Santoso, U., Nuraida, L., & Ardiansyah. (2015). *Tinjauan ilmiah teknologi pengolahan tempe kedelai*. Perhimpunan Ahli Teknologi Pangan Indonesia.
- Rahmadian, Y., Supriyadi, Santoso, U., Mahmudah, N.A., & Ichsan, O.A.N. (2019). Non-volatile taste components and amino acid profile of jengkol (*Pithecellobium jiringa*) seed flour after steam blanching. *International Journal of Food Properties*, 22(1), 1536–1547. <https://doi.org/10.1080/10942912.2019.1657445>.
- Ramadhani, A.R. (2015). *Karakteristik organoleptik bubuk flavor kepala ikan tenggiri dengan bahan pengisi tepung terigu*. [Universitas Padjajaran].
- Ribeiro, L., Miguel, M., Evangelista, S., Martins, P., Mullem, J., Belizario, M., & Schwan, R. (2017). Behavior of yeast inoculated during semi-dry coffee fermentation and the effect on chemical and sensorial properties of the final beverage. *Food Research International*, 92, 26–32. <https://doi.org/10.1016/j.foodres.2016.12.011>.
- Rifqi, M., Rohmayanti, T., Sania, F.R., & Hapsari, U.D. (2023). Profil sensori pada roti tawar dengan penambahan tepung kulit buah naga merah

- dengan menggunakan metode Rate-All-That-Apply (RATA). *Jurnal Agroindustri Halal*, 9(3), 332–342.
- Rinawan, B.E. (2015). *Pengaruh macam pembungkus (daun dan plastik) terhadap profil pertumbuhan bakteri asam laktat dan bakteri proteolitik indigenous pada fermentasi tempe*. [Universitas Gadjah Mada].
- Rokhmah, L.N., Anam, C., Handajani, S., & Rachmawati, D. (2009). Kajian kadar asam fitat dan kadar protein selama pembuatan tempe kara benguk (*Mucuna pruriens*) dengan variasi pengecilan ukuran dan lama fermentasi. *Biofarmasi*, 7(1), 1–9. <https://doi.org/10.13057/biofar/f070101>.
- Roland, W.S.U., Vincken, J.P., Gouka, R.J., Van Buren, L., Gruppen, H., & Smit, G., (2011). Soy isoflavones and other isoflavonoids activate the human bitter taste receptors hTAS2R14 and hTAS2R39. *Journal of Agricultural and Food Chemistry*, 59(21), 11764–11771. <https://doi.org/10.1021/jf202816u>.
- Romulo, A., & Surya, R. (2021). Tempe: A traditional fermented food of Indonesia and its health benefits. *International Journal of Gastronomy and Food Science*, 26, 1–9. <https://doi.org/10.1016/j.ijgfs.2021.100413>.
- Rosida, Prianto, A.D., Munarko, H., & Kurnianto, M.A. (2023). Variation of fermentation time of crystal guava leaf tea (*Psidium guajava* L.) on antioxidant activity and sensory profile with Rate-All-That-Apply method. *FOODSCITECH*, 6(1), 42–50.
- Rosyidi, M., & Fahrudin, A. (2022). Rancang bangun cabinet dryer dengan variasi kecepatan blower menggunakan pembakaran arang pada tanaman cabai. *Procedia of Engineering and Life Science*, 3, 1–10.
- Ruiz-Terán, F., & Owens, J.D. (1999). Fate of oligosaccharides during production of soya bean tempe. *Journal of the Science of Food and Agriculture*, 79(2), 249–252. [https://doi.org/10.1002/\(SICI\)1097-0010\(199902\)79:2<249:AID-JSFA180>3.0.CO;2-1](https://doi.org/10.1002/(SICI)1097-0010(199902)79:2<249:AID-JSFA180>3.0.CO;2-1).
- Saldivar, X., Wang, Y.J., Chen, P., & Mauromoustakos, A. (2010). Effects of blanching and storage conditions on soluble sugar contents in vegetable soybean. *LWT - Food Science and Technology*, 43, 1368–1372. <https://doi.org/10.1016/j.lwt.2010.04.017>.
- Salim, R. (2017). Analisis jenis kemasan terhadap kadar protein dan kadar air pada tempe. *Jurnal Katalisator*, 2(2), 106. <https://doi.org/10.22216/jk.v2i2.2531>.
- Santhirasegaram, V., George, D.S., Anthony, K.K., Singh, H.K.B., Saruan, N.M., Razali, Z., & Somasundram, C. (2016). Effects of soybean processing and packaging on the quality of commonly consumed local delicacy tempe. *Journal of Food Quality*, 39, 675–684. <https://doi.org/10.1111/jfq.12252>.
- Sarrette, M., Nout, M.J.R., Gervais, P., & Rombouts, F.M. (1992). Effect of water activity on production and activity of *Rhizopus oligosporus*

- polysaccharidases. *Applied Microbiology and Biotechnology*, 37, 420–425. <https://doi.org/10.1007/BF00180961>.
- Sayuti. (2015). Pengaruh bahan kemasan dan lama inkubasi terhadap kualitas tempe kacang gude sebagai sumber belajar IPA. *BIOEDUKASI (Jurnal Pendidikan Biologi)*, 6(2), 148–158. <https://doi.org/10.24127/bioedukasi.v6i2.345>.
- Setyaningsih, D., Anton, A., & Maya, P.S. (2010). *Analisis sensori untuk Industri Pangan dan Agro*. Bogor. Institut Pertanian Bogor Press.
- Seulalae, A.V., Prangdimurti, E., Adawiyah, D.R., & Nurjanah. (2023). Evaluasi tingkat keasinan relatif dan profil sensori garam rumput laut menggunakan metode magnitude estimation dan Rate-All-That-Apply (RATA). *JPHPI*, 26(1), 54–66.
- Sielberberg, M. (2012). *Chemistry: The Molecular Nature of Matter and Change 6th edn*. New York, N.Y., McGraw Hill.
- Silva, J.D., Prudencio, S., Carrão-Panizzi, M., Gregorut, C., Fonseca, F., & Mattoso, L. (2012). Study on the flavour of soybean cultivars by sensory analysis and electronic tongue. *International Journal of Food Science and Technology*, 47, 1630–1638. <https://doi.org/10.1111/j.1365-2621.2012.03013.x>.
- Sitanggang, A.B., Firdausi, N.Z., & Budijanto, S. (2021). Antioxidant activity of a mixture between water-soluble tempeh extract and whey powder that has undergone a Maillard reaction. *Future of Food: Journal on Food, Agriculture and Society*, 9(5), 1–12. <https://doi.org/10.17170/kobra-202110144897>.
- Soegiharto, I.S. (1995). *Mempelajari pembuatan cookies dengan substitusi tepung tempe*. [Institut Pertanian Bogor].
- Song, J.Y., An, G.H., & Kim, C.J. (2003). Color, texture, nutrient contents, and sensory values of vegetable soybeans [*Glycine max* (L.) Merrill] as affected by blanching. *Food Chemistry*, 83(1), 69–74. [https://doi.org/10.1016/S0308-8146\(03\)00049-9](https://doi.org/10.1016/S0308-8146(03)00049-9).
- Steinkraus, K.H. (1983). *Handbook of indigenous fermented foods*. New York, USA. Marcel Dekker, Inc.
- Suhartanti, P.D., Handajani, S., & Nandariyah. (2019). Physical characteristics of the seeds of soybean (*Glycine max*) varieties and the effect of fermentation time on the chemical characteristics of tempeh. *Cell Biology & Development*, 3(1), 13–18.
- Sumasto, C.Y., & Budiasih, S. (1997). Penelitian pengaruh pembungkus plastik terhadap kualitas tempe kedele dengan variasi jenis plastik dan lobang plastik. *Majalah Barang Kulit, Karet dan Plastik*, 12(25), 48–53.
- Sun, L., Zhang, Z., Xin, G., Sun, B., Bao, X., Wei, Y., Zhao, X., & Xu, H. (2020). Advances in umami taste and aroma of edible mushrooms. *Trends in Food Science and Technology*, 96, 176–187. <https://doi.org/10.1016/j.tifs.2019.12.018>.

- Suprpto. (2004). Pengaruh lama blanching terhadap kualitas stik ubi jalar (*Ipomoea batatas* L.) dari tiga varietas. *Prosiding Temu Teknis Nasional Tenaga Fungsional Pertanian Tahun 2004*, 220-228.
- Suprihatin. (2010). *Teknologi Fermentasi*. UNESA Press. Surabaya.
- Suprijono, M., & Sutedja, A. (2008). Efek metode blanching uap dalam pembuatan biskuit tepung tempe terhadap penerimaan konsumen. *Prosiding Seminar Nasional: Peningkatan Keamanan Pangan Menuju Pasar Global*, 152–159. <http://repository.wima.ac.id/8393/1/%286%29>.
- Supriyono. (2003). *Memproduksi tempe*. Jakarta. Direktorat Pendidikan Menengah Kejurusan. Departemen Pendidikan Nasional.
- Sweers, L.J.H., Keppler, J.K., Feng, S., Zea, J.A., Veen, H.V.B. De, Timmermans, R. A.H., Boom, R.M., Fogliano, V., Lakemond, C.M.M., & Mishyna, M. (2024). High hydrostatic pressure for decontamination of soluble insect proteins prevents protein denaturation better than blanching. *Innovative Food Science and Emerging Technologies*, 96(103743).
- Syukri, D., Silvy, D., & Ramadani, S.F. (2022). Effect of various cooking methods on quality and sensory characteristics of tempeh made from soybeans and corn. *And. Int. J. Agric. Nat. Sci.* 3(2), 87–113.
- Tanuwidjaja, L., Hariyadi, H.R., & Mulyati, Y. (1991). Faktor-faktor yang mempengaruhi kualitas tempe di Indonesia. *Jurnal Ekonomi Pertanian dan Agribisnis (JEPA)*, 1(1), 31–34.
- Taylor, M., Hershey, H., Levine, R., Coy, K., & Olivelle, S. (1981). Improved method of resolving nucleotides by reversed-phase high-performance liquid chromatography. *Journal of Chromatography*, 219(1), 133–139.
- Tikk, M., Tikk, K., Tørngren, M.A., Meinert, L., Aaslyng, M.D., Karlsson, A.H., & Andersen, H.J. (2006). Development of inosine monophosphate and its degradation products during aging of pork of different qualities in relation to basic taste and retronasal flavor perception of the meat. *Journal of Agricultural and Food Chemistry*, 54, 7769–7777. <https://doi.org/10.1021/jf060145a>.
- Trifunovic, N. (2002). *Small community water supplies/IRC technical paper series 40*. Ed. Smet, J & van Wijk, C. Chapter 20: Water transmission. The Netherlands. Delft.
- Tseng, Y.H., Lee, Y.L., Li, R.C., & Mau, J.L. (2005). Non-volatile flavour components of *Ganoderma tsugae*. *Food Chemistry*, 90, 409–415. <https://doi.org/10.1016/j.foodchem.2004.03.054>.
- Umami, S., Jaya, I.K.S., Darawati, M., & Widiada, I.G.N. (2018). Kajian sifat organoleptik dan masa simpan tempe kedelai dengan beberapa jenis kemasan. *Jurnal Gizi Prima*, 3(2), 142–148.
- Utami, R., Wijaya, C.H., & Lioe, H.N. (2016). Taste of water-soluble extracts obtained from over-fermented tempe. *International Journal of Food Properties*, 19(9), 2063–2073. <https://doi.org/10.1080/10942912.2015.1104509>.

- Varzakas, T., Mahn, A., Perez, C., Miranda, M., & Barrientos, H. (2015). *Blanching*. In T. Varzakas & C. Tzia (Eds.), *Handbook of Food Processing: Food Preservation* (pp. 1–26). CRC Press. <https://doi.org/10.1201/b19397-2>.
- Vidal, L., Jaeger, S.R., Antúnez, L., Giménez, A., & Ares, G. (2016). Product spaces derived from projective mapping and CATA questions: Influence of replicated assessments and increased number of study participants. *Journal of Sensory Studies*, 31(5), 373–381. <https://doi.org/10.1111/joss.12220>.
- Wang, T.Y., Li, Q., & Bi, K.S. (2018). Bioactive flavonoids in medicinal plants: Structure, activity and biological fate. *Asian Journal of Pharmaceutical Sciences*, 13, 12–23. <https://doi.org/10.1016/j.ajps.2017.08.004>.
- Watanabe, N., Endo, Y., Fujimoto, K., & Aoki, H. (2006). Tempeh-like fermented soybean (GABA-tempeh) has an effective influence on lipid metabolism in rats. *Journal of Oleo Science*, 55(8), 391–396. <https://doi.org/10.5650/jos.55.391>.
- Webb, J., Bolhuis, D.P., Cicerale, S., Hayes, J.E., & Keast, R. (2015). The relationships between common measurements of taste function. *Chemosensory Perception*, 8(1), 11–18. <https://doi.org/10.1007/s12078-015-9183-x>.
- Wihandini, D.A., Arsanti, L., & Wijanarka, A. (2012). Sifat fisik, kadar protein dan uji organoleptik tempe kedelai hitam dan tempe kedelai kuning dengan berbagai metode pemasakan. *Nutrisia*, 14(1), 34–43.
- Winarno, F.G. (2004). *Kimia pangan dan gizi*. Gramedia Pustaka Utama. Jakarta.
- Winarno, F.G., Winarno, W., & Winarno, A.D.A. (2017). *Tempe: kumpulan fakta menarik berdasarkan penelitian*. Gramedia. Jakarta.
- Wulandari, D., Komar, N., & Sumarlan, S.H. (2013). Perekayasaan pangan berbasis produk lokal Indonesia (studi kasus sosis berbahan baku tempe kedelai). *Jurnal Bioproses Komoditas Tropis*, 1(2), 73–82.
- Xie, C., Zeng, H., Li, J., & Qin, L. (2019). Comprehensive explorations of nutritional, functional and potential tasty components of various types of sufu, a Chinese fermented soybean appetizer. *Food Sci. Technol. Campinas*, 39(1), 105–114. <https://doi.org/10.1590/fst.37917>.
- Xiao, H., Pan, Z., Deng, L., El-Mashad, H., Yang, X., Mujumdar, A., Gao, Z., & Zhang, Q. (2017). Recent developments and trends in thermal blanching-a comprehensive review. *Information Processing in Agriculture*, 4, 101–127. <http://dx.doi.org/10.1016/j.inpa.2017.02.001>.
- Yamaguchi, S., Tomoko, Y., Shingo, I., & Tsunehiko, N. (1971). Measurement of the relative taste intensity of some L-a-amino acids and 5'-nucleotides. *Journal of Food Science*, 36, 846–849.
- Yang, J., Lin, H., & Mau, J. (2001). Non-volatile taste components of several commercial mushrooms. *Food Chemistry*, 72, 465–471.



- Yao, J., Zhao, W., Bai, X., Wan, P., Liu, J., & Chen, D.W. (2023). Non-volatile taste active compounds in the meat of river snail (*Sinotaia quadrata*) determined by <sup>1</sup>H NMR, e-tongue and sensory analysis. *International Journal of Gastronomy and Food Science*, 34, 1–9. <https://doi.org/10.1016/j.ijgfs.2023.100803>.
- Yue, J., Zhang, Y., Jin, Y., Deng, Y., & Zhao, Y. (2016). Impact of high hydrostatic pressure on non-volatile and volatile compounds of squid muscles. *Food Chemistry*, 194, 12–19. <https://doi.org/10.1016/j.foodchem.2015.07.134>.
- Zhang, L., Zhang, L., & Xu, Y. (2020). Effects of *Tetragenococcus halophilus* and *Candida versatilis* on the production of aroma-active and umami-taste compounds during soy sauce fermentation. *Journal of the Science of Food and Agriculture*, 100, 2782–2790. <https://doi.org/10.1002/jsfa.10310>.
- Zhao, C.J., Schieber, A., & Gänzle, M.G. (2016). Formation of taste-active amino acids, amino acid derivatives and peptides in food fermentations – A review. *Food Research International*, 89, 39–47. <https://doi.org/10.1016/j.foodres.2016.08.042>.
- Zioga, E., Tøstesen, M., Madsen, S.K., Shetty, R., & Bang-Berthelsen, C.H. (2022). Bringing plant-based Cli-meat closer to original meat experience: insights in flavor. *Future Foods*, 5, 1–13. <https://doi.org/10.1016/j.fufo.2022.100138>.