

## DAFTAR PUSTAKA

- Akmaluddin, A., Agustin, M.V., Kurniawan A.M. 2019. Stratigraphy and Foraminiferal Biostratigraphy of Sentolo Formation in Sedayu Area: Local Unconformity Identification in Early Pliocene. *Journal of Applied Geology.*, 3(2), pp.32.
- Anda, M., Sarwani, M., 2012. Mineralogy, chemical composition, and dissolution of fresh ash eruption: new potential source of nutrients. *Soil Sci. Soc. Am. J.* 76, 733–747.
- Anda, M., Suryani, E., Husnain, and Subardja, D. 2015. Strategy to reduce fertilizer application in volcanic paddy soils: nutrient reserves approach from parent materials. *Soil Tillage Res.* 150, 10–20.
- Barrow, N.J. 2022. How understanding soil chemistry can lead to better phosphate fertilizer practice: a 68 year journey (so far). *Plant Soil* 476, 117–131
- Blakemore, L.C., Searly, P.L., Daly, B.K., 1981. *Methods for Chemical Analysis of Soils.* NZ Bureau Scientific Report No. 80, Lower Hutt, New Zealand.
- BBSDLP. 2017a. Atlas Peta Tanah Semi Detail Skala 1:50.000, Kabupaten Sleman, Provinsi Daerah Istimewa Yogyakarta. Badan Penelitian dan Pengembangan Pertanian
- BBSDLP. 2017b. Atlas Peta Tanah Semi Detail Skala 1:50.000, Kabupaten Bantul, Provinsi Daerah Istimewa Yogyakarta. Badan Penelitian dan Pengembangan Pertanian
- Camus, G., A. Gourgaud, P.C. Mossand-Bernhommier, and P.M. Vincenn. 2000. Merapi (central Java, Indonesia): An outline of the structural and magmatological evolution, with a special emphasis to the major pyroclastic events. *J. Volcanol. Geotherm. Res.* 100:139–163.
- Caubet, M., Cornu, S., Saby, N.P., Meunier, J.-D. (2020). Agriculture increases the bioavailability of silicon, a beneficial element for crop, in temperate soils. *Sci. Rep.* 10, 1–11.
- Churchman, J. and Lowe, D.J. 2012. Alteration, formation, and occurrence of minerals in soils. In: Huang, P.M., Li. Y., Sumner Y.E. (eds), *Handbook of Soil Sciences.* 2nd Edition. Vol 1: Properties and Processes. CRC Press. Boca Raton, pp 20.1-20.72.
- Cornelis J-T, Dumon M, Tolossa AR, Delvaux B, Deckers J, Van Ranst E. 2014. The effect of pedological conditions on the sources and sinks of silicon in the vertic planosols in south western ethiopia. *Catena* 112:131–8.
- Coskun D, Deshmukh R, Sonah H, Menzies JG, Reynolds O, Ma JF, Kronzucker HJ, Be'langer RR. 2019. The controversies of silicon's role in plant biology. *New Phytol* 221:67–85. <https://doi.org/10.1111/nph.15343>.
- Darmawan, Kyuma K, Saleh A, Subagjo H, Masunaga T, Wakatsuki T 2006: Effect of long-term intensive rice cultivation on the available silica content of sawah soils: Java Island, Indonesia. *Soil Sci. Plant Nutr.*, 52, 745–753.

- Debona D, Rodrigues FA, Datnoff LE. Silicon's Role in Abiotic and Biotic Plant Stresses. *Annu Rev Phytopathol.* 2017.
- DeMaster DJ. 1981. The supply and accumulation of silica in the marine environments. *Geochim Cosmochim Acta* 45:1715–32.
- de Tombeur F, Turner BL, Laliberté E, Lambers H, Cornelis J-T. 2020. Silicon dynamics during 2 million years of soil development in a coastal dune chronosequence under a Mediterranean climate. *Ecosystems* 23:1614–1630.
- de Tombeur F, Cornelis JT, Laliberte E, Lambers H, Mahy G, Faucon MP, Turner BL. 2021. Impact of ecosystem water balance and soil parent material on silicon dynamics: insights from three long-term chronosequences. *Biogeochemistry* 156:335–350.
- Dietzel, M. Interaction of polysilicic and monosilicic acid with mineral surfaces. In *Water-Rock Interaction*; Stober, I., Bucher, K., Eds.; Springer: Dordrecht, The Netherlands, 2002; pp. 207–235.
- Gertisser R, Charbonnier SJ, Keller J, Quidelleur X (2012b) The geological evolution of Merapi volcano, Central Java, Indonesia. *Bull Volcanol* 74:1213–1233.
- Guntzer, F., Keller, C., Meunier, J.D., 2012. Benefits of plant silicon for crops: a review. *Agron. Sustain. Dev.* 32, 201–213.
- Haysom, M.B.C & Chapman, L.S., 1975. Some aspects of the calcium silicate trials at Mackay. *Proceedings of the Australian Sugar Cane Technology* 42, 117–122.
- Henriet, C., Bodarw' e, L., Dorel, M., Draye, X., Delvaux, B., 2008. Leaf silicon content in banana (*Musa* spp.) reveals the weathering stage of volcanic ash soils in Guadeloupe. *Plant Soil* 313, 71–82.
- Herbillon, A.J., 1986. Chemical estimation of weatherable minerals in the diagnostic horizons of the low activity clay soils. In: Beinroth, F.H., Camargo, M.N., Eswaran, H. (Eds.), *Proceedings of the Eight International Soil Classification Workshop: Classification, Characterization and Utilization of Oxisols, Part 1. EM-BRAPA, Rio de Janeiro*, pp. 39–48.
- Holmgren, G. G. S. (1967). A rapid citrate-dithionite extractable iron procedure. *Soil Science Society of America Journal*, 31, 210–211.
- Huang, Y. C., and Z. Y. Hseu. 2021. Silicon Availability in Relation to Soil Properties in Inceptisols on Uncultivated Lands and Paddy Fields in Taiwan. *Geoderma Regional* 26: e00406
- Husein, S., D. Karnawati, S. Pramumijoyo, and A. Ratdomopurbo (2007) Kontrol Geologi terhadap Respon Lahan dalam Gempabumi Yogyakarta 27 Mei 2006: upaya pembuatan peta zonasi mikro di daerah Bantul. *Proceeding Seminar Nasional 2007 Geotechnics for Earthquake Engineering, Universitas Katolik Parahyangan, Bandung*, pp. 6.1–6.12.
- Husnain Wakatsuki T, Setyorini D, Hermansah Sato K, Masunage T (2008) Silica availability in soils and river in two watersheds on Java Island, Indonesia. *Soil Sci Plant Nutr* 54:916–927

- Imaizumi K, Yoshidai S. Edaphological studies on the silicon supplying power of paddy soils. *Bull Natl Inst Agric Sci Tokyo*. 1958; B8:261–304.
- Ismangil and Hanudin, E. 2005. Degradasi mineral batuan oleh asam-asam organik. *Jurnal Tanah dan Lingkungan* 5(1): 1-17
- Kanivets, S. 2017. The Factors and Conditions of Soil Formation: A Critical Analysis of Equivalence. In: D. Dent, Y Dmytruk (Eds). *Soil Science Working for a Living Applications of soil science to present-day problems*. Springer Cham. 3–8
- Kawaguchi K, Kyuma K 1977: *Paddy Soil in Tropical Asia. Their Material, Nature and Fertility*. The University Press of Hawaii, Honolulu.
- Klotzbücher T, Marxen A, Vetterlein D, Schneiker J, Tuörke M, Sinh NV et al (2015a) Plant-available silicon in paddy soils as a key factor for sustainable rice production in Southeast Asia. *Basic Appl Ecol* 16:665–673
- Klotzbücher A, Klotzbücher T, Jahn R et al. 2018. Effects of Sifertilization on Si in soil solution, Si uptake by rice, and resistance of rice to biotic stresses in Southern Vietnam. *Paddy Water Environ* 16:243–252.
- Lindsay, W.L., 1979. *Chemical equilibria in soils*. John Wiley and Sons Ltd.
- Meharg C, Meharg AA (2015) Silicon, the silver bullet for mitigating biotic and abiotic stress, and improving grain quality, in rice? *Environ Exp Bot* 120:8–17
- Meunier, J.D.; Keller, C.; Guntzer, F.; Riotte, J.; Braun, J.J.; Anupama, K. Assessment of the 1% Na<sub>2</sub>CO<sub>3</sub> technique to quantify the phytolith pool. *Geoderma* 2014, 216, 30–35.
- Meunier, J.-D., Sandhya, K., Prakash, N.B., Borschneck, D., Dussouillez, P., 2018. pH as a proxy for estimating plant-available Si? A case study in rice fields in Karnataka (South India). *Plant Soil* 432, 143–155.
- Nesbitt H, Young G (1982) Early Proterozoic climates and plate motions inferred from major element chemistry of lutites. *Nature* 299:715–717
- Nordt LC, Driese SD (2010) New weathering index improves paleorainfall estimates from Vertisols. *Geology* 38:407–410
- Nurcholis, M. Isdiyanto, R. Krismawan, H. Satria F.A. 2023. Land suitability study for sweet sorghum (*Sorghum bicolor* (L.) Moench) farming in Pleret Yogyakarta. *OP Conf. Ser.: Earth Environ. Sci.* 1180 012001
- Nursyamsi D., K. Idris, S. Sabiham, D.A. Rachim, and A. Sofyan. 2008. Dominant Soil Characteristics Influencing Available Potassium on Smectitic Soils. *Indonesian Journal of Agriculture* 1(2): 121-131
- Nurwihastuti, D.W.; Sartohadi, J.; Mardiatno, D.; Nehren, U. 2014. Understanding of earthquake damage pattern through geomorphological approach: A case study of 2006 earthquake in Bantul, Yogyakarta, Indonesia. *World J. Eng. Technol.* 2, 61–70.

- Maryanto, S. (2012). Limestone diagenetic records based on petrographic data of Sentolo Formation at Hargorejo traverse, Kokap, Kulonprogo. *Indonesian Journal on Geoscience*, 7(2), 87-99.
- Mehra OP, Jackson ML. 1960. Iron oxide removal from soils and clays by a dithionite-citrate system buffered with sodium bicarbonate. *Clays Clay Miner* 7:317–27.
- Pandita, Hita & Hartono, Hill & Pambudi, Setyo & Winarti, Winarti. (2021). Depositional model of Volcaniclastic-Carbonate Facies of Sentolo Formation at Miocene. 10.4108/eai.30-8-2021.2311527.
- Rahardjo W, Sukandarrumidi, Rosidi HMD. (1995). Peta Geologi Lembar Yogyakarta Jawa (1408-2 dan 1407-5) Skala 1: 100.000. Pulitbang Geologi, Bandung.
- Rayes M.L. 2000. Karakteristik, Genesis, dan Klasifikasi Tanah Sawah Berasal dari Bahan Volkan Merapi. Disertasi. Institut Pertanian Bogor. Bogor
- Saputra, A.; Gomez, C.; Delikostidis, I.; Zavar-Reza, P.; Hadmoko, D.S.; Sartohadi, J.; Setiawan, M.A. Determining Earthquake Susceptible Areas Southeast of Yogyakarta, Indonesia—Outcrop Analysis from Structure from Motion (SfM) and Geographic Information System (GIS). *Geosciences* 2018, 8, 132.
- Schaller J, Puppe D, Kaczorek D, Ellerbrock R, Sommer M. 2021. Silicon cycling in soils re-visited. *Plants*. 10(2):295.
- Soil Survey Staff. 2014. Keys to Soil Taxonomy. 12nd ed. USDA Natural Resources Conservation Service. Washington DC.
- Sommer, M., Kaczorek, D., Kuzyakov, Y., Breuer, J., 2006. Silicon pools and fluxes in soils and landscapes—a review. *J. Plant Nutr. Soil Sci.* 169, 310–329.
- Sposito, G. 2008. *The Chemistry of Soils*. 2nd Ed. Oxford University Press, New York. 255 p.
- Sukarman, S. Ritung, M. Anda, dan E. Suryani. 2016. Pedoman Pengamatan Tanah di Lapangan. Balai Besar Penelitian dan Pengembangan Sumberdaya Lahan Pertanian, Badan Penelitian dan Pengembangan Pertanian. 136 hlm.
- Surono, T.B., Sudarno, I., 1992. Geological Map of the Surakarta–Giritontro Quadrangle. Geological Research and Development Centre, Indonesia, Bandung.
- Tavares, T.R., Nunes, L.C., Alves, E.E.N., de Almeida, E., Maldaner, L.F., Krug, F.J., de Carvalho, H.W.P., Molin, J.P., 2019. Simplifying sample preparation for soil fertility analysis by x-ray fluorescence spectrometry. *Sensors*, 19, 5066
- Tubana BS, Babu T, Datnoff LE (2016) A review of silicon in soils and plants and its role in US agriculture: history and future perspectives. *Soil Sci* 181:393–411
- Vander Linden, C., Delvaux, B., 2019. The weathering stage of tropical soils affects the soil-plant cycle of silicon, but depending on land use. *Geoderma* 351, 209–220.
- van Reeuwijk, L.P., 2002. Procedure for soil analyses. In: Technical Paper 9, 6th edition. ISRIC, Wageningen, The Netherlands, pp. 101.



- Wijayanti, H. D. K. (2022). Stratigrafi Kontak Formasi Semilir dan Ngglangran Pada Jalur Pilangrejo, Nglipar, Gunung Kidul. *JGE (Jurnal Geofisika Eksplorasi)*, 8(02), 137-151
- Wu, W.; Limmer, M.A.; Seyfferth, A.L. Quantitative assessment of plant-available silicon extraction methods in rice paddy soils under different management. *Soil Sci. Soc. Am. J.* 2020, 84, 618–626.

## LAMPIRAN

### Lampiran 1. Deskripsi Profil

P1

Location name: Triwidadi, Pajangan District, Bantul Regency

Vegetation: Teak Tree and grass

Land Use: Secondary Forest

Parent Material: Limestone and Marl

Geological age: Middle Miocene–Pliocene

Topography: Undulating

Soil Classification: Very Fine, Smectitic, Superactive, Isohyperthermic, Typic Haprendolls

### Morfologi Profil Tanah

Depth (cm)	Horizons	Soil Profile Description
0-21/34	A	Very dark brown (10YR 2/2) moist; gravelly clay; moderate, coarse, granular structure; firm, very plastic; many fine and medium roots; about 15% by volume gravel size limestone fragment; violently effervescent (HCl); moderately alkaline (pH 8); gradual, wavy boundary.
21/34-52/60	Bw/2Cr	Bw Part: very dark grayish brown (10YR 3/2) clay, dark grayish (7.5YR 4/1) moist; weak fine angular blocky; firm; very plastic; few fine and very fine roots; common pressure faces; about 7% by volume fine gravel size limestone fragment; strongly effervescent (HCl); moderately alkaline (pH 8);  2Cr part: Pale brown (2,5Y 8/2) silty loam, massive, friable, very plastic; very few very fine roots; violently effervescent (HCl); moderately alkaline (pH 8); clear, irregular boundary.
52/60-220+	R	Limestone bedrock with some cracks

P2

Location name: Argodadi, Sedayu District, Bantul Regency

Vegetation: paddy-chilli (rotation)

Land Use: upland paddy field

Parent Material: Colluvium Limestone-Aluvium

Geological age: Quaternary (?)

Topography: Gently sloping

Soil Classification: Very fine, smectitic, Superactive, Isohyperthermic, Mollic Fluvaquents

Morfologi Profil Tanah

Depth (cm)	Horizons	Soil Profile Description
0–18/22	Ap	Very dark gray (2.5Y 3/1) gravelly clay, moist; moderate, coarse, angular clods; firm, very plastic; many fine and medium roots; about 25% by volume fine gravel size limestone fragment; strongly effervescent; moderately alkaline (pH 8); clear wavy boundary.
18/22–75/82	Cg1	Gray (5Y 5/1) gravelly clay, moist; massive; friable, very plastic, very few very fine and fine roots; about 15% by volume coarse gravel size limestone fragment; strongly effervescent; moderately alkaline (pH 8); diffuse wavy boundary.
75/82–141	2Cg2	Dark Yellowish Brown (10YR 4/4) gravelly clay loam, Gray (2,5Y 6/1) moist; massive; friable, moderately plastic; very few very fine roots; 25% gravel; slightly effervescent; moderately alkaline (pH 8); gradual smooth boundary.
141–150	3Cg3	Gray (2.5Y 5/1) silty clay loam, moist; massive; friable, moderately plastic; 10% gravel; very slightly effervescent; moderately alkaline (pH 8).

P3

Location name: Sumberrahayu, Moyudan District, Bantul Regency

Vegetation: paddy

Land Use: lowland paddy field

Parent Material: Clay Alluvium

Geological age: Quaternary (?)

Topography: Nearly level

Soil Classification: Fine, smectitic, superactive, Isohyperthermic, Typic Epiaquerts

Morfologi Profil Tanah

Depth (cm)	Horizons	Soil Profile Description
0–22/25	Apg1	Very dark grayish Brown (2.5Y 3/2) clay, dark red (2,5YR 3/6) moist; massive; firm, very plastic; many fine and medium roots; effervescent; slightly acid (pH 6,5); gradual wavy boundary.
22/25–40/44	Apg2	Gray (2.5Y 5/1) clay, grayish brown (2.5Y 5/2), moist, many dark red (2,5YR 3/6) mottles; massive; firm, very plastic, few very fine roots; moderately alkaline (pH 8); clear wavy boundary.
43/48–80/84	2B(ss)g	Gray (5 Y 5/1) clay, dark gray (5Y 4/1) moist; weak, medium, angular blocky; firm, very plastic; few very fine roots; many stress surfaces; common fine prominent brownish yellow (10YR 6/8) masses of oxidized iron with clear boundaries lining pores; few pitted concretions of calcium carbonate; moderately alkaline (pH 8); clear wavy boundary.
80/84–102/110	2B(ss)g	Dark gray (5Y 4/1) clay, gray (5 Y 5/1) moist; strong; friable, weak, medium, wedge structure; very plastic; many slickensides; many yellowish brown (10YR 5/6) masses of oxidized iron with diffuse boundaries along slickenside faces; few pitted concretions of calcium carbonate; few very dark brown (10YR 2/2) iron-manganese nodules across; strongly alkaline (pH 8,5); clear wavy boundary.
102/110–115/120	2Bkg	Pale Yellow (2.5Y 8/2) clay coating with carbonate, moist; moderate fine angular blocky structure, firm, very plastic; violently effervescent (HCl) carbonate masses; moderately alkaline (pH 8); abrupt broken boundary
115/120–160+	3Cg	Gray (5Y 5/1) clay loam, light yellowish brown (7.5YR 6/4); massive; firm, plastic; many iron-manganese nodules across; 10% gravel; very slightly effervescent (HCl); moderately alkaline (pH 8)

P4

Location name: Margoluwih, Seyegan District, Sleman Regency

Vegetation: paddy

Land Use: lowland paddy field

Parent Material: Quaternary Volcanic Deposit

Geological age: Quaternary

Topography: Nearly level

Soil Classification: Fine-loamy, mixed, superactive, isohyperthermic, Typic Fragiaquepts

Morfologi Profil Tanah

Depth (cm)	Horizons	Soil Profile Description
0-22	Apg	Dark grayish Brown (10YR 4/2) loam, pale brown (10YR 6/3) especially in lower part, moist; massive; friable, slightly plastic; many fine and medium roots; moderately acid (pH 6); clear smooth boundary.
22-33	Ad	Very dark grayish Brown (10YR 3/2) sandy loam, light yellowish brown (10YR 6/4) iron oxide layer with 3-6 cm thick in lower part, moist; massive; extremely firm, slightly plastic; few fine roots; neutral (pH 7); clear smooth boundary.
33-47	Bg1	Very dark grayish brown (10YR 3/2) loam, moist; weak, fine, angular blocky; firm, slightly plastic; few fine roots; many fine, distinct, very dark brown (10YR 2/2) Mn mottles; neutral (pH 7); clear smooth boundary.
47-65	Bg2	Gray (10YR 4/1) sandy loam, moist; weak, fine, angular blocky; firm, slightly plastic; few fine roots; many fine, faint, dark brown (10YR 3/3) Mn mottles; neutral (pH 7); gradual smooth boundary.
65-78	Bg3	Gray (10YR 4/1) sandy loam, moist; weak, fine, angular blocky; very firm, slightly plastic; few fine roots; many fine to medium, distinct, dark brown (10YR 3/3) Fe/Mn masses, neutral (pH 7); abrupt smooth boundary
78-94	Bsm1	Weak red (2.5YR 4/2) to reddish brown (2.5YR 4/4) loam cemented by iron oxide, Gray (10YR 4/1) moist; massive; rigid, slightly plastic; few fine roots; neutral (pH 7); abrupt smooth boundary
94-110	Bsm2	Very dark grey (10YR 3/1) to dark brown (10YR 3/3) clay loam cemented by Mn/Fe oxide, reddish brown (2.5YR 4/4) moist; massive; rigid, slightly plastic; few fine roots; neutral (pH 7); diffuse smooth boundary
110-150	2Bg4	Dark brown (10YR 3/3), Very dark gray (10YR 3/1) clay loam moist; weak, fine, angular blocky; firm, slightly plastic; few fine roots; neutral (pH 7)

P5

Location name: Wukirsari, Imogiri District, Bantul Regency

Vegetation: Teak Tree, Grass

Land Use: Secondary Forest

Parent Material: Sandstone, Tertiary Volcanic Tuff

Geological age: Early Miocene-Middle Miocene

Topography: Undulating

Soil Classification: Fine, smectitic, superactive, isohyperthermic, Lithic Udorthents

Morfologi Profil Tanah

Depth (cm)	Horizons	Soil Profile Description
0-20	A	Dark brown (7.5YR 3/4) clay loam, moist; weak, fine, subangular blocky; firm, slightly plastic; many fine, medium, and coarse roots; moderately acid (pH 5,5); abrupt smooth boundary.
20-24/50	AC	Pale Brown (10YR 6/3) clay loam, pale red (2,5YR 4/4), Pale yellow (2,5Y 8/2), moist; massive; firm, slightly plastic; many fine, medium, and coarse roots; many iron oxide masses in root channels; moderately acid (pH 5,5); abrupt irregular boundary.
24/50+	R	Sandstone and Tuff bedrock

P6

Location name: Segoroyoso, Imogiri District, Bantul Regency

Vegetation: paddy

Land Use: lowland paddy field

Parent Material: Clay Alluvium

Geological age: Quaternary (?)

Topography: Nearly level

Soil Classification: Fine, smectitic, superactive, Isohyperthermic, Typic Epiaquerts

Morfologi Profil Tanah

Depth (cm)	Horizons	Soil Profile Description
0-27/30	Apg	Very dark gray (7.5YR 3/1) silty clay, dark gray (10YR 4/1) moist; moderate, medium, prismatic structure parting to moderate, coarse, subangular blocky structure; firm, very plastic; many fine and medium roots; common fine, distinct, brown (7,5YR 4/3) iron mottles; slightly acid (pH 6); clear wavy boundary.
27/30-56/62	Bg	Very dark grayish brown (10YR 3/2) silty clay loam, dark gray (10YR 4/1) moist; weak, medium, subangular blocky structure; firm, very plastic; few fine roots; many fine, distinct, very dark grayish (10YR 3/1) Mn mottles throughout, few fine, faint, brown (7,5YR 4/3) iron mottles in upper part; neutral (pH 7); gradual wavy boundary.
56/62-90/93	B(ss)g	Dark gray (10YR 4/1) clay, moist; weak, medium, subangular blocky structure; firm, very plastic; few fine roots; many stress surfaces; many slickensides; many fine, distinct, very dark grayish (10YR 3/1) Mn mottles throughout; neutral (pH 7); clear wavy boundary.
90/93-110	BC	Very dark grayish brown (10YR 3/2) clay, moist; weak, medium, subangular blocky structure; firm, very plastic; few fine roots; neutral (pH 7); abrupt wavy boundary.
110-170+	2B(ss)g	Dark greyish (2.5Y 4/1) clay, moist; weak, fine, wedge structure parting to weak, fine, angular blocky, firm, very plastic; many Mn concretion; many slickensides; slightly alkaline (pH 7,5)

P7

Location name: Pleret, Imogiri District, Bantul Regency

Vegetation: paddy

Land Use: lowland paddy field

Parent Material: Quaternary Volcanic Deposit

Geological age: Quaternary

Topography: Nearly level

Soil Classification: Coarse-loamy over sandy-skeletal, mixed, isohyperthermic, Typic Epiaquepts


Morfologi Profil Tanah

Depth (cm)	Horizons	Soil Profile Description
0-17	Ap	Gray (10YR 4/1) sandy clay loam, moist; weak, fine, granular; friable, slightly plastic; many fine and medium roots; moderately acid (pH 6); clear smooth boundary.
17-40	Bg1	Very dark grayish Brown (10YR 3/2) sandy loam, gray (10YR 4/1), moist; weak, medium, granular; very friable, slightly plastic; 3-5% medium gravel; common fine roots; many fine, distinct, yellowish red (5YR 4/6) Fe mottles; slightly acid (pH 6,5); clear smooth boundary.
40-48	Bg2	Very dark grayish Brown (10YR 3/2) sandy loam, gray (10YR 4/1), moist; weak, medium, granular; very friable, slightly plastic; 3-5% medium gravel; few fine roots; many fine, distinct, black (10YR 2/1) Mn mottles; slightly acid (pH 6,5); clear smooth boundary.
48-68	C1	Gray (10YR 4/1) gravelly coarse sand, about 35% fine gravel and 15% coarse gravel; single grain; loose, nonplastic; few fine roots; slightly acid (pH 6,5); clear smooth boundary.
68-110	C2	Gray (10YR 4/1) very cobbly coarse sand, about 50% cobbles and 15% coarse gravel; single grain; loose, nonplastic; few fine roots; slightly acid (pH 6,5); abrupt smooth boundary.
110-121	2C3	Yellow (2,5Y 7/8) cobble deposit; clear smooth boundary.
121-170	3C4	Gray (10YR 4/1) coarse sand, moist; single grain; loose, nonplastic; slightly acid (pH 6,5)




### Lampiran 3. Korelasi Spearman

#### Tanah Halus



#### Tanah Kasar





UNIVERSITAS  
GADJAH MADA

**Faktor-Faktor Pedologis yang Memengaruhi Dinamika Silikon pada Tanah-Tanah di Cekungan Yogyakarta**

Padana Aperta Barus, Dr. Ir. Eko Hanudin, M.P. ; Dr.rer.nat. Ir. I Wayan Warmada, IPM.

Universitas Gadjah Mada, 2024 | Diunduh dari <http://etd.repository.ugm.ac.id/>