

DAFTAR PUSTAKA

- Alongi, D. M., Murdiyarso, D., Fourqurean, J. W., Kauffman, J. B., Hutahaean, A., Crooks, S., ... & Wagey, T. (2016). Indonesia's blue carbon: a globally significant and vulnerable sink for seagrass and *mangrove* carbon. *Wetlands ecology and management*, *24*, 3-13.
- Armstrong, R. A. (1993). Remote sensing of submerged vegetation canopies for biomass estimation. *International Journal of Remote Sensing*, *14*(3), 621–627. <https://doi.org/10.1080/01431169308904363>
- Bekkby, T., Rinde, E., Erikstad, L., Bakkestuen, V., Longva, O., Christensen, O., ... & Isachsen, P. E. (2008). Spatial probability modelling of eelgrass (*Zostera marina*) distribution on the west coast of Norway. *ICES Journal of Marine Science*, *65*(7), 1093-1101.
- Burdige, D. J. (2007). Preservation of organic matter in marine sediments: controls, mechanisms, and an imbalance in sediment organic carbon budgets. *Chemical reviews*, *107*(2), 467-485.
- Breiman, L. (2001). Random forests. *Machine learning*, *45*, 5-32.
- Carpenter, S., Byfield, V., Felgate, S. L., Price, D. M., Andrade, V., Cobb, E., Strong, J., Lichtschlag, A., Brittain, H., Barry, C., Fitch, A., Young, A., Sanders, R., & Evans, C. (2022). Using Unoccupied Aerial Vehicles (UAVs) to Map Seagrass Cover from Sentinel-2 Imagery. *Remote Sensing*, *14*(3), 477. <https://doi.org/10.3390/rs14030477>
- Chen, J., Sasaki, J., Guo, Z., & Endo, M. (2023). UAV-based seagrass wrack orthophotos classification for estimating blue carbon. *Estuarine, Coastal and Shelf Science*, *293*, 108476.
- Chmura, G. L., Anisfeld, S. C., Cahoon, D. R., & Lynch, J. C. (2003). Global carbon sequestration in tidal, saline wetland soils. *Global Biogeochemical Cycles*, *17*(4). <https://doi.org/10.1029/2002gb001917>
- Christie, H., Norderhaug, K. M., & Fredriksen, S. (2009). Macrophytes as habitat for fauna. *Marine ecology progress series*, *396*, 221-233.

- Costanza, R., D'arge, R., De Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg G', K., Naeem, S., O'Neill, R. V., Paruelo J', J., Raskin, R. G., Sutton, P., & Van Den Belt, M. (1998). The value of the world's ecosystem services and natural capitaP. In *Ecological Economics* (Vol. 25).
- Danoedoro, P. (2012). Pengantar penginderaan jauh digital. *Penerbit Andi, Yogyakarta*.
- Den Hartog, C. (1970). The sea-grasses of the world. *North-Holland, Amsterdam*.
- Duarte, C. M., & Cebrián, J. (1996). The fate of marine autotrophic production. *Limnology and Oceanography*, 41(8), 1758–1766. <https://doi.org/10.4319/lo.1996.41.8.1758>
- Duarte, C. M., Dennison, W. C., Orth, R. J. W., & Carruthers, T. J. B. (2008). The charisma of coastal ecosystems: Addressing the imbalance. In *Estuaries and Coasts* (Vol. 31, Issue 2, pp. 233–238). Springer New York. <https://doi.org/10.1007/s12237-008-9038-7>
- Duarte, C. M., Middelburg, J. J., & Caraco, N. (2005). Major role of marine vegetation on the oceanic carbon cycle. In *Biogeosciences* (Vol. 2). www.biogeosciences.net/bg/2/1/
- European Space Agency. (2015). *The Sentinel missions*. (Diakses pada 27 November 2023 dari https://www.esa.int/Applications/Observing_the_Earth/Copernicus/The_Sentinel_missions)
- Fonseca, M. S., & Bell, S. S. (1998). Influence of physical setting on seagrass landscapes near Beaufort, North Carolina, USA. *Marine Ecology Progress Series*, 171, 109-121.
- Fourqrean, J. W., Duarte, C. M., Kennedy, H., Marbà, N., Holmer, M., Mateo, M. A., ... & Serrano, O. (2012). Seagrass ecosystems as a globally significant carbon stock. *Nature geoscience*, 5(7), 505-509.
- Freund, Y., & Schapire, R. E. (1996, July). Experiments with a new boosting algorithm. In *icml* (Vol. 96, pp. 148-156).

- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote sensing of Environment*, 202, 18-27.
- Griffiths, P., Van Der Linden, S., Kuemmerle, T., & Hostert, P. (2013). Erratum: A pixel-based landsat compositing algorithm for large area land cover mapping (IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing). In *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* (Vol. 6, Issue 5, pp. 2088–2101). <https://doi.org/10.1109/JSTARS.2012.2228167>
- Gutman, G., Byrnes, R. A., Masek, J., Covington, S., Justice, C., Franks, S., & Headley, R. (2008). Towards monitoring land-cover and land-use changes at a global scale: The Global Land Survey 2005. *Photogrammetric Engineering and Remote Sensing*, 74(1), 6-10.
- Hernawan, U. E., Sjafrie, N. D., Supriyadi, I. H., Suyarso, I. M., & Anggraini, K. (2017). Status padang lamun Indonesia. *Pus. Penelit. Oseanografi (LIPI), Lemb. Ilmu Pengetah. Indones.*
- Hedley, J. D., Harborne, A. R., & Mumby, P. J. (2005). Simple and robust removal of sun glint for mapping shallow-water benthos. *International Journal of Remote Sensing*, 26(10), 2107–2112. <https://doi.org/10.1080/01431160500034086>
- Hidayat, W., Warpala, I. S., & Dewi, N. S. R. (2018). Komposisi jenis lamun (seagrass) dan karakteristik biofisik perairan di kawasan Pelabuhan Desa Celukanbawang Kecamatan Gerokgak Kabupaten Buleleng Bali. *Jurnal Pendidikan Biologi Undiksha*, 5(3), 133-145.
- Hossain, M. S., Bujang, J. S., Zakaria, M. H., & Hashim, M. (2015). The application of remote sensing to seagrass ecosystems: an overview and future research prospects. *International Journal of Remote Sensing*, 36(1), 61-114.
- Howard, J., Hoyt, S., Isensee, K., Telszewski, M., Pidgeon, E. (Eds.) (2014). Coastal Blue Carbon: Methods For Assessing Carbon Stocks And Emissions Factors In *Mangroves, Tidal Salt Marshes, And Seagrasses. Conservation International*,

Intergovernmental Oceanographic Commission Of UNESCO, International Union For Conservation Of Nature. Arlington, Virginia, USA.

- Jin, X. L., Diao, W. Y., Xiao, C. H., Wang, F. Y., Chen, B., Wang, K. R., & Li, S. K. (2013). Estimation of wheat agronomic parameters using new spectral indices. *PLoS One*, 8(8), e72736.
- Kay, S., Hedley, J. D., & Lavender, S. (2009). Sun glint correction of high and low spatial resolution images of aquatic scenes: A review of methods for visible and near-infrared wavelengths. In *Remote Sensing* (Vol. 1, Issue 4, pp. 697–730). <https://doi.org/10.3390/rs1040697>
- Kiswara, W., Kumoro, E. D., Kawaroe, M., & Rahadian, N. P. (2010). Transplanting *Enhalus Acoroides* (L.F) Royle With Different Length Rhizome On The Muddy Substrate And High Water Dynamic At Banten Bay, Indonesia. *Marine Research in Indonesia*, 35(2), 1–7. <https://doi.org/10.14203/mri.v35i2.472>
- Knudby, A., & Nordlund, L. (2011). Remote sensing of seagrasses in a patchy multi-species environment. *International Journal of Remote Sensing*, 32(8), 2227–2244. <https://doi.org/10.1080/01431161003692057>
- Koedsin, W., Intararuang, W., Ritchie, R. J., & Huete, A. (2016). An integrated field and remote sensing method for mapping seagrass species, cover, and biomass in southern Thailand. *Remote Sensing*, 8(4), 292.
- Lamb, J. B., Van De Water, J. A., Bourne, D. G., Altier, C., Hein, M. Y., Fiorenza, E. A., ... & Harvell, C. D. (2017). Seagrass ecosystems reduce exposure to bacterial pathogens of humans, fishes, and invertebrates. *Science*, 355(6326), 731-733.
- Li, B., Zhao, H., & Lv, Z. (2010, October). Parallel ISODATA clustering of remote sensing images based on MapReduce. In *2010 International Conference on Cyber-Enabled Distributed Computing and Knowledge Discovery* (pp. 380-383). IEEE.
- Lillesand, T., Kiefer, R. W., & Chipman, J. (2015). *Remote sensing and image interpretation*. John Wiley & Sons.

- Lima, M., Bergamo, T. F., Ward, R. D., & Joyce, C. B. (2023). A review of seagrass ecosystem services: providing nature-based solutions for a changing world. *Hydrobiologia*, 850(12), 2655-2670.
- Lyons, M., Phinn, S., & Roelfsema, C. (2011). Integrating Quickbird multi-spectral satellite and field data: Mapping bathymetry, seagrass cover, seagrass species and change in Moreton Bay, Australia in 2004 and 2007. *Remote Sensing*, 3(1), 42–64. <https://doi.org/10.3390/rs3010042>
- Memarsadeghi, N., Mount, D. M., Netanyahu, N. S., & Le Moigne, J. (2007). A Fast Implementation Of The Isodata Clustering Algorithm. *International Journal of Computational Geometry & Applications*, 17(01), 71–103. doi:10.1142/s0218195907002252
- Mashoreng, S., Isyrini, R., & Gosari, B. A. J. (2013). Estimasi Potensi Penyerapan Karbon Oleh Komunitas Lamun Di Pulau Barranglompo Makassar. *Lembaga Penelitian Dan Pengabdian Masyarakat, Universitas Hasanuddin*.
- Mateo, M. A., Romero, J., Pérez, M., Littler, M. M., & Littler, D. S. (1997). Dynamics of millenary organic deposits resulting from the growth of the Mediterranean seagrass *Posidonia oceanica*. *Estuarine, Coastal and Shelf Science*, 44(1), 103-110.
- McKenzie, L. J., Finkbeiner, M. A., & Kirkman, H. (2001). Methods for mapping seagrass distribution. *Global seagrass research methods*, 101-121.
- McKenzie, L. J., Yoshida, R. L., Mellors, J. E., & Coles, R. G. (2009). Seagrass-watch. In *Proceeding of a workshop for monitoring seagrass habitats in Indonesia. The Nature Conservancy, Coral Triangle Center, Sanur, Bali* (pp. 29-32).
- Munir, M., & Wicaksono, P. (2019, December). Support vector machine for seagrass percent cover mapping using PlanetScope image in Labuan Bajo, East Nusa Tenggara. In *Sixth International Symposium on LAPAN-IPB Satellite* (Vol. 11372, pp. 582-587). SPIE.
- Nahirnick, N. K., Reshitnyk, L., Campbell, M., Hessian-Lewis, M., Costa, M., Yakimishyn, J., & Lee, L. (2019). Mapping with confidence; delineating seagrass

- habitats using Unoccupied Aerial Systems (UAS). *Remote Sensing in Ecology and Conservation*, 5(2), 121-135.
- Pendleton, L., Donato, D. C., Murray, B. C., Crooks, S., Jenkins, W. A., Sifleet, S., ... & Baldera, A. (2012). Estimating global “blue carbon” emissions from conversion and degradation of vegetated coastal ecosystems.
- Philips, C. R., & Menez, E. G. (1988). *Seagrass. Smith Sonian Institutions Press, Washington DC.*
- Phinn, S., Roelfsema, C., Dekker, A., Brando, V., & Anstee, J. (2008). Mapping seagrass species, cover and biomass in shallow waters: An assessment of satellite multi-spectral and airborne hyper-spectral imaging systems in Moreton Bay (Australia). *Remote Sensing of Environment*, 112(8), 3413–3425. <https://doi.org/10.1016/j.rse.2007.09.017>
- Rahayu, Y. P., Solihuddin, T., Kusumaningtyas, M. A., Afi Ati, R. N., Salim, H. L., Rixen, T., & Hutahaeon, A. A. (2019). The sources of organic matter in seagrass sediments and their contribution to carbon stocks in the Spermonde Islands, Indonesia. *Aquatic Geochemistry*, 25(3), 161-178.
- Rahmi, K. I. N., & Febrianti, N. (2020). Pemanfaatan data sentinel-2 untuk analisis indeks area terbakar (burned area). *Jurnal Penginderaan Jauh Indonesia*, 2(1), 1-6.
- Rais, M., Inaku, D. F., Moka, W. J., Mashoreng, S., Satari, D. Y., & Rukminasari, N. (2023). Estimasi Stok Karbon Padang Lamun menggunakan Citra Spot-7 di Perairan Pulau Kodingarenglompo, Sangkarrang, Kota Makassar. *Jurnal Kelautan Tropis*, 26(2), 387-398.
- Ralph, P. J., Crosswell, J. R., Cannard, T., & Steven, A. D. (2018). Estimating seagrass blue carbon and policy implications: The Australian perspective. *Seagrasses of Australia: structure, ecology and conservation*, 743-758.
- Roelfsema, C. M., Lyons, M., Kovacs, E. M., Maxwell, P., Saunders, M. I., Samper-Villarreal, J., & Phinn, S. R. (2014). Multi-temporal mapping of seagrass cover, species and biomass: A semi-automated object based image analysis approach.

- Remote Sensing of Environment*, 150, 172–187.
<https://doi.org/10.1016/j.rse.2014.05.001>
- Roelfsema, C. M., Phinn, S. R., Udy, N., & Maxwell, P. (2009). An integrated field and remote sensing approach for mapping seagrass cover, moreton bay, Australia. *Journal of Spatial Science*, 54(1), 45–62.
<https://doi.org/10.1080/14498596.2009.9635166>
- Rustam, A., Sudirman, N., Ati, R. N. A., Salim, H. L., & Rahayu, Y. P. (2017). Seagrass ecosystem carbon stock in the small islands: case study in Spermonde island, South Sulawesi, Indonesia. *Jurnal Segara*, 13(2).
- Setyanto, R. A., Widianingsih, W., & Setyati, W. A. (2023). Simpanan Karbon Lamun di Pulau Sintok dan Pulau Menjangan Besar, Karimunjawa. *Journal of Marine Research*, 12(1), 110-115.
- Scornet, E. (2017). Tuning parameters in random forests. *ESAIM. Proceedings and Surveys*, 60.
- Short, F. T. (2003). *World atlas of seagrasses*. Univ of California Press.
- Sifleet, S., Pendleton, L., & Murray, B. C. (2011). *State of the Science on Coastal Blue Carbon A Summary for Policy Makers*.
- Simpson, J., Bruce, E., Davies, K. P., & Barber, P. (2022). A blueprint for the estimation of seagrass carbon stock using remote sensing-enabled proxies. *Remote Sensing*, 14(15), 3572.
- Siregar, A. Z. (2021). Implementasi Metode Regresi Linier Berganda Dalam Estimasi Tingkat Pendaftaran Mahasiswa Baru. *Kesatria: Jurnal Penerapan Sistem Informasi (Komputer dan Manajemen)*, 2(3), 133-137.
- Skilbeck, C. G., Trevathan-Tackett, S., Apichanangkool, P., & Macreadie, P. I. (2017). Sediment sampling in estuaries: site selection and sampling techniques. *Applications of paleoenvironmental techniques in estuarine studies*, 89-120.
- Sorzano, C. O. S., Vargas, J., & Montano, A. P. (2014). A survey of dimensionality reduction techniques. *arXiv preprint arXiv:1403.2877*.

- Stekoll, M. S., Deysher, L. E., & Hess, M. (2006). A remote sensing approach to estimating harvestable kelp biomass. *Journal of Applied Phycology*, 18(3–5), 323–334. <https://doi.org/10.1007/s10811-006-9029-7>
- Supriadi, S., Kaswadji, R. F., Bengen, D. G., & Hutomo, M. (2014). Carbon stock of seagrass community in Barranglompo Island, Makassar (stok karbon pada komunitas lamun di Pulau Barranglompo, Makassar). *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, 19(1), 1-10.
- Talavera, L., Vila-Concejo, A., Webster, J. M., Smith, C., Duce, S., Fellowes, T. E., ... & Hacker, J. (2021). Morphodynamic controls for growth and evolution of a rubble coral island. *Remote Sensing*, 13(8), 1582.
- Tou, J.T., & Gonzales, R.C. (1974). *Pattern Recognition Principles*.
- Unsworth, R. K., van Keulen, M., & Coles, R. G. (2014). Seagrass meadows in a globally changing environment. *Marine Pollution Bulletin*, 83(2), 383-386.
- Wahyudi, A. A. J., Rahmawati, S., Irawan, A., Hadiyanto, H., Prayudha, B., Hafizt, M., ... & Kiswara, W. (2020). Assessing carbon stock and sequestration of the tropical seagrass meadows in Indonesia. *Ocean Science Journal*, 55, 85-97.
- Wicaksono, P. (2015). *Remote Sensing Model Development For Seagrass and Mangroves Carbon Stock Mapping*. Yogyakarta: Fakultas Geografi Universitas Gadjah Mada
- Wicaksono, P. (2016). Improving the accuracy of Multispectral-based benthic habitats mapping using image rotations: the application of Principle Component Analysis and Independent Component Analysis. *European Journal of Remote Sensing*, 49(1), 433-463.
- Wicaksono, P., Wulandari, S. A., Lazuardi, W., & Munir, M. (2021). Sentinel-2 images deliver possibilities for accurate and consistent multi-temporal benthic habitat maps in optically shallow water. *Remote Sensing Applications: Society and Environment*, 23, 100572.
- Wicaksono, P., & Lazuardi, W. (2021). Assessment of PlanetScope images for benthic habitat and seagrass species mapping in a complex optically shallow water

- environment. In *Fine Resolution Remote Sensing of Species in Terrestrial and Coastal Ecosystems* (pp. 143-169). Routledge.
- Wicaksono, P., Aryaguna, P. A., & Lazuardi, W. (2019). Benthic habitat mapping model and cross validation using machine-learning classification algorithms. *Remote Sensing*, *11*(11), 1279.
- Wicaksono, P., Lazuardi, W., & Munir, M. (2019). Integrating Image At Different Spatial Resolutions And Field Data For Seagrass Percent Cover Mapping. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives*, *42*(4/W19), 487–492. <https://doi.org/10.5194/isprs-archives-XLII-4-W19-487-2019>
- Wicaksono, P., Fauzan, M. A., & Asta, S. G. W. (2020). Assessment of Sentinel-2A multispectral image for benthic habitat composition mapping. *IET Image Processing*, *14*(2), 279-288.
- Wicaksono, P., Maishella, A., & Hafizt, M. (2022). Multitemporal seagrass carbon assimilation and aboveground carbon stock mapping using Sentinel-2 in Labuan Bajo 2019–2020. *Remote Sensing Applications: Society and Environment*, *27*, 100803.
- Wicaksono, P., Hafizt, M., Harahap, S. D., & Nandika, M. R. (2024). Integrating Sentinel-2 and PlanetScope Image with Drone-based Seagrass Data for Seagrass Percent Cover Mapping. In *IOP Conference Series: Earth and Environmental Science* (Vol. 1291, No. 1, p. 012012). IOP Publishing.
- Wright, J. P., & Jones, C. G. (2006). The Concept Of Organism As Ecosystem Engineers Ten Years On: Progress, Limitations, And Challenges. *BioScience*, *56*(3), 203–209. www.biosciencemag.org
- Yang, B., Hawthorne, T. L., Hessing-Lewis, M., Duffy, E. J., Reshitnyk, L. Y., Feinman, M., & Searson, H. (2020). Developing an introductory uav/drone mapping training program for seagrass monitoring and research. *Drones*, *4*(4), 1–18. <https://doi.org/10.3390/drones4040070>

Zhou, X., Zhu, X., Dong, Z., & Guo, W. (2016). Estimation of biomass in wheat using random forest regression algorithm and remote sensing data. *The Crop Journal*, 4(3), 212-219.