

DAFTAR PUSTAKA

- Amaral, M. K., Péllico Netto, S., Lingnau, C., & Figueiredo Filho, A. (2015). Evaluation of the morisita index for determination of the spatial distribution of species in a fragment of araucaria forest. *Applied Ecology and Environmental Research*, 13(2). https://doi.org/10.15666/aeer/1302_361372
- Amos, J. N., Harrison, K. A., Radford, J. Q., White, M., Newell, G., Nally, R. Mac, Sunnucks, P., & Pavlova, A. (2014). Species- and sex-specific connectivity effects of habitat fragmentation in a suite of woodland birds. *Ecology*, 95(6). <https://doi.org/10.1890/13-1328.1>
- Bednar-Friedl, B., Behrens, D. A., & Getzner, M. (2012). Optimal Dynamic Control of Visitors and Endangered Species in a National Park. *Environmental and Resource Economics*, 52(1). <https://doi.org/10.1007/s10640-011-9515-5>
- BirdLifeInternational. (2018). THE IUCN RED LIST OF THREATENED SPECIES™. *The IUCN Red List of Threatened Species 2018: E.T22679440A131749282*. <https://doi.org/10.2305/IUCN.UK.2018>
- Bonesi, L., Rushton, S., & Macdonald, D. (2002). The combined effect of environmental factors and neighbouring populations on the distribution and abundance of *Arvicola terrestris*. An approach using rule-based models. *Oikos*, 99(2). <https://doi.org/10.1034/j.1600-0706.2002.990202.x>
- Boyce, M. S., & Merrill, E. H. (1993). Wildlife-Habitat Relationships: Concepts and Applications . Michael L. Morrison , Bruce G. Marcot , R. William Mannan . *The Quarterly Review of Biology*, 68(2). <https://doi.org/10.1086/418137>
- Brickle, N. W. (2002). Habitat use, predicted distribution and conservation of green peafowl (*Pavo muticus*) in Dak Lak province, Vietnam. *Biological Conservation*, 105(2). [https://doi.org/10.1016/S0006-3207\(01\)00182-3](https://doi.org/10.1016/S0006-3207(01)00182-3)
- BTNB. (2014). *Rencana Pengelolaan Taman Nasional Baluran Tahun 2014-2023*.

19(5), 1–114.

Coulloudon, B., Eshelman, K., Gianola, J., Habich, N., Hughes, L., Johnson, C., Pellant, M., Podborny, P., Rasmussen, A., Robles, B., Shaver, P., Spehar, J., & Willoughby, J. (1996). Sampling vegetation attributes: interagency technical reference. *BLM Technical Reference*.

Davis, N. E., Di Stefano, J., Coulson, G., Whelan, J., & Wright, J. (2016). Vegetation management influences habitat use by mammalian herbivores in shrub-encroached grassy woodland. *Wildlife Research*, 43(5). <https://doi.org/10.1071/WR15218>

Dong, F., Kuo, H. C., Chen, G. L., Wu, F., Shan, P. F., Wang, J., Chen, D., Lei, F. M., Hung, C. M., Liu, Y., & Yang, X. J. (2021). Population genomic, climatic and anthropogenic evidence suggest the role of human forces in endangerment of green peafowl (*Pavo muticus*). *Proceedings of the Royal Society B: Biological Sciences*, 288(1948). <https://doi.org/10.1098/rspb.2021.0073>

Draper, N. R., & Smith, H. (2014). Applied regression analysis. In *Applied Regression Analysis*. <https://doi.org/10.1002/9781118625590>

Durazzo, A., Lucarini, M., Zaccardelli, M., & Santini, A. (2020). Forest, foods, and nutrition. In *Forests* (Vol. 11, Issue 11). <https://doi.org/10.3390/f11111182>

Eddyono, F., Darusman, D., Sumarwan, U., & Sunarminto, T. (2021). Tourism competitiveness on the utilization zone of National Park in Indonesia. *Jurnal Penelitian Kehutanan Wallacea*, 10(2). <https://doi.org/10.18330/jwallacea.2021.vol10iss2pp145-164>

FABRICANTE, J. R., & OLIVEIRA, C. R. (2013). Estrutura populacional de *Melocactus ernestii* Vaupel subsp. *ernestii* (Cactaceae) J. *SCIENTIA PLENA*, 9(6).

Flores, B. M., & Holmgren, M. (2021). White-Sand Savannas Expand at the Core of the Amazon After Forest Wildfires. *Ecosystems*, 24(7). <https://doi.org/10.1007/s10021-021-00607-x>

Franklin, J., & Miller, J. A. (2010). Mapping species distributions: Spatial inference

- and prediction. In *Mapping Species Distributions: Spatial Inference and Prediction*. <https://doi.org/10.1017/CBO9780511810602>
- Gamon, J. A., Somers, B., Malenovsky, Z., Middleton, E. M., Rascher, U., & Schaepman, M. E. (2019). Assessing Vegetation Function with Imaging Spectroscopy. In *Surveys in Geophysics* (Vol. 40, Issue 3). <https://doi.org/10.1007/s10712-019-09511-5>
- Gaston, K. J., & Fuller, R. A. (2007). Biodiversity and extinction: Losing the common and the widespread. *Progress in Physical Geography*, *31*(2), 213–225. <https://doi.org/10.1177/0309133307076488>
- Grinnell, J. (1917). The Niche-Relationships of the California Thrasher. *The Auk*, *34*(4). <https://doi.org/10.2307/4072271>
- Gu, Y., Hunt, E., Wardlow, B., Basara, J. B., Brown, J. F., & Verdin, J. P. (2008). Evaluation of MODIS NDVI and NDWI for vegetation drought monitoring using Oklahoma Mesonet soil moisture data. *Geophysical Research Letters*, *35*(22). <https://doi.org/10.1029/2008GL035772>
- Guichard, F. (2017). Regular patterns link individual behavior to population persistence. In *Proceedings of the National Academy of Sciences of the United States of America* (Vol. 114, Issue 30). <https://doi.org/10.1073/pnas.1709063114>
- Guisan, A., & Zimmermann, N. E. (2000). Predictive habitat distribution models in ecology. *Ecological Modelling*, *135*(2–3). [https://doi.org/10.1016/S0304-3800\(00\)00354-9](https://doi.org/10.1016/S0304-3800(00)00354-9)
- Habib, M. (2021). Evaluation of DEM interpolation techniques for characterizing terrain roughness. *Catena*, *198*. <https://doi.org/10.1016/j.catena.2020.105072>
- Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E., Sexton, J. O., Austin, M. P., Collins, C. D., Cook, W. M., Damschen, E. I., Ewers, R. M., Foster, B. L., Jenkins, C. N., King, A. J., Laurance, W. F., Levey, D. J., Margules, C. R., ... Townshend, J. R. (2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, *1*(2). <https://doi.org/10.1126/sciadv.1500052>

- Harjadi, B. (2017). Detection of water abundance in Baluran National Park with landsat satellite imagery analysis. *Jurnal Penelitian Kehutanan Wallacea*, 6(1), 53. <https://doi.org/10.18330/jwallacea.2017.vol6iss1pp53-60>
- Heikkinen, R. K., Luoto, M., Virkkala, R., Pearson, R. G., & Körber, J. H. (2007). Biotic interactions improve prediction of boreal bird distributions at macro-scales. *Global Ecology and Biogeography*, 16(6). <https://doi.org/10.1111/j.1466-8238.2007.00345.x>
- Henderson, P. A. (2021). Southwood's ecological methods. In *Southwood's Ecological Methods*. <https://doi.org/10.1093/oso/9780198862277.001.0001>
- Hernowo, J. B. (1999). *Habitat Local Distribution Of Javan Green Peafowl (*Pavo muticus* Linnaeus 1758) In Baluran National Park, East Java*.
- Hernowo, J. B. (2017). Determinant factors of the javan green peafowl (*Pavo muticus muticus* Linnaeus, 1758) habitat in Baluran and Alas Purwo National Parks, East Java, Indonesia. *Tropical Drylands*, 1(1). <https://doi.org/10.13057/tropdrylands/t010108>
- Hernowo, J. B., & Alikodra, H. S. (2018). The grouping system and local distribution pattern analysis of Javan green peafowl (*Pavo muticus muticus*, linnaeus 1758) population in Baluran and Alas Purwo National Parks, East Java, Indonesia. *Biodiversitas*, 19(5), 1690–1695. <https://doi.org/10.13057/biodiv/d190514>
- Hernowo, J. B., Alikodra, H. S., Mardiasuti, A., & Kusmana, C. (2011). Population analysis of the javan green peafowl (*Pavo muticus muticus* Linnaeus 1758) in Baluran and Alas Purwo National Parks, East Java. *Biodiversitas Journal of Biological Diversity*, 12(2). <https://doi.org/10.13057/biodiv/d120207>
- Hernowo, J. B., Kusmana, C., Alikodra, H. S., & Mardiasuti, A. (2018). Analysis of the Javan Green Peafowl (*Pavo muticus muticus* linnaeus 1758) habitat in Baluran and Alas Purwo National Park, East Java. *HAYATI Journal of Biosciences*, 25(3), 101–114. <https://doi.org/10.4308/hjb.25.3.101>
- Hernowo, J. B., Mardiasuti, A. N. I., Alikodra, H. S., & Kusmana, C. E. C. E. P. (2011). Behavior Ecology of the Javan Green Peafowl (*Pavo muticus muticus*

- Linnaeus 1758) in Baluran and Alas Purwo National Park, East Java. *HAYATI Journal of Biosciences*, 18(4), 164–176. <https://doi.org/10.4308/hjb.18.4.164>
- Hijmans, R. J. (2020). raster: Geographic analysis and modeling with raster data. : *R Package Version 3.1-5*, 3.1(5).
- Hijmans, R. J., Bivand, R., Dyba, K., Pebesma, E., & Sumner, M. D. (2024). Spatial Data Analysis. R Package ‘terra’. *October*, 3(500), 1–6. <https://rspatial.github.io/terra/><https://rspatial.org/>, [https://rspatial.github.io/terra/ BugReports](https://rspatial.github.io/terra/BugReports)
- Hijmans, R. J., & Elith, J. (2013). Species distribution modeling with R Introduction. *October*.
- Hudjimartsu, S., Prasetyo, L., Setiawan, Y., & Suyamto, D. (2017). Illumination Modelling for Topographic Correction of Landsat 8 and Sentinel-2A Imageries. *Proceedings - UKSim-AMSS 11th European Modelling Symposium on Computer Modelling and Simulation, EMS 2017*. <https://doi.org/10.1109/EMS.2017.27>
- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., & Ferreira, L. G. (2002). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment*, 83(1–2). [https://doi.org/10.1016/S0034-4257\(02\)00096-2](https://doi.org/10.1016/S0034-4257(02)00096-2)
- Hunter JR, M. L., & Gibbs, J. (2007). *Fundamental of Conservation Biology* (third). Blackwell Publishing.
- Hutchinson, G. E. (1957). Concluding remarks cold spring harbor symposia on quantitative biology. *GS SEARCH*, 22.
- Ji, L., Zhang, L., & Wylie, B. (2009). Analysis of dynamic thresholds for the normalized difference water index. *Photogrammetric Engineering and Remote Sensing*, 75(11). <https://doi.org/10.14358/PERS.75.11.1307>
- Jose V, S., & Nameer, P. O. (2020). The expanding distribution of the Indian Peafowl (*Pavo cristatus*) as an indicator of changing climate in Kerala, southern India: A modelling study using MaxEnt. *Ecological Indicators*, 110. <https://doi.org/10.1016/j.ecolind.2019.105930>

- Khaemba, W. M. (2001). Spatial point pattern analysis of aerial survey data to assess clustering in wildlife distributions. *ITC Journal*, 3(2). [https://doi.org/10.1016/S0303-2434\(01\)85005-4](https://doi.org/10.1016/S0303-2434(01)85005-4)
- Knuff, A. K., Staab, M., Frey, J., Dormann, C. F., Asbeck, T., & Klein, A. M. (2020). Insect abundance in managed forests benefits from multi-layered vegetation. *Basic and Applied Ecology*, 48. <https://doi.org/10.1016/j.baae.2020.09.002>
- Liu, Y., Han, L., Xie, Y., Wen, Y., & Zhang, R. (2007). The status and habitat use of green peafowl *Pavo muticus* in Shuangbai Konglonghe Nature Reserve, China. In *International Journal of Galliformes Conservation* (Vol. 1).
- Ma, Q. Y. (2009). Analysis of the negative binomial distribution and test of population pattern. *Beijing Linye Daxue Xuebao/Journal of Beijing Forestry University*, 31(3).
- Macarthur, R. H., & Wilson, E. O. (2021). The Theory of Island Biogeography. In *A Century in Books: Princeton University Press 1905–2005*. <https://doi.org/10.2307/3799353>
- Manel, S., Ceri Williams, H., & Ormerod, S. J. (2001). Evaluating presence-absence models in ecology: The need to account for prevalence. *Journal of Applied Ecology*, 38(5). <https://doi.org/10.1046/j.1365-2664.2001.00647.x>
- Matsushita, B., Yang, W., Chen, J., Onda, Y., & Qiu, G. (2007). Sensitivity of the Enhanced Vegetation Index (EVI) and Normalized Difference Vegetation Index (NDVI) to topographic effects: A case study in high-density cypress forest. *Sensors*, 7(11). <https://doi.org/10.3390/s7112636>
- McComb, B. C. (2007). *Wildlife Habitat Management: Concept and Applications in Forestry*. CRC Press.
- McFeeters, S. K. (1996). The use of the Normalized Difference Water Index (NDWI) in the delineation of open water features. *International Journal of Remote Sensing*, 17(7). <https://doi.org/10.1080/01431169608948714>
- McGowan, P. J. K., & Kirwan, G. M. (2020). Green Peafowl (*Pavo muticus*). In *Birds of the World*. <https://doi.org/10.2173/bow.grepea1.01>

- Mkenda, P. A., Ndakidemi, P. A., Mbega, E., Stevenson, P. C., Arnold, S. E. J., Gurr, G. M., & Belmain, S. R. (2019). Multiple ecosystem services from field margin vegetation for ecological sustainability in agriculture: Scientific evidence and knowledge gaps. *PeerJ*, 2019(11). <https://doi.org/10.7717/peerj.8091>
- Mo, D., Yan, E., Lin, H., Sun, H., Li, J., & Zhang, G. (2012). Development and validation of 2-band EVI with MODIS data in Southeast China. *Proceedings of 2012 International Conference on Measurement, Information and Control, MIC 2012, 1*. <https://doi.org/10.1109/MIC.2012.6273306>
- Morisita, M. (1962). $I\sigma$ -Index, a measure of dispersion of individuals. *Researches on Population Ecology*, 4(1). <https://doi.org/10.1007/BF02533903>
- Morrison, M. L., Marcot, B. G., & Mannan, R. W. (2002). Wildlife-habitat relationships: concepts and applications. In *NCASI Technical Bulletin* (Vol. 2, Issue 781). <https://doi.org/10.2307/4002558>
- Moudrý, V., Cord, A. F., Gábor, L., Laurin, G. V., Barták, V., Gdulová, K., Malavasi, M., Rocchini, D., Stereńczak, K., Prošek, J., Klápště, P., & Wild, J. (2023). Vegetation structure derived from airborne laser scanning to assess species distribution and habitat suitability: The way forward. *Diversity and Distributions*, 29(1). <https://doi.org/10.1111/ddi.13644>
- Moustakas, A., Wiegand, K., Meyer, K. M., Ward, D., & Sankaran, M. (2012). perspective: Learning new tricks from old trees: revisiting the savanna question. *Frontiers of Biogeography*, 2(2). <https://doi.org/10.21425/f5fbg12335>
- Noon, B. R. (1981). Techniques for sampling avian habitats. In *The use of multivariate statistics in studies of wildlife habitat*.
- Norbury, G., & van Overmeire, W. (2019). Low structural complexity of nonnative grassland habitat exposes prey to higher predation. *Ecological Applications*, 29(2). <https://doi.org/10.1002/eap.1830>
- Nuttall, M. N., Griffin, O., Fewster, R. M., McGowan, P. J. K., Abernethy, K., O'Kelly, H., Nut, M., Sot, V., & Bunnefeld, N. (2022). Long-term monitoring of wildlife populations for protected area management in Southeast Asia.

- Conservation Science and Practice*, 4(2), 1–17. <https://doi.org/10.1111/csp2.614>
- Nuttall, M., Nut, M., Ung, V., & O’Kelly, H. (2017). Abundance estimates for the endangered Green Peafowl *Pavo muticus* in Cambodia: Identification of a globally important site for conservation. *Bird Conservation International*, 27(1). <https://doi.org/10.1017/S0959270916000083>
- Ober, H. K., & Hayes, J. P. (2008). Influence of forest riparian vegetation on abundance and biomass of nocturnal flying insects. *Forest Ecology and Management*, 256(5). <https://doi.org/10.1016/j.foreco.2008.06.010>
- Osborne, C. P., Charles-Dominique, T., Stevens, N., Bond, W. J., Midgley, G., & Lehmann, C. E. R. (2018). Human impacts in African savannas are mediated by plant functional traits. In *New Phytologist* (Vol. 220, Issue 1). <https://doi.org/10.1111/nph.15236>
- Pearce, J. L., & Boyce, M. S. (2006). Modelling distribution and abundance with presence-only data. In *Journal of Applied Ecology* (Vol. 43, Issue 3). <https://doi.org/10.1111/j.1365-2664.2005.01112.x>
- Pebesma, E., & Bivand, R. (2023). Spatial Data Science: With Applications in R. In *Spatial Data Science: With Applications in R*. <https://doi.org/10.1201/9780429459016>
- Pudyatmoko, S. (2019). Habitat dan Interaksi Spatio-Temporal Merak Hijau dengan Sapi dan Herbivora Besar di Taman Nasional Baluran. *Jurnal Ilmu Kehutanan*, 13(1). <https://doi.org/10.22146/jik.46142>
- Redlich, S., Martin, E. A., Wende, B., & Steffan-Dewenter, I. (2018). Landscape heterogeneity rather than crop diversity mediates bird diversity in agricultural landscapes. *PLoS ONE*, 13(8). <https://doi.org/10.1371/journal.pone.0200438>
- Rocha, A. V., & Shaver, G. R. (2009). Advantages of a two band EVI calculated from solar and photosynthetically active radiation fluxes. *Agricultural and Forest Meteorology*, 149(9). <https://doi.org/10.1016/j.agrformet.2009.03.016>
- Saridnirun, G., Sukumal, N., Grainger, M. J., & Savini, T. (2021). Living with human encroachment: Status and distribution of Green Peafowl in northern stronghold

- of Thailand. *Global Ecology and Conservation*, 28.
<https://doi.org/10.1016/j.gecco.2021.e01674>
- Shwe, N. M., Sukumal, N., Oo, K. M., Dowell, S., Browne, S., & Savini, T. (2021). Importance of isolated forest fragments and low intensity agriculture for the long-term conservation of the green peafowl *Pavo muticus*. *ORYX*, 55(2).
<https://doi.org/10.1017/S0030605319000267>
- Siebert, F., & Dreber, N. (2019). Forb ecology research in dry African savannas: Knowledge, gaps, and future perspectives. In *Ecology and Evolution* (Vol. 9, Issue 13). <https://doi.org/10.1002/ece3.5307>
- Silva, C. P., García, C. E., Estay, S. A., Barbosa, O., & Chapman, M. G. (2015). Bird richness and abundance in response to urban form in a Latin American City: Valdivia, Chile as a Case Study. *PLoS ONE*, 10(9).
<https://doi.org/10.1371/journal.pone.0138120>
- Srinivasan, U., Raza, R. H., & Quader, S. (2010). The nuclear question: Rethinking species importance in multi-species animal groups. *Journal of Animal Ecology*, 79(5). <https://doi.org/10.1111/j.1365-2656.2010.01707.x>
- Sukumal, N., Dowell, S. D., & Savini, T. (2017). Micro-habitat selection and population recovery of the Endangered Green Peafowl *Pavo muticus* in western Thailand: Implications for conservation guidance. *Bird Conservation International*, 27(3), 414–430. <https://doi.org/10.1017/S095927091600037X>
- Sukumal, N., Grainger, M. J., & Savini, T. (2020). Lower levels of human disturbance correspond with longer-term persistence of Endangered Green Peafowl *Pavo muticus* populations. *Bird Conservation International*, 30(2), 210–219. <https://doi.org/10.1017/S0959270919000443>
- Sukumal, N., McGowan, P. J. K., & Savini, T. (2015). Change in status of green peafowl *Pavo muticus* (Family Phasianidae) in Southcentral Vietnam: A comparison over 15years. *Global Ecology and Conservation*, 3, 11–19.
<https://doi.org/10.1016/j.gecco.2014.10.007>
- Takandjandji, M., & Sawitri, R. (2011).

POPULASI_BURUNG_MERAK_HIJAU_Pavo_muticus_Linnaeus_. *Jurnal Penelitian Ilmu Hutan Dan Konservasi Alam*, 8, 13–24.

Teng, J., Xia, S., Liu, Y., Yu, X., Duan, H., Xiao, H., & Zhao, C. (2021). Assessing habitat suitability for wintering geese by using Normalized Difference Water Index (NDWI) in a large floodplain wetland, China. *Ecological Indicators*, 122. <https://doi.org/10.1016/j.ecolind.2020.107260>

Tryjanowski, P., Jankowiak, Czechowski, P., Dulisz, B., Golawski, A., Grzywaczewski, G., Indykiewicz, P., Kwieciński, Z., Mitrus, C., Nowakowski, J. J., Polakowski, M., Siekiera, J., Sparks, T. H., & Wuczyński, A. (2022). Summer water sources for temperate birds: use, importance, and threats. *European Zoological Journal*, 89(1), 913–926. <https://doi.org/10.1080/24750263.2022.2101699>

Turchin, P., & Batzli, G. O. (2001). Availability of food and the population dynamics of arvicoline rodents. *Ecology*, 82(6). [https://doi.org/10.1890/0012-9658\(2001\)082\[1521:AOFATP\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[1521:AOFATP]2.0.CO;2)

U.S.D.A. (2010). Sampling vegetation. *Teaching*.

Umar, H. (2013). Metode Penelitian Untuk Skripsi Dan Tesis. In *Metode penelitian untuk skripsi dan tesi* (Vol. 1, Issue August).

Van Balen, S., Prawiradilaga, D. M., & Indrawan, M. (1995). *THE DISTRIBUTION AND STATUS OF GREEN PEA FOWL Pavo muticus IN JAVA*.

Varman, K. S., & Sukumar, R. (1995). The line transect method for estimating densities of large mammals in a tropical deciduous forest: An evaluation of models and field experiments. *Journal of Biosciences*, 20(2). <https://doi.org/10.1007/BF02703274>

Vasconcelos, R. P., Batista, M. I., & Henriques, S. (2017). Current limitations of global conservation to protect higher vulnerability and lower resilience fish species. *Scientific Reports*, 7(1). <https://doi.org/10.1038/s41598-017-06633-x>

Venables, W. N., & Ripley, B. D. (2002). Modern Applied Statistics with S Fourth edition by. In *World* (Vol. 53, Issue March).

- Wiegand, H. (1971). Pielou, E. C. An introduction to mathematical ecology. Wiley Interscience. John Wiley & Sons, New York 1969. VIII + 286 S., 32 Abb., Preis 140 s. *Biometrische Zeitschrift*, 13(3).
<https://doi.org/10.1002/bimj.19710130308>
- Win, L., Sukumal, N., Shwe, N. M., & Savini, T. (2023). Teak plantations are complementary habitat for the Green Peafowl movement within the fragmented Bago Yoma range, south-central Myanmar. *Ornithological Applications*, 125(1).
<https://doi.org/10.1093/ornithapp/duac051>
- Yorzinski, J. L. (2014). Peafowl antipredator calls encode information about signalers. *The Journal of the Acoustical Society of America*, 135(2).
<https://doi.org/10.1121/1.4861340>
- Zha, Y., Gao, J., & Ni, S. (2003). Use of normalized difference built-up index in automatically mapping urban areas from TM imagery. *International Journal of Remote Sensing*, 24(3). <https://doi.org/10.1080/01431160304987>