

DAFTAR PUSTAKA

- Arora, J., Mendelson, A.A. and Fox-Robichaud, A. (2023) 'Sepsis: Network Pathophysiology and implications for Early Diagnosis', *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, (March), pp. 613–624. Available at: <https://doi.org/10.1152/ajpregu.00003.2023>.
- Bardou, M., Quenot, J.P. and Barkun, A. (2015) 'Stress-related mucosal disease in the critically ill patient', *Nature Reviews Gastroenterology and Hepatology*. Nature Publishing Group, pp. 98–107. Available at: <https://doi.org/10.1038/nrgastro.2014.235>.
- Berger, M.M. *et al.* (2005) 'Enteral nutrition in critically ill patients with severe hemodynamic failure after cardiopulmonary bypass', *Clinical Nutrition*, 24(1), pp. 124–132.
- Bernardi, M. *et al.* (2015) 'Mechanisms of decompensation and organ failure in cirrhosis: from peripheral arterial vasodilation to systemic inflammation hypothesis', *Journal of hepatology*, 63(5), pp. 1272–1284.
- Brisard, L. *et al.* (2014) 'Impact of early enteral versus parenteral nutrition on mortality in patients requiring mechanical ventilation and catecholamines: study protocol for a randomized controlled trial (NUTRIREA-2)', *Trials*, 15, pp. 1–13.
- Butterworth, J.F., Mackey, D. and Wasnick, J.D. (2013) 'The Practice of Anesthesiology', 7(3).
- Chapman, M.J. *et al.* (2009) 'Glucose absorption and gastric emptying in critical illness', *Critical Care*, 13(4). Available at: <https://doi.org/10.1186/cc8021>.
- Chee, F. and Fernando, T. (2007) *Closed-loop control of blood glucose*. Springer.
- Van Cromphaut, S., Vanhorebeek, I. and Berghe, G. (2008) 'Glucose Metabolism and Insulin Resistance in Sepsis', *Current Pharmaceutical Design*, 14(19), pp. 1887–1899. Available at: <https://doi.org/10.2174/138161208784980563>.
- Dahlan, M.S. (2011) *Statistik untuk kedokteran dan kesehatan*. Penerbit Salemba.
- Deane, A.M. *et al.* (2011) 'Glucose absorption and small intestinal transit in critical illness', *Critical care medicine*, 39(6), pp. 1282–1288.
- Deane, A.M. *et al.* (2014) 'The effects of critical illness on intestinal glucose sensing, transporters, and absorption', *Critical Care Medicine*, 42(1), pp. 57–65. Available at: <https://doi.org/10.1097/CCM.0b013e318298a8af>.
- DeMers, D. and Wachs, D. (2021) 'Physiology, mean arterial pressure', in *StatPearls [Internet]*. StatPearls Publishing.
- Deng, J. *et al.* (2023) 'Comprehensive Management of Blood Pressure in Patients with Septic AKI', *Journal of Clinical Medicine*, 12(3), p. 1018.

- Erdogan, M. and Findikli, H.A. (2021) 'The Relationship Between Hypoalbuminemia and Mortality in Geriatric Patients with Sepsis', *Journal of Critical and Intensive Care*, 12(3), p. 80.
- Evans, L. *et al.* (2021) 'Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021', *Intensive care medicine*, 47(11), pp. 1181–1247.
- Fordtran, J.S. *et al.* (no date) *Sugar Absorption Tests, with Special Reference to 3-0-Methyl-d-Glucose and d-Xylose*. Available at: <https://annals.org/>.
- Furukawa, M. *et al.* (2019) 'Sepsis patients with complication of hypoglycemia and hypoalbuminemia are an early and easy identification of high mortality risk', *Internal and Emergency Medicine*, 14(4), pp. 539–548. Available at: <https://doi.org/10.1007/s11739-019-02034-2>.
- Geboes, K., Geboes, K.P. and Maleux, G. (2001) 'Vascular anatomy of the gastrointestinal tract', *Best Practice and Research: Clinical Gastroenterology*, 15(1), pp. 1–14. Available at: <https://doi.org/10.1053/bega.2000.0152>.
- Grissom, C.K. *et al.* (2010) 'A modified sequential organ failure assessment score for critical care triage', *Disaster medicine and public health preparedness*, 4(4), pp. 277–284.
- Gromova, L. V, Fetissov, S.O. and Gruzdkov, A.A. (2021) 'Mechanisms of glucose absorption in the small intestine in health and metabolic diseases and their role in appetite regulation', *Nutrients*, 13(7), p. 2474.
- Gül, F. *et al.* (2017) 'Changing definitions of sepsis', *Turkish journal of anaesthesiology and reanimation*, 45(3), p. 129.
- Gunst, J. *et al.* (2023) 'Tight blood-glucose control without early parenteral nutrition in the ICU', *New England Journal of Medicine*, 389(13), pp. 1180–1190.
- Hall, J.E. (2017) *Guyton and Hall*. 12th edn, *Nucl. Phys*. 12th edn. Edited by Elsevier.
- Van Haren, F.M.P. *et al.* (2007) 'Gastrointestinal perfusion in septic shock', *Anaesthesia and intensive care*, 35(5), pp. 679–694.
- Harper, D. and Chandler, B. (2016) 'Splanchnic circulation', *BJA Education*, 16(2), pp. 66–71. Available at: <https://doi.org/10.1093/bjaceaccp/mkv017>.
- Hutajulu, S. V (2017) 'Feeding', *Dalam: Pangalila FJV, Mansjoer A, penyunting. Penatalaksanaan sepsis dan syok septik (optimalisasi Fashugbid)*. Jakarta: Perdici, pp. 28–32.
- Jakob, S.M. (2002) 'Clinical review: splanchnic ischaemia', *Critical Care*, 6(4), pp. 1–7.
- Jenkins, B., Calder, P.C. and Marino, L. V. (2022) 'A systematic review of the definitions and prevalence of feeding intolerance in critically ill adults', *Clinical Nutrition ESPEN*, 49, pp.

- Jimenez, C.E.A.O. (2023) ‘Sepsis and Microcirculation’, *The Sepsis Codex-E-Book*, p. 29.
- de Jong, P.R., González-Navajas, J.M. and Jansen, N.J.G. (2016) ‘The digestive tract as the origin of systemic inflammation’, *Critical care*, 20, pp. 1–12.
- Kato, R. and Pinsky, M.R. (2015) ‘Personalizing blood pressure management in septic shock’, *Annals of intensive care*, 5(1), pp. 1–10.
- Kemenkes, R.I. (2010) ‘Pedoman Penyelenggaraan Pelayanan Intensive Care Unit (ICU) di Rumah Sakit’, *Jakarta: Depkes RI* [Preprint].
- Khwannimit, B., Bhurayanontachai, R. and Vattanavanit, V. (2022) ‘Ability of a modified Sequential Organ Failure Assessment score to predict mortality among sepsis patients in a resource-limited setting’, *Acute and Critical Care*, 37(3), pp. 363–371.
- Koepsell, H. (2020) ‘Glucose Transporters In The SmallI’, pp. 1207–1248.
- Landzinski, J. *et al.* (2008) ‘Gastric motility function in critically ill patients tolerant vs intolerant to gastric nutrition’, *Journal of Parenteral and Enteral Nutrition*, 32(1), pp. 45–50.
- László, I. *et al.* (2015) ‘Sepsis: From Pathophysiology to Individualized Patient Care’, *Journal of Immunology Research*, 2015. Available at: <https://doi.org/10.1155/2015/510436>.
- Leone, M. *et al.* (2015) ‘Optimizing mean arterial pressure in septic shock: a critical reappraisal of the literature’, *Critical care*, 19(1), pp. 1–7.
- Luttikhof, J. *et al.* (2013) ‘the role of gastrointestinal hormones in the treatment of delayed gastric emptying in critically ill patients’, *Alimentary pharmacology & therapeutics*, 38(6), pp. 573–583.
- Mao, Z. *et al.* (2020) ‘Association between serum lactate levels and enteral feeding intolerance in septic patients treated with vasopressors: a retrospective cohort study’, *Annals of Translational Medicine*, 8(19).
- Marik, P.E. and Bellomo, R. (2013) ‘Stress hyperglycemia : an essential survival response !’, pp. 1–7.
- Matheson, P.J., Wilson, M.A. and Garrison, R.N. (2000) ‘Regulation of intestinal blood flow’, *Journal of Surgical Research*, 93(1), pp. 182–196.
- McClave, S.A. *et al.* (2020) ‘Gastrointestinal dysfunction and feeding intolerance in critical illness: do we need an objective scoring system?’, *Current gastroenterology reports*, 22, pp. 1–8.
- McNicol, L. *et al.* (2013) ‘Pilot alternating treatment design study of the splanchnic metabolic effects of two mean arterial pressure targets during cardiopulmonary bypass’, *British Journal of Anaesthesia*, 110(5), pp. 721–728. Available at: <https://doi.org/10.1093/bja/aes493>.

- Mendez, C.E. *et al.* (2013) 'Increased glycemic variability is independently associated with length of stay and mortality in noncritically ill hospitalized patients', *Diabetes care*, 36(12), pp. 4091–4097.
- Michael A. Gropper (2020) *Miller's Anesthesia*. Elsevier. Available at: www.elsevier.com/permissions.
- Montori, V.M., Bistrian, B.R. and McMahon, M.M. (2002) 'Hyperglycemia in acutely ill patients', *Jama*, 288(17), pp. 2167–2169.
- Putowski, Z. *et al.* (2023) 'Correlation Between Mean Arterial Pressure and Capillary Refill Time in Patients with Septic Shock: A Systematic Review and Meta-analysis', *Journal of Intensive Care Medicine*, p. 08850666231168038.
- Revelly, J.P. *et al.* (2001) 'Early metabolic and splanchnic responses to enteral nutrition in postoperative cardiac surgery patients with circulatory compromise', *Intensive Care Medicine*, 27(3), pp. 540–547. Available at: <https://doi.org/10.1007/s001340100855>.
- Rudiger, A. and Singer, M. (2007) 'Mechanisms of sepsis-induced cardiac dysfunction', *Critical Care Medicine*, 35(6), pp. 1599–1608. Available at: <https://doi.org/10.1097/01.CCM.0000266683.64081.02>.
- Sastroasmoro, S. and Ismael, S. (2014) 'Dasar-dasar metodologi penelitian klinis 5th edn', *Sagung Seto, Jakarta* [Preprint].
- Sim, J.A. *et al.* (2013) 'Mesenteric blood flow, glucose absorption and blood pressure responses to small intestinal glucose in critically ill patients older than 65 years', *Intensive care medicine*, 39, pp. 258–266.
- Simoës Covello, L.H. *et al.* (2020) 'Vasopressors and Nutrition Therapy: Safe Dose for the Outset of Enteral Nutrition?', *Critical Care Research and Practice*, 2020. Available at: <https://doi.org/10.1155/2020/1095693>.
- Singer, M. *et al.* (2016) 'The third international consensus definitions for sepsis and septic shock (Sepsis-3)', *Jama*, 315(8), pp. 801–810.
- Singer, P. *et al.* (2019) 'ESPEN guideline on clinical nutrition in the intensive care unit', *Clinical Nutrition*, 38(1), pp. 48–79. Available at: <https://doi.org/10.1016/j.clnu.2018.08.037>.
- Someya, N. *et al.* (2008) 'Blood flow responses in celiac and superior mesenteric arteries in the initial phase of digestion', *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 294(6), pp. R1790–R1796.
- Suzuki, T. *et al.* (2017) 'Sepsis-induced cardiac dysfunction and β -adrenergic blockade therapy for sepsis', *Journal of Intensive Care*, 5(1), pp. 1–10.
- Taeb, A.M., Hooper, M.H. and Marik, P.E. (2017) 'Sepsis: current definition, pathophysiology, diagnosis, and management', *Nutrition in Clinical Practice*, 32(3), pp. 296–308.

- Takeishi, S. *et al.* (2016) 'Hypoglycemia and glycemic variability are associated with mortality in non-intensive care unit hospitalized infectious disease patients with diabetes mellitus', *Journal of Diabetes Investigation*, 7(3), pp. 429–435.
- Thazhath, S.S. *et al.* (2014) 'Glucose absorption in small intestinal diseases', *Expert review of gastroenterology & hepatology*, 8(3), pp. 301–312.
- Thibault, R. *et al.* (2011) 'Cardiogenic shock and nutrition: safe?', *Intensive care medicine*, 37, pp. 35–45.
- Tirone, T.A. and Brunicardi, F.C. (2001) 'Overview of glucose regulation', *World Journal of Surgery*, 25(4), pp. 461–467. Available at: <https://doi.org/10.1007/s002680020338>.
- Trahair, L.G. *et al.* (2014) 'Impact of gastric emptying to the glycemic and insulinemic responses to a 75-g oral glucose load in older subjects with normal and impaired glucose tolerance', *Physiological reports*, 2(11), p. e12204.
- Wang, J. *et al.* (2021) 'Hypoglycemia and mortality in sepsis patients: A systematic review and meta-analysis', *Heart and Lung*, 50(6), pp. 933–940. Available at: <https://doi.org/10.1016/j.hrtlng.2021.07.017>.
- Wang, L. *et al.* (2022) 'Association of Vasopressors Dose Trajectories with Enteral Nutrition Tolerance in Patients with Shock: A Prospective Observational Study', *Nutrients*, 14(24). Available at: <https://doi.org/10.3390/nu14245393>.
- Wang, Q. and Zhang, G. (2024) 'Gastrointestinal symptoms in sepsis shock patients: diastolic arterial pressure matter'.
- Weimann, A. *et al.* (2017) 'ESPEN guideline: clinical nutrition in surgery', *Clinical nutrition*, 36(3), pp. 623–650.
- Wentowski, C., Mewada, N. and Nielsen, N.D. (2019) 'Sepsis in 2018: a review', *Anaesthesia & Intensive Care Medicine*, 20(1), pp. 6–13.
- Wischmeyer, P.E. (2020) 'Enteral Nutrition Can Be Given to Patients on Vasopressors', *Critical Care Medicine*, 48(1), pp. 122–125. Available at: <https://doi.org/10.1097/CCM.00000000000003965>.
- Xu, J.Y. *et al.* (2015) 'A high mean arterial pressure target is associated with improved microcirculation in septic shock patients with previous hypertension: A prospective open label study', *Critical Care*, 19(1), pp. 1–8. Available at: <https://doi.org/10.1186/s13054-015-0866-0>.
- Yahyapoor, F. *et al.* (2021) 'The prevalence and possible causes of enteral tube feeding intolerance in critically ill patients: a cross-sectional study', *Journal of Research in Medical Sciences*, 26(1), p. 60.