

DAFTAR PUSTAKA

Adegbite, B.R. *et al.* (2021) ‘A comparison of different scores for diagnosis and mortality prediction of adults with sepsis in Low-and-Middle-Income Countries: a systematic review and meta-analysis’, *eClinicalMedicine*, 42. Available at: <https://doi.org/10.1016/j.eclinm.2021.101184>.

Agnello, L. *et al.* (2021) ‘The value of a complete blood count (Cbc) for sepsis diagnosis and prognosis’, *Diagnostics*, 11(10), pp. 1–19. Available at: <https://doi.org/10.3390/diagnostics11101881>.

Akilli, N.B. *et al.* (2014) ‘Prognostic importance of neutrophil-lymphocyte ratio in critically ill patients: Short- and long-term outcomes’, *American Journal of Emergency Medicine*, 32(12), pp. 1476–1480. Available at: <https://doi.org/10.1016/j.ajem.2014.09.001>.

Angus, D.C. *et al.* (2001) ‘Epidemiology of severe sepsis in the United States: Analysis of incidence, outcome, and associated costs of care’, *Critical Care Medicine*, 29(7), pp. 1303–1310. Available at: <https://doi.org/10.1097/00003246-200107000-00002>.

Aung ZNH, Maung NL, Hlaing SW, Aung SM, Aung ZP, Min S *et al.* (2023). Sepsis and Septic Shock: Are Initial mSOFA Score, Point-of-Care Lactate, Neutrophil-Lymphocyte Ratio and Platelet Count Good Predictors for Mortality? *Int J Med Sci Clin Res Stud*, 03(01):56–62. doi:10.47191/ijmscrs/v3-i1-12.

Baradari, A.G. *et al.* (2016) ‘Comparison of Proposed Modified and Original Sequential Organ Failure Assessment Scores in Predicting ICU Mortality: A Prospective, Observational, Follow-Up Study’, *Scientifica*, 2016. Available at: <https://doi.org/10.1155/2016/7379325>.

Bu, X. *et al.* (2019) ‘Relation of neutrophil-to-lymphocyte ratio to acute kidney injury in patients with sepsis and septic shock: A retrospective study’, *International Immunopharmacology*, 70(169), pp. 372–377. Available at: <https://doi.org/10.1016/j.intimp.2019.02.043>.

Burdick, H. *et al.* (2020) ‘Effect of a sepsis prediction algorithm on patient mortality, length of stay and readmission: A prospective multicentre clinical outcomes evaluation of real-world patient data from US hospitals’, *BMJ Health and Care Informatics*, 27(1), pp. 1–8. Available at: <https://doi.org/10.1136/bmjhci-2019-100109>.

Bermejo-Martín JF, Tamayo E, Ruiz G, Andaluz-Ojeda D, Herrán-Monge R, Muriel-Bombín A *et al.* (2014). Circulating neutrophil counts and mortality in septic shock. *Crit Care*, 18(1):407. doi:10.1186/cc13728.

Can, E., Hamilcikan, Ş. and Cen, C. (2018) ‘The Value of Neutrophil to Lymphocyte Ratio and Platelet to Lymphocyte Ratio for Detecting Early-onset Neonatal Sepsis’, *Journal of Pediatric Hematology/Oncology*, 40(4), pp. e229–

e232. Available at: <https://doi.org/10.1097/MPH.0000000000001059>.

Dahlan, M.S. (2013) *Besar Sampel dan Cara Pengambilan Sampel*. Tiga. Jakarta: Salemba Medika.

Dharmawan, A. *et al.* (2021) 'The Outcome of Patients with Sepsis at Tarakan Hospital Central Jakarta in 2018', *Muhammadiyah Medical Journal*, 2(2), p. 49. Available at: <https://doi.org/10.24853/mmj.2.2.49-54>.

Drăgoescu, A.N. *et al.* (2022) 'Neutrophil to Lymphocyte Ratio (NLR)—A Useful Tool for the Prognosis of Sepsis in the ICU', *Biomedicines*, 10(1), pp. 1–9. Available at: <https://doi.org/10.3390/biomedicines10010075>.

Eguia, E. *et al.* (2019) 'Risk factors for infection and evaluation of Sepsis-3 in patients with trauma', *American Journal of Surgery*, 218(5), pp. 851–857. Available at: <https://doi.org/10.1016/j.amjsurg.2019.03.005>.

El Said, A.M., Fayed, A.M. and El-Reweny, E.M. (2020) 'Comparative study between complete blood picture indices and presepsin as early prognostic markers in septic shock patients', *Egyptian Journal of Anaesthesia*, 36(1), pp. 118–125. Available at: <https://doi.org/10.1080/11101849.2020.1789403>.

Emadi, S.A. *et al.* (2022) 'SOFA and modified SOFA score for accessing outcomes among trauma patients in intensive care unit', *International Journal of Surgery Open*, 47(July), p. 100559. Available at: <https://doi.org/10.1016/j.ijso.2022.100559>.

Evans, L. *et al.* (2021) 'Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021', *Intensive Care Medicine*, 47(11), pp. 1181–1247. Available at: <https://doi.org/10.1007/s00134-021-06506-y>.

Farkas, J.D. (2020) 'The complete blood count to diagnose septic shock', *Journal of Thoracic Disease*, 2(Suppl 1), pp. S16–S21. Available at: <https://doi.org/10.21037/jtd.2019.12.63>.

Fujishima, S., 2016. Organ dysfunction as a new standard for defining sepsis. *Inflammation and Regeneration*, 36(24), pp. 1-6.

Grissom, C.K. *et al.* (2010) 'A modified Sequential Organ Failure Assessment score for critical care triage', *Disaster Medicine and Public Health Preparedness*, 4(4), pp. 277–284. Available at: <https://doi.org/10.1001/dmp.2010.40>.

Haloho, A.B., Heru Irwanto, F. and Gusnaintin, R. (2017) 'Uji Diagnostik Rasio Neutrofil-Limfosit dibanding dengan Procalcitonin sebagai Biomarker Infeksi Bakteri Pasien Sepsis', *Anesthesia & Critical Care*, 35(3), pp. 121–131.

Ham, S.Y. *et al.* (2020) 'Prognostic value of neutrophil/lymphocyte ratio and mean platelet volume/platelet ratio for 1-year mortality in critically ill patients', *Scientific Reports*, 10(1), pp. 1–8. Available at: <https://doi.org/10.1038/s41598-020-78476-y>.

Hou, J. *et al.* (2021) 'TREM2 sustains macrophage-hepatocyte metabolic coordination in nonalcoholic fatty liver disease and sepsis', *Journal of Clinical*

Investigation, 131(4). Available at: <https://doi.org/10.1172/JCI135197>.

Huang, Z. *et al.* (2020) ‘Prognostic value of neutrophil-to-lymphocyte ratio in sepsis: A meta-analysis’, *American Journal of Emergency Medicine*, 38(3), pp. 641–647. Available at: <https://doi.org/10.1016/j.ajem.2019.10.023>.

Imaeda, T. *et al.* (2021) ‘Trends in the incidence and outcome of sepsis using data from a Japanese nationwide medical claims database-the Japan Sepsis Alliance (JaSA) study group-’, *Critical Care*, 25(1), pp. 1–9. Available at: <https://doi.org/10.1186/s13054-021-03762-8>.

de Jager, C.P.C. *et al.* (2010) ‘Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia better than conventional infection markers in an emergency care unit’, *Critical Care*, 14(5). Available at: <https://doi.org/10.1186/cc9309>.

Jarczak, D., Kluge, S. and Nierhaus, A. (2021) ‘Sepsis—Pathophysiology and Therapeutic Concepts’, *Frontiers in Medicine*, 8(May), pp. 1–22. Available at: <https://doi.org/10.3389/fmed.2021.628302>.

Kari, K.A. *et al.* (2023) ‘Prevalence and Outcome of Sepsis: Mortality and Prolonged Intensive Care Unit Stay among Sepsis Patients Admitted to a Tertiary Centre in Malaysia’, *Malaysian Journal of Medical Sciences*, 30(6), pp. 120–132. Available at: <https://doi.org/10.21315/mjms2023.30.6.12>.

Kaushik, R. *et al.* (2018) ‘Diagnostic and prognostic role of neutrophil-to-lymphocyte ratio in early and late phase of sepsis’, *Indian Journal of Critical Care Medicine*, 22(9), pp. 660–663. Available at: https://doi.org/10.4103/ijccm.IJCCM_59_18.

Khwannimit, B., Bhurayanontachai, R. and Vattanavanit, V. (2022) ‘Ability of a modified Sequential Organ Failure Assessment score to predict mortality among sepsis patients in a resource-limited setting’, *Acute and Critical Care*, 37(3), pp. 363–371. Available at: <https://doi.org/10.4266/acc.2021.01627>.

Kim, J. *et al.* (2019) ‘Epidemiology of sepsis in Korea: A population-based study of incidence, mortality, cost and risk factors for death in sepsis’, *Clinical and Experimental Emergency Medicine*, 6(1), pp. 49–63. Available at: <https://doi.org/10.15441/ceem.18.007>.

Kurniawan, F. *et al.* (2021) ‘Platelet-lymphocyte ratio and sepsis outcomes in children’, *Paediatrica Indonesiana(Paediatrica Indonesiana)*, 61(6), pp. 322–327. Available at: <https://doi.org/10.14238/pi61.6.2021.322-7>.

Lambden, S. *et al.* (2019) ‘The SOFA score - Development, utility and challenges of accurate assessment in clinical trials’, *Critical Care*, 23(1), pp. 1–9. Available at: <https://doi.org/10.1186/s13054-019-2663-7>.

Lei, S. *et al.* (2022) ‘Prevalence of sepsis among adults in China: A systematic review and meta-analysis’, *Frontiers in Public Health*, 10. Available at: <https://doi.org/10.3389/fpubh.2022.977094>.

Li, A. *et al.* (2022) ‘Epidemiology, Management, and Outcomes of Sepsis in ICUs among Countries of Differing National Wealth across Asia’, *American Journal of Respiratory and Critical Care Medicine*, 206(9), pp. 1107–1116. Available at: <https://doi.org/10.1164/rccm.202112-2743OC>.

Li, Y. *et al.* (2021) ‘The Value of NLR Combined SOFA in Assessing the Prognosis of Sepsis Patients’, pp. 1–10. Available at: <https://doi.org/https://doi.org/10.21203/rs.3.rs-671911/v1>.

Li X, Chen Y, Yuan Q, Zhou H, Lu L, Guo R (2024). Neutrophil-to-Lymphocyte ratio, Monocyte-to-Lymphocyte ratio, Platelet-to-Lymphocyte ratio associated with 28-day mortality in sepsis patients with coronary artery disease. *BMC Infect Dis*, 24(1):749. doi:10.1186/s12879-024-09516-5.

Litell, J.M. *et al.* (2021) ‘Most emergency department patients meeting sepsis criteria are not diagnosed with sepsis at discharge’, *Academic Emergency Medicine*, 28(7), pp. 745–752. Available at: <https://doi.org/10.1111/acem.14265>.

Liu, S. *et al.* (2021) ‘Effects of Neutrophil-to-Lymphocyte Ratio Combined With Interleukin-6 in Predicting 28-Day Mortality in Patients With Sepsis’, *Frontiers in Immunology*, 12(March), pp. 1–9. Available at: <https://doi.org/10.3389/fimmu.2021.639735>.

Liu, X. *et al.* (2016) ‘Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Patients with Sepsis: A Prospective Observational Study’, *Mediators of Inflammation*, 2016. Available at: <https://doi.org/10.1155/2016/8191254>.

Liu, Y.C. *et al.* (2019) ‘Quick Sequential Organ Failure Assessment as a prognostic factor for infected patients outside the intensive care unit: a systematic review and meta-analysis’, *Internal and Emergency Medicine*, 14(4), pp. 603–615. Available at: <https://doi.org/10.1007/s11739-019-02036-0>.

Liu, Y.C. *et al.* (2022) ‘Frequency and mortality of sepsis and septic shock in China: a systematic review and meta-analysis’, *BMC Infectious Diseases*, 22(1), pp. 1–12. Available at: <https://doi.org/10.1186/s12879-022-07543-8>.

Loftus, T.J. *et al.* (2020) ‘Delayed interhospital transfer of critically ill patients with surgical sepsis’, *Journal of Trauma and Acute Care Surgery*, 88(1), pp. 169–175. Available at: <https://doi.org/10.1097/TA.0000000000002476>.

Lorente, L. *et al.* (2022) ‘Association between neutrophil-to-lymphocyte ratio in the first seven days of sepsis and mortality’, *Enfermedades Infecciosas y Microbiologia Clinica*, 40(5), pp. 235–240. Available at: <https://doi.org/10.1016/j.eimc.2020.11.004>.

Mandal, R.K. and Valenzuela, P.B. (2018) ‘Neutrophil-Lymphocyte count ratio on admission as a predictor of Bacteremia and In Hospital Mortality among Sepsis and Septic shock In Patients at Rizal Medical Center’, *Asian Journal of Medical Sciences*, 9(3), pp. 36–40. Available at: <https://doi.org/10.3126/ajms.v9i3.19030>.

Mankowski, R.T. *et al.* (2020) ‘Older Sepsis Survivors Suffer Persistent Disability

Burden and Poor Long-Term Survival’, *Journal of the American Geriatrics Society*, 68(9), pp. 1962–1969. Available at: <https://doi.org/10.1111/jgs.16435>.

McSweeney, D., 2015. Identifying Acute Organ Dysfunction as a Marker of Severe Sepsis. Master of Science in Nursing Student Scholarship, p. 69.

Merdji, H. *et al.* (2023) ‘Sex and gender differences in intensive care medicine’, *Intensive Care Medicine*. Springer Berlin Heidelberg, pp. 1155–1167. Available at: <https://doi.org/10.1007/s00134-023-07194-6>.

Muhammad, R. *et al.* (2022) ‘Increase Mortality Rate in Sepsis Patients with qSOFA Score Greater than Two’, 72(November 2021), pp. 2174–2177.

Nasir, N. *et al.* (2015) ‘Mortality in sepsis and its relationship with gender’, *Pakistan Journal of Medical Sciences*, 31(5), pp. 1201–1206. Available at: <https://doi.org/10.12669/pjms.315.6925>.

Octora, M. *et al.* (2021) ‘Predictive Score Model of Clinical Outcomes Sepsis in Intensive Care Unit Tertier Referral Hospital of Eastern Indonesia’, *Open Access Macedonian Journal of Medical Sciences*, 9(Apache Ii), pp. 1710–1716. Available at: <https://doi.org/10.3889/oamjms.2021.7780>.

Oda, S. *et al.* (2000) ‘Comparison of Sepsis-related Organ Failure Assessment (SOFA) score and CIS (cellular injury score) for scoring of severity for patients with multiple organ dysfunction syndrome (MODS)’, *Intensive Care Medicine*, 26(12), pp. 1786–1793. Available at: <https://doi.org/10.1007/s001340000710>.

Oh AR, Ryu J-A, Lee SJ, Kim CS, Lee SM (2024). Association between Inflammation-Based Prognostic Markers and Mortality in Patients Admitted to Intensive Care Units. *Diagnostics*, 14(16):1709. doi:10.3390/diagnostics14161709.

Pathania, R. *et al.* (2024) ‘Neutrophil-to-lymphocyte ratio as marker of organ dysfunction severity in patients with sepsis in the intensive care unit: An observational study’, *Journal of Clinical and Scientific Research*, 13(1), pp. 35–39. Available at: https://doi.org/10.4103/jcsr.jcsr_259_22.

Purba, A.K.R. *et al.* (2020) ‘The burden and costs of sepsis and reimbursement of its treatment in a developing country: An observational study on focal infections in Indonesia’, *International Journal of Infectious Diseases*, 96, pp. 211–218. Available at: <https://doi.org/10.1016/j.ijid.2020.04.075>.

Rahmatinejad, Z. *et al.* (2019) ‘Predictive performance of the SOFA and mSOFA scoring systems for predicting in-hospital mortality in the emergency department’, *American Journal of Emergency Medicine*, 37(7), pp. 1237–1241. Available at: <https://doi.org/10.1016/j.ajem.2018.09.011>.

Rajbhandari, P., Shakya, Y.L. and Upreti, A.R. (2019) ‘Neutrophil to Lymphocytic Count and SOFA Score as Prognostic Marker in Patient With Sepsis’, *Nepal Medical Journal*, 1(2), pp. 43–47.

Ramachandran, G. (2014) ‘Gram-positive and gram-negative bacterial toxins in sepsis: A brief review’, *Virulence*, 5(1), pp. 213–218. Available at: <https://doi.org/10.4161/viru.27024>.

Raymond, N.J. *et al.* (2019) ‘Modified Sequential Organ Failure Assessment sepsis score in an emergency department setting: Retrospective assessment of prognostic value’, *EMA - Emergency Medicine Australasia*, 31(3), pp. 339–346. Available at: <https://doi.org/10.1111/1742-6723.13154>.

Rello Jordi, Valenzuela-Sanchez, F., Ruiz-Rodriguez, M. & Moyano, S., (2017). Sepsis: A Review Of Advances in Management. *Adv Ther*, Volume 34, pp. 2393-2411.

Rehman, F.U. *et al.* (2020) ‘Neutrophils to Lymphocyte Ratio: Earliest and Efficacious Markers of Sepsis’, *Cureus*, 12(10), pp. 1–7. Available at: <https://doi.org/10.7759/cureus.10851>.

Rudd, K.E. *et al.* (2020) ‘Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study’, *The Lancet*, 395(10219), pp. 200–211. Available at: [https://doi.org/10.1016/S0140-6736\(19\)32989-7](https://doi.org/10.1016/S0140-6736(19)32989-7).

Sakr, Y. *et al.* (2013) ‘The influence of gender on the epidemiology of and outcome from severe sepsis’, *Critical Care*, 17(2). Available at: <https://doi.org/10.1186/cc12570>.

Saliccioli JD, Marshall DC, Pimentel MA, Santos MD, Pollard T, Celi LA *et al.* (2015). The association between the neutrophil-to-lymphocyte ratio and mortality in critical illness: an observational cohort study. *Crit Care*, 19(1):13. doi:10.1186/s13054-014-0731-6.

Shukeri, W.F.W.M., Nor, M.B.M. and Ralib, A.M. (2022) ‘Sepsis and Its Impact on Outcomes in Elderly Patients Admitted to a Malaysian Intensive Care Unit’, *Malaysian Journal of Medical Sciences*, 29(3), pp. 145–150. Available at: <https://doi.org/10.21315/mjms2022.29.3.14>.

Shi, Y., Yang, C., Chen, L., Cheng, M. and Xie, W. (2022). Predictive value of neutrophil-to-lymphocyte and platelet ratio in in-hospital mortality in septic patients. *Heliyon*, 8(11), p.e11498. doi:<https://doi.org/10.1016/j.heliyon.2022.e11498>.

Shrestha, G.S., Maharjan, A., Shrestha, R. and Nepal, G. (2021). Comparison of modified Sequential Organ Failure Assessment (mSOFA) score with Sequential Organ Failure Assessment (SOFA) score to predict mortality in critically ill patients. *Bangladesh Critical Care Journal*, 9(2), pp.74–78. doi:<https://doi.org/10.3329/bccj.v9i2.56153>.

Siloși, C.A. *et al.* (2018) ‘Sepsis and identification of reliable biomarkers for postoperative period prognosis’, *Romanian Journal of Morphology and Embryology*, 59(1), pp. 77–91.

Singer, M., Deutschman, C. S., Seymour, C. W. & Shankar-Hari, M., (2016). The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*, Volume 315, pp. 801-10.

Taylor, S.P. *et al.* (2020) 'Association between adherence to recommended care and outcomes for adult survivors of sepsis', *Annals of the American Thoracic Society*, 17(1), pp. 89–97. Available at: <https://doi.org/10.1513/AnnalsATS.201907-514OC>.

Terradas, R. *et al.* (2012) 'Eosinophil count and neutrophil-lymphocyte count ratio as prognostic markers in patients with bacteremia: A retrospective cohort study', *PLoS ONE*, 7(8), pp. 1–8. Available at: <https://doi.org/10.1371/journal.pone.0042860>.

Viglino, D., Maignan, M. and Debaty, G. (2016) 'A modified sequential organ failure assessment score using the richmond agitation-sedation scale in critically ill patients', *Journal of Thoracic Disease*, 8(3), pp. 311–313. Available at: <https://doi.org/10.21037/jtd.2016.02.61>.

Wang, X. *et al.* (2023) 'Application Prospect of the SOFA Score and Related Modification Research Progress in Sepsis', *Journal of Clinical Medicine*, 12(10). Available at: <https://doi.org/10.3390/jcm12103493>.

Wardi, G. *et al.* (2021) 'Age-related incidence and outcomes of sepsis in California, 2008–2015', *Journal of Critical Care*, 62, pp. 212–217. Available at: <https://doi.org/10.1016/j.jcrc.2020.12.015>.

Wei Z, Zhao L, Yan J, Wang X, Li Q, Ji Y *et al.* (2024). Dynamic monitoring of neutrophil/lymphocyte ratio, APACHE II score, and SOFA score predict prognosis and drug resistance in patients with *Acinetobacter baumannii*–calcoaceticus complex bloodstream infection: a single-center retrospective study. *Front Microbiol*, 15. doi:10.3389/fmicb.2024.1296059.

Westerdijk, K., Simons, K.S., Zegers, M., Wever, P.C., Pickkers, P. and de Jager, C.P.C. (2019). The value of the neutrophil-lymphocyte count ratio in the diagnosis of sepsis in patients admitted to the Intensive Care Unit: A retrospective cohort study. *PLOS ONE*, 14(2), p.e0212861. doi:<https://doi.org/10.1371/journal.pone.0212861>.

Wolfertz, N. *et al.* (2022) 'Epidemiology, management, and outcome of infection, sepsis, and septic shock in a German emergency department (EpiSEP study)', *Frontiers in Medicine*, 9(October). Available at: <https://doi.org/10.3389/fmed.2022.997992>.

Xie, J. *et al.* (2020) 'The Epidemiology of Sepsis in Chinese ICUs: A National Cross-Sectional Survey', *Critical Care Medicine*, 48(3), pp. E209–E218. Available at: <https://doi.org/10.1097/CCM.0000000000004155>.

Yang, Y. *et al.* (2010) 'The effect of comorbidity and age on hospital mortality and length of stay in patients with sepsis', *Journal of Critical Care*, 25(3), pp. 398–405.

Available at: <https://doi.org/10.1016/j.jcrc.2009.09.001>.

Zhang, H.B. *et al.* (2016) 'Diagnostic values of red cell distribution width, platelet distribution width and neutrophil-lymphocyte count ratio for sepsis', *Experimental and Therapeutic Medicine*, 12(4), pp. 2215–2219. Available at: <https://doi.org/10.3892/etm.2016.3583>.

Zhang, M.Q. *et al.* (2021) 'Sex- and Gender-Dependent Differences in Clinical and Preclinical Sepsis', *Shock*, 56(2), pp. 178–187. Available at: <https://doi.org/10.1097/SHK.0000000000001717>.

Zhang, Y., Peng, W. and Zheng, X. (2024) 'The prognostic value of the combined neutrophil-to-lymphocyte ratio (NLR) and neutrophil-to-platelet ratio (NPR) in sepsis', *Scientific Reports*, 14(1), pp. 1–10. Available at: <https://doi.org/10.1038/s41598-024-64469-8>.

Zhang, Z. *et al.* (2019) 'Prolonged length of stay in the emergency department and increased risk of hospital mortality in patients with sepsis requiring ICU admission', *Emergency Medicine Journal*, 36(2), pp. 82–87. Available at: <https://doi.org/10.1136/emmermed-2018-208032>.