

DAFTAR PUSTAKA

- Andrew, E., Emyedu, A., Atumanya, P., Okello, E., Wabule, A., Ssemogerere, L., Mukisa, J., & Sendagire, C. (2022). Prevalence and Complications of Residual Neuromuscular Blockade in the Post Anesthesia Care Units of Uganda. *Research Article Journal of Anesthesia and Clinical Research J Clin Trials*, 13(2), 1001045–1001046. <https://doi.org/10.35248/2155-6148.22.13.1045>
- Ardyan Wardhana, Juni Kurniawaty, Y. U. (2019). Basics of CPB. *Indian Journal of Anaesthesia*, 49(4), 257–262. <https://doi.org/10.4103/ija.IJA>
- Azam Danish, M. (2021). Preoxygenation and Anesthesia: A Detailed Review. *Cureus*. <https://doi.org/10.7759/cureus.13240>
- Baillard, C., Clec'h, C., Catineau, J., Salhi, F., Gehan, G., Cupa, M., & Samama, C. M. (2005). Postoperative residual neuromuscular block: A survey of management. *British Journal of Anaesthesia*, 95(5), 622–626. <https://doi.org/10.1093/bja/aei240>
- Baurain, M. (1991). Effects of Residual Concentrations of Isoflurane on the Reversal of Vecuronium-induced Neuromuscular Blockade. *Anesthesiology*, 74, 474–478.
- Baurain, M. J., Hoton, F., D'hollander, A. A., & Cantraine, F. R. (1996). Is recovery of neuromuscular transmission complete after the use of neostigmine to antagonize block produced by rocuronium, vecuronium, atracurium and pancuronium? In *British Journal of Anaesthesia* (Vol. 77).
- Bellini, L., Banzato, T., Contiero, B., & Zotti, A. (2014). Evaluation of sedation and clinical effects of midazolam with ketamine or dexmedetomidine in pet rabbits. *Veterinary Record*, 175(15), 372–372.
- Bohringer, C., & Liu, H. (2019). Is it always necessary to reverse the neuromuscular blockade at the end of surgery? In *Journal of Biomedical Research* (Vol. 33, Issue 4, pp. 217–220). Nanjing Medical University and Chungbuk National University Press. <https://doi.org/10.7555/JBR.33.20180123>
- Boles, J. M., Bion, J., Connors, A., Herridge, M., Marsh, B., Melot, C., Pearl, R., Silverman, H., Stanchina, M., Vieillard-Baron, A., & Welte, T. (2007).

- Weaning from mechanical ventilation. In *European Respiratory Journal* (Vol. 29, Issue 5, pp. 1033–1056). <https://doi.org/10.1183/09031936.00010206>
- Bowdle, A., Bussey, L., Michaelsen, K., Jelacic, S., Nair, B., Togashi, K., & Hulvershorn, J. (2020). Counting train-of-four twitch response: comparison of palpation to mechanomyography, acceleromyography, and electromyography. *British Journal of Anaesthesia*, 124(6), 712–717. <https://doi.org/10.1016/j.bja.2020.02.022>
- Bronsert, M. R., Henderson, W. G., Monk, T. G., Richman, J. S., Nguyen, J. D., Sum-Ping, J. T., Mangione, M. P., Higley, B., & Hammermeister, K. E. (2017). Intermediate-acting nondepolarizing neuromuscular blocking agents and risk of postoperative 30-day morbidity and mortality, and long-term survival. *Anesthesia and Analgesia*, 124(5), 1476–1483. <https://doi.org/10.1213/ANE.0000000000001848>
- Brull, K. (2017). Current Status of Neuromuscular Reversal and Monitoring: Posttetanic Neuromonitoring and Other Considerations. *Anesthesiology*, 127(4), 723–724. <https://doi.org/10.1097/ALN.0000000000001806>
- Brull, S. J., & Kopman, A. F. (2017). Current Status of Neuromuscular Reversal and Monitoring. In *Anesthesiology* (Vol. 126, Issue 1, pp. 173–190). Lippincott Williams and Wilkins. <https://doi.org/10.1097/ALN.0000000000001409>
- Brull, S. J., & Murphy, G. S. (2010). Residual neuromuscular block: Lessons unlearned. Part ii: Methods to reduce the risk of residual weakness. In *Anesthesia and Analgesia* (Vol. 111, Issue 1, pp. 129–140). Lippincott Williams and Wilkins. <https://doi.org/10.1213/ANE.0b013e3181da8312>
- Brull, S., & Meistelman, C. (2019). Pharmacology of Neuromuscular Blocking Drugs. In *Miller's Anesthesia* (9th ed., pp. 792–831). Elsevier.
- Bulka, C. M., Terekhov, M. A., Martin, B. J., Dmochowski, R. R., Hayes, R. M., & Ehrenfeld, J. M. (2016). Nondepolarizing Neuromuscular Blocking Agents, Reversal, and Risk of Postoperative Pneumonia. *Anesthesiology*, 125(4), 647–655. <https://doi.org/10.1097/ALN.0000000000001279>

- Butterly, A., Bittner, E. A., George, E., Sandberg, W. S., Eikermann, M., & Schmidt, U. (2010). Postoperative residual curarization from intermediate-acting neuromuscular blocking agents delays recovery room discharge. *British Journal of Anaesthesia*, 105(3), 304–309. <https://doi.org/10.1093/bja/aeq157>
- Butterworth, J. F., Mackey, D. C., Wasnick, J. D., Morgan, G. E., Mikhail, M. S., & Morgan, G. Edward. (2018). *Morgan & Mikhail's clinical anesthesiology*.
- Cakar, N., Tuorul, M., Demirarslan, A., Nahum, A., Adams, A., Esen, F., & Telci, L. (2001). Time required for partial pressure of arterial oxygen equilibration during mechanical ventilation after a step change in fractional inspired oxygen concentration. *Intensive Care Medicine*, 27(4), 655–659. <https://doi.org/10.1007/s001340100900>
- Caldwell, J. E., Robertson, E. N., & Baird, W. L. M. (1986). ANTAGONISM OF PROFOUND NEUROMUSCULAR BLOCKADE INDUCED BY VECURONIUM* OR ATRACURIUM Comparison of Neostigmine with Edrophonium. In *Br.J. Anaesth* (Vol. 58). <http://bjaoxfordjournals.org/>
- Chawla, R., & Todi, S. (2019). ICU Protocols. In *ICU Protocols: A Step-wise Approach, Vol I*. Springer Singapore. <https://doi.org/10.1007/978-981-15-0898-1>
- Cheng, C. R., Sessler, D. I., & Apfel, C. C. (2005). Does neostigmine administration produce a clinically important increase in postoperative nausea and vomiting? *Anesthesia and Analgesia*, 101(5), 1349–1355. <https://doi.org/10.1213/01.ANE.0000180992.76743.C9>
- Chuanxiang, Yao Shanghmg., Nie Hui, & Lu Bin. (2004). Effects of Isoflurane on the Actions of Neuromuscular Blockers on the Muscle Nicotine Acetylcholine Receptors. *Journal of Huazhong University of Science and Technology*, 21(6), 605–606.
- Crul, J. F. (1982). The Effect of Inhalation Anaesthetics on Skeletal and Smooth Muscle. *Springer-Verlag Berlin Heidelberg, Inhalation Anaesthesia Today and Tomorrow*.

- Donati, F. (2000). Neuromuscular Blocking Drugs for the New Millennium: Current Practice, Future Trends-Comparative Pharmacology of Neuromuscular Blocking Drugs. *ANESTH ANALG*, 90, 2–6.
- Driessen, J. J., Crul, J. F., Jansen, R., & van Egmond, J. (1986). Isoflurane and Neuromuscular Blocking Drugs. *Anaesthesiology Intensive Care Medicine/Anaesthesiologie Und Intensivmedizin*, 182.
- Eikermann, M., Groeben, H., Hüsing, J., & Peters, J. (2003). Accelerometry of Adductor Pollicis Muscle Predicts Recovery of Respiratory Function from Neuromuscular Blockade. In *Anesthesiology* (Vol. 98). <http://pubs.asahq.org/anesthesiology/article-pdf/98/6/1333/653565/0000542-200306000-00006.pdf>
- Farhan, H., Moreno-Duarte, I., McLean, D., & Eikermann, M. (2014a). Residual Paralysis: Does it Influence Outcome After Ambulatory Surgery? *Current Anesthesiology Reports*, 4(4), 290–302. <https://doi.org/10.1007/s40140-014-0073-6>
- Fortier, L. P., Mckeen, D., Turner, K., De Médicis, É., Warriner, B., Jones, P. M., Chaput, A., Pouliot, J. F., & Galarneau, A. (2015). The RECITE study: A Canadian prospective, multicenter study of the incidence and severity of residual neuromuscular blockade. *Anesthesia and Analgesia*, 121(2), 366–372. <https://doi.org/10.1213/ANE.0000000000000757>
- Fu, E. S., Downs, J. B., Schweiger, J. W., Miguel, R. V., & Smith, R. A. (2004). Supplement oxygen impairs detection of hypoventilation by pulse oximetry. *Chest*, 126(5), 1552–1558. <https://doi.org/10.1378/chest.126.5.1552>
- Grayling, M., & Sweeney, B. P. (2007). Recovery from neuromuscular blockade: A survey of practice. *Anaesthesia*, 62(8), 806–809. <https://doi.org/10.1111/j.1365-2044.2007.05101.x>
- Grosse-Sundrup, M., Henneman, J. P., Sandberg, W. S., Bateman, B. T., Uribe, J. V., Nguyen, N. T., Ehrenfeld, J. M., Martinez, E. A., Kurth, T., & Eikermann, M. (2012). Intermediate acting non-depolarizing neuromuscular blocking agents and risk of postoperative respiratory complications: Prospective

- propensity score matched cohort study. *BMJ (Online)*, 345(7880).
<https://doi.org/10.1136/bmj.e6329>
- Hahn, S. (2012). Understanding noninferiority trials. In *Korean Journal of Pediatrics* (Vol. 55, Issue 11, pp. 403–407).
<https://doi.org/10.3345/kjp.2012.55.11.403>
- Hedenstierna, G., & Edmark, L. (2015). Effects of anesthesia on the respiratory system. In *Best Practice and Research: Clinical Anaesthesiology* (Vol. 29, Issue 3, pp. 273–284). Bailliere Tindall Ltd.
<https://doi.org/10.1016/j.bpa.2015.08.008>
- Herbstreit, F., Zigran, D., Ochterbeck, C., rgen Peters, J., & Eikermann, M. (2010). PERIOPERATIVE MEDICINE Neostigmine/Glycopyrrolate Administered after Recovery from Neuromuscular Block Increases Upper Airway Collapsibility by Decreasing Genioglossus Muscle Activity in Response to Negative Pharyngeal Pressure. In *Anesthesiology* (Vol. 113).
www.anesthesiology.org
- Hsu, C., Hsu, Y., Huang, G., Lu, C., Ho, S., Liaw, W., Tsai, Y., Lin, C., Tsai, C., & Lin, T. (2013). *Iso fl urane compared with fentanyl-midazolam- based anesthesia in patients undergoing heart transplantation. May*.
- Islam, M. A., Shahida, S. M., Islam, M. R., Haque, M. M., Rashid, M. H., & Rashid, M. H. (2011). Different doses of atropine on heart rate fluctuations during reversal of neuromuscular blockade. *Mymensingh Medical Journal : MMJ*, 20(4), 595–599.
- Jemea B, Nomo S, N., C, I., Tamchom D, D., G, C., B, M., S, N., Minkande J, Z., & F, B. (2022). Postoperative Residual Curarization in Difficult Environment for the Practice of General Anesthesia: Inventory of the Yaounde/Essos Hospital Center. *Anesthesia and Critical Care*, 04(01).
<https://doi.org/10.26502/acc.039>
- Jubb, A., & Ford, P. (2009). Extubation after Anaesthesia: A Systematic Review. *Update in Anaesthesia*, 30–36. www.anaesthesiologists.org
- Kaufhold, N., Schaller, S. J., Stäuble, C. G., Baumüller, E., Ulm, K., Blobner, M., & Fink, H. (2016). Sugammadex and neostigmine dose-finding study for

reversal of residual neuromuscular block at a train-of-four ratio of 0.2 (SUNDRO20). *British Journal of Anaesthesia*, 116(2), 233–240. <https://doi.org/10.1093/bja/aev437>

Kim, K. S., Cheong, M. A., Lee, H. J., & Lee, J. M. (2004). Tactile assessment for the reversibility of rocuronium-induced neuromuscular blockade during propofol or sevoflurane anesthesia. *Anesthesia and Analgesia*, 99(4), 1080–1085. <https://doi.org/10.1213/01.ANE.0000130616.57678.80>

Kopman, A. F. (2010). Managing neuromuscular block: Where are the guidelines? In *Anesthesia and Analgesia* (Vol. 111, Issue 1, pp. 9–10). Lippincott Williams and Wilkins. <https://doi.org/10.1213/ANE.0b013e3181cdb0a5>

Kopman, A. F. (2016). Residual Neuromuscular Blockade and Adverse Postoperative Outcomes: An Update. *Current Anesthesiology Reports*, 6(2), 178–184. <https://doi.org/10.1007/s40140-016-0151-z>

Kopman, A. F., Zank, L. M., Ng, J., & Neuman, G. G. (2004). Antagonism of Cisatracurium and Rocuronium Block at a Tactile Train-of-Four Count of 2: Should Quantitative Assessment of Neuromuscular Function Be Mandatory? *Anesthesia and Analgesia*, 98(1), 102–106. <https://doi.org/10.1213/01.ANE.0000094985.19305.E9>

Lee, S., Ro, Y.J., Koh, W.U., Nishiyama, T., Yang, H. (2016). The neuromuscular effect of rocuronium under sevoflurane-remifentanyl or propofol-remifentanyl anesthesia: a randomized clinical comparative study in an Asia population. *BMC Anesthesiology*.

Lemmens, H. J., El-Orbany, M. I., Berry, J., Ben Morte Jr, J., & Martin, G. (2010). *Reversal of profound vecuronium-induced neuromuscular block under sevoflurane anesthesia: sugammadex versus neostigmine*. <http://www.biomedcentral.com/1471-2253/10/15>

Lowry, D. W., Mirakhur, R. K., McCarthy, G. J., Carroll, M. T., & McCourt, K. C. (1998). Neuromuscular Effects of Rocuronium During Sevoflurane, Isoflurane, and Intravenous Anesthesia. *Anesth Analg*, 87, 936–940.

- Malik, V., Smith, D., & Lee-Chiong, T. (2012). Respiratory physiology during sleep. In *Sleep Medicine Clinics* (Vol. 7, Issue 3, pp. 497–505). W.B. Saunders. <https://doi.org/10.1016/j.jsmc.2012.06.011>
- McLean, D. J., Diaz-Gil, D., Farhan, H. N., Ladha, K. S., Kurth, T., & Eikermann, M. (2015). Dose-dependent Association between Intermediate-acting Neuromuscular-blocking Agents and Postoperative Respiratory Complications. *Anesthesiology*, 122, 1201–1213.
- Miller RD. (2015). *Miller's Anesthesia* (8th ed.). Elsevier/Saunders.
- Murphy, G. S., Szokol, J. W., Avram, M. J., Greenberg, S. B., Marymont, J. H., Vender, J. S., Gray, J., Landry, E., & Gupta, D. K. (2011). *Intraoperative Acceleromyography Monitoring Reduces Symptoms of Muscle Weakness and Improves Quality of Recovery in the Early Postoperative Period*. www.anesthesiology.org.
- Murphy, G. S., Szokol, J. W., Avram, M. J., Greenberg, S. B., Shear, T. D., Vender, J. S., Parikh, K. N., Patel, S. S., & Patel, A. (2015). Residual Neuromuscular Block in the Elderly. *Anesthesiology*, 123, 1322–1336.
- Murphy, G. S., Szokol, J. W., Marymont, J. H., Franklin, M., Avram, M. J., & Vender, J. S. (2005). Residual paralysis at the time of tracheal extubation. *Anesthesia and Analgesia*, 100(6), 1840–1845. <https://doi.org/10.1213/01.ANE.0000151159.55655.CB>
- Murphy, G. S., Szokol, J. W., Marymont, J. H., Greenberg, S. B., Avram, M. J., & Vender, J. S. (2008). Residual neuromuscular blockade and critical respiratory events in the postanesthesia care unit. *Anesthesia and Analgesia*, 107(1), 130–137. <https://doi.org/10.1213/ane.0b013e31816d1268>
- Murphy, GS., Brull, SJ. (2010). Residual neuromuscular Block: Lessons Unlearned. Part II: Metode to reduce the risk of residual weakness. *International Anesthesia Research Society*, 1, 129–131.
- Murphy, GS., Brull, SJ. (2021). Quantitative Neuromuscular Monitoring and Postoperative Outcomes: A Narrative Review. *The American Society of Anesthesiologists*, 348–349.

- MUSLU ERDEM, V., SAYAN, H. E., & KORFALI, S. G. (2021). Comparison of the effects of isoflurane and sevoflurane on surgical stress in intracranial tumor surgery. *The European Research Journal*, 7(2), 142–151. <https://doi.org/10.18621/eurj.648843>
- Naguib, M., Kopman, A. F., & Ensor, J. E. (2007). Neuromuscular monitoring and postoperative residual curarisation: A meta-analysis. *British Journal of Anaesthesia*, 98(3), 302–316. <https://doi.org/10.1093/bja/ael386>
- Naguib, M., Kopman, A. F., Lien, C. A., Hunter, J. M., Lopez, A., & Brull, S. J. (2010). A survey of current management of neuromuscular block in the United States and Europe. In *Anesthesia and Analgesia* (Vol. 111, Issue 1, pp. 110–119). Lippincott Williams and Wilkins. <https://doi.org/10.1213/ANE.0b013e3181c07428>
- Naguib, M., Lien, CA., & Meistelman, C. (2015). Pharmacology of Neuromuscular Blocking Drugs. In *Miller's Anesthesia* (8th ed., pp. 958–994). Elsevier/Saunders.
- Naik, S. S., Rajappa, G. C., Rao, R. N., & B, Y. (2015). Incidence of Residual Neuromuscular Blockade At Tracheal Extubation: Comparison of Atracurium With Vecuronium. *Journal of Evolution of Medical and Dental Sciences*, 4(94), 15947–15951. <https://doi.org/10.14260/jemds/2015/2323>
- Nasibova, E. M., & Polukhov, R. S. (2021). Comparative study of the potentiating effect of isoflurane and sevoflurane on the duration of muscle relaxation during surgical interventions in one-day surgery in children. *Pediatric Anesthesia and Critical Care Journal*, 9(1), 21–29. <https://doi.org/10.14587/paccj.2021.4>
- Nolan, J. P. (2012). Anaesthesia and neuromuscular block. In P. Bennet, M. Brown, & P. Sharma (Eds.), *Clinical Pharmacology* (11th ed., pp. 295–310). Elsevier. <https://doi.org/10.1016/B978-0-7020-4084-9.00058-6>
- Parotto, M., Brown, C. A., Sakles, J., Mick, N., Mosier, J., & Braude, D. (2023). Safe Extubation in the Emergency Airway Patient. In *The Walls Manual Of Emergency Airway Management* (6th ed.). Wolters Kluwer.

- Plaud, B., Baillard, C., Bourgain, J. L., Bouroche, G., Desplanque, L., Devys, J. M., Fletcher, D., Fuchs-Buder, T., Lebuffe, G., Meistelman, C., Motamed, C., Raft, J., Servin, F., Sirieix, D., Slim, K., Velly, L., Verdonk, F., & Debaene, B. (2020). Guidelines on muscle relaxants and reversal in anaesthesia. *Anaesthesia Critical Care and Pain Medicine*, 39(1), 125–142. <https://doi.org/10.1016/j.accpm.2020.01.005>
- Rodney G, Raju PKBC, & Ball DR. (2015). *Editorial Not just monitoring; a strategy for managing neuromuscular blockade*. www.aagbi.org
- Stoelting RK, Shafer SL, Rathmell JP, & Flood P. (2015). *Pharmacology and Physiology in Anesthetic Practice* (3rd ed.). Wolters Kluwer.
- Sturgill, E. L., & Campbell, N. (2015). Neuromuscular Blocking and Reversal Agents. In *Basic Clinical Anesthesia* (5th ed., pp. 151–158). Springer.
- Tan, X., & Lu, Y. (2022). *Simple grip strength device for assessing the degree of recovery from residual neuromuscular blockade after general anaesthesia : An observational study*. 1–13.
- Thilen, S. R., & Bhananker, S. M. (2016). Qualitative Neuromuscular Monitoring: How to Optimize the Use of a Peripheral Nerve Stimulator to Reduce the Risk of Residual Neuromuscular Blockade. *Current Anesthesiology Reports*, 6(2), 164–169. <https://doi.org/10.1007/s40140-016-0155-8>
- Thilen, S. R., Sherpa, J. R., James, A. M., Cain, K. C., Treggiari, M. M., & Bhananker, S. M. (2023). Management of Muscle Relaxation With Rocuronium and Reversal With Neostigmine or Sugammadex Guided by Quantitative Neuromuscular Monitoring. *Anesthesia & Analgesia, Publish Ahead of Print*(3), 536–544. <https://doi.org/10.1213/ane.00000000000006511>
- Tramè, M. R., & Fuchs-Buder, T. (1999). Omitting antagonism of neuromuscular block: effect on postoperative nausea and vomiting and risk of residual paralysis. A systematic review. In *British Journal of Anaesthesia* (Vol. 82, Issue 3).
- Uzawa, K., Seki, H., & Yorozu, T. (2021). Residual paralysis caused by 50 mg rocuronium after reversal with 4 mg/kg sugammadex: a case report. *BMC Anesthesiology*, 21(1). <https://doi.org/10.1186/s12871-021-01379-6>

- Vanlinthout, L. E. H., Booij, L. H. D. J., Van Egmond, J., & Robertson, E. N. (1996). Effect of isoflurane and sevoflurane on the magnitude and time course of neuromuscular block produced by vecuronium, pancuronium and atracurium. *British Journal of Anaesthesia*, 76(3), 389–395. <https://doi.org/10.1093/bja/76.3.389>
- Videira, R. L. R., & Vieira, J. E. (2011). What rules of thumb do clinicians use to decide whether to antagonize nondepolarizing neuromuscular blocking drugs? *Anesthesia and Analgesia*, 113(5), 1192–1196. <https://doi.org/10.1213/ANE.0b013e31822c986e>
- Walker, E., & Nowacki, A. S. (2011). Understanding equivalence and noninferiority testing. In *Journal of General Internal Medicine* (Vol. 26, Issue 2, pp. 192–196). <https://doi.org/10.1007/s11606-010-1513-8>
- Wardhana, A., Kurniawaty, J., & Uyun, Y. (2019). Optimised reversal without train-of-four monitoring versus reversal using quantitative train-of-four monitoring: An equivalence study. *Indian Journal of Anaesthesia*, 63(5), 361. https://doi.org/10.4103/ija.ija_94_19
- Willis, D. (2019). *Anesthesia Providers' Knowledge and Utilization of a Quantitative Monitor for Neuromuscular Function Recovery*.
- Wulf, H., Ledowski, T., & Linstedt, U. (1997). *Neuromuscular effects during isoflurane, block-of rocuroni-desflurane, and sevoflurane anaesthesia*.
- Yang, L., Hu, N., Chang, H., Yang, D., & Zuo, Y. (2023). Routine administration of neostigmine after recovery of spontaneous breathing versus neuromuscular monitor-guided administration of neostigmine in pediatric patients: a parallel, randomized, controlled study. *Trials*, 24(1), 1–6. <https://doi.org/10.1186/s13063-023-07066-w>
- Zhong, B. (2009). How to Calculate Sample Size in Randomized Controlled Trial? *Journal of Thoracic Disease*, 1, 51–54.