

## DAFTAR PUSTAKA

- Asali, F., & Phillips, C., 2012. The management of urogynaecological problems in pregnancy and the early postpartum period. *Obstet. Gynaecol.* 14: 153–158. doi:10.1111/j.1744-4667.2012.00111.x
- Aydin, S., & Aydin, C.A., 2015. Evaluation of labor-related pelvic floor changes 3 months after delivery : a 3D transperineal ultrasound study. *Int. Urogynecol. J.* doi:10.1007/s00192-015-2774-x
- Santoso, B., 2012. Budi Iman Santoso Assessment (BISA): a model for predicting levator ani injury after vaginal delivery. *Medical Journal of Indonesia*, p.102
- Barber, M.D., Walters, M.D., & Bump, R.C., 2005. Short forms of two condition- specific quality-of-life questionnaires for women with pelvic floor disorders (PFDI-20 and PFIQ-7). *Am. J. Obstet. Gynecol.* 193: 103–113. doi:10.1016/j.ajog.2004.12.025
- Barca, J.A. *et al.*, 2021, Pelvic Floor Morbidity Following Vaginal Delivery versus Cesarean Delivery: Systematic Review and Meta-Analysis<sup>2</sup>, *Journal of Clinical Medicine*, 10(8), p. 1652. Available at: <https://doi.org/10.3390/jcm10081652>.
- Blomquist, J.L., Muñoz, A., Carroll, M., & Handa, V.L., 2018. Association of Delivery Mode With Pelvic Floor Disorders After Childbirth. *JAMA* 21204. doi:10.1001/jama.2018.18315
- Corton, M.M., 2009. Anatomy of Pelvic Floor Dysfunction. *Obstet. Gynecol. Clin. North Am.* 36: 401–419. doi:10.1016/j.ogc.2009.09.002
- Cunningham, F.G., Leveno, K.J., & Bloom, S.L., 2013. *Williams Obstetrics* 23rd Edition-idw.
- Dave, A., Joshi, R., Goel, V., & Dave, A., 2017. Original Research Article MR imaging evaluation for the assessment of pelvic organ prolapse : a newer technique. *Int. J. Reprod. Contraception, Obstet. Gynecol.* 6: 3400–3404. doi:<http://dx.doi.org/10.18203/2320-1770.ijrcog20173451>
- Dietz, H.P., 2004. Ultrasound imaging of the pelvic floor . Part I: two-dimensional aspects. *Ultrasound Obs. Gynecol* 23: 80–92. doi:10.1002/uog.939
- Dietz, H.P., & Shek, K.L., 2009. Levator defects can be detected by 2D translabial ultrasound. *Int. Urogynecol. J.* 20: 807–811. doi:10.1007/s00192-009-0839-4
- Dietz, H.P., 2017. Pelvic floor ultrasound: a review. *Clinical obstetrics and gynecology*, 60(1), pp.58-81.

- Dixit, P., Shek, K.L. and Dietz, H.P., 2014. How common is pelvic floor muscle atrophy after vaginal childbirth?. *Ultrasound in Obstetrics & Gynecology*, 43(1), pp.83-88.
- Dumolin, C., Hay-Smith, E., & Mac Habée-Séguin, G., 2019. Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Res. Nurs. Heal.* 42: 234–235. doi:10.1002/nur.21946
- Gutiñas, A., Galocha, C., Madurga, R., Kirk, J., Usandizaga, R. and Ángel Rodríguez-Zambrano, M., 2022. Application of pelvic floor ultrasound during pregnancy to detect patients at risk of cesarean section due to failure of labor progression in a Spanish population. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 269, pp.102-107.
- Gyhagen, M., Bullarbo, M., Nielsen, T.F., & Milsom, I., 2013. The prevalence of urinary incontinence 20 years after childbirth: A national cohort study in singleton primiparae after vaginal or caesarean delivery. *BJOG An Int. J. Obstet. Gynaecol.* 120: 144–151. doi:10.1111/j.1471-0528.2012.03301.x
- Halle, T.K., Gunvor, J.S., Kari, H., Engh, M.E., & Siafarikas, F., 2020. Change in prevalence of major levator ani muscle defects from 6 weeks to 1 year postpartum , and maternal and obstetric risk factors : A longitudinal ultrasound study. *Acta Obstet. Ginecol. Scand.* 1403–1410. doi:10.1111/aogs.13878
- Jain, P., & Parsons, M., 2011. The effects of obesity on the pelvic floor. *Obstet. Gynaecol.* 13: 133–142. doi:10.1576/toag.13.3.133.27664
- Jelovsek, J.E., Ms, K.C., Gyhagen, M., Hagen, S., Wilson, D., Kattan, M.W., et al., 2018. Predicting risk of pelvic floor disorders 12 and 20 years after delivery. *Am. J. Obstet. Gynecol.* 218: 222.e1-222.e19. doi:10.1016/j.ajog.2017.10.014
- Jelovsek, J.E., Piccorelli, A., Barber, P.M.D., Tunitsky-bitton, E., & Kattan, M.W., 2013. Prediction Models for Postpartum Urinary and Fecal Incontinence in Primiparous Women. *female Pelvic Med. Reconstr. Surg.* 19: 110–118. doi:10.1097/SPV
- Jeon, J., & Na, S., 2017. Vacuum extraction vaginal delivery: Current trend and safety. *Obstet. Gynecol. Sci.* 60: 499–505. doi:10.5468/ogs.2017.60.6.499
- JNPK-KR, POGI, IDAI, IBI, & PPNI, 2008. Buku Acuan Pelatihan Klinik Asuhan Persalinan Normal, Asuhan Esensial, Pencegahan dan Penanggulangan Segera Komplikasi Persalinan dan Bayi Baru Lahir, Revisi 1. *JNPK-KR Depkes RI Jakarta*.
- Jundt, K., Peschers, U., & Kentenich, H., 2015. MEDIZIN: CME: Diagnostik und Therapie der Weiblichen Beckenbodendysfunktion. *Dtsch. Arztebl.*

*Int.* 112. doi:10.3238/arztebl.2015.0564

- Kamel, R. *et al.*, 2019. Contraction of the levator ani muscle during Valsalva maneuver (coactivation) is associated with a longer active second stage of labor in nulliparous women undergoing induction of labor, *American Journal of Obstetrics and Gynecology*, 220(2), p. 189.e1-189.e8. Available at: <https://doi.org/10.1016/j.ajog.2018.10.013>.
- Kementerian Kesehatan, 2018. Data dan Informasi, Profil Kesehatan Indonesia 2017. *Kementeri. Kesehat. Republik Indones. Jakarta.*
- Kementerian Kesehatan, 2016a. Ringkasan Eksekutif Data dan Informasi Kesehatan Provinsi DI Yogyakarta. *Kementeri. Kesehat. Republik Indones. Jakarta.*
- Kementerian Kesehatan, 2016b. Indonesian Case Based Groups (INA-CBG's) dan non Indonesian Case Based. *Peratur. MENTERI Kesehat. REPUBLIK Indones. 52.*
- Kementerian Kesehatan, WHO, IDI, POGI, & IBI, 2013. Buku Saku Pelayanan Kesehatan Ibu di Fasilitas Kesehatan Dsar dan Rujukan, Pedoman bagi Tenaga KesehatanN, 1st ed. *Kementreian Kesehat. Republik Indones. Jakarta.*
- Kochenour, N., Peterson, C., Adashi, C., & Burrows, P., 1997. Human Reproduction: Clinical, Pathologic and Pharmacologic Correlations. *Knowl. Weav. Spencer S. Eccles Heal. Sci. Libr. Univ. Utah.*
- Leeman, L., Rogers, R., Borders, N., Teaf, D., & Qualls, C., 2016. The Effect of Perineal Lacerations on Pelvic Floor Function and Anatomy at 6 Months Postpartum in a Prospective Cohort of Nulliparous Women. *Birth* 43: 293–302. doi:10.1111/birt.12258
- Martínez Franco, E. *et al.*, 2023, Factors involved in changes in the levator ani during pregnancy, *International Urogynecology Journal*, 34(8), pp. 1933–1938. Available at: <https://doi.org/10.1007/s00192-023-05487-4>.
- Masenga, G.G., Shayo, B.C., & Rasch, V., 2018. Prevalence and risk factors for pelvic organ prolapse in Kilimanjaro , Tanzania : A population based study in Tanzanian rural community. *PLoS One* 13: 1–13. doi:<https://doi.org/10.1371/journal.pone.0195910>
- Memon, H.U., & Handa, V.L., 2013. Vaginal childbirth and pelvic floor disorders. *Women's Heal.* 9: 265–277. doi:10.2217/whe.13.17
- Meston, C.M., 2003. Validation of the female sexual function index (Fsfi) in women with female orgasmic disorder and in women with hypoactive sexual desire disorder. *J. Sex Marital Ther.* 29: 39–46. doi:10.1080/713847100
- Pangastuti, N., 2018. Instrumen prediktor disfungsi dasar panggul pasca

persalinan vaginal.

- Pangastuti, N., Sari, D.C.R., Santoso, B.I., Agustiningsih, D., & Emilia, O., 2018. Gambaran Faktor Risiko Prolaps Organ Panggul Pasca Persalinan Vaginal di Daerah Istimewa Yogyakarta Risk Factors for Pelvic Organ Prolapse in Women with History of Vaginal Delivery in Yogyakarta. *Maj. Kedokt. Bandung* 50: 102–108.
- Pregazzi, R., Sartore, A., Bortoli, P., Grimaldi, E., Ricci, G., & Guaschino, S., 2002. Immediate postpartum perineal examination as a predictor of puerperal pelvic floor dysfunction. *Obstet. Gynecol.* 99: 581–584. doi:10.1016/S0029-7844(01)01763-X
- Pudasaini, S. and Dangal, G., 2023, Clinical Profile of Patients of Pelvic Organ Prolapse, *J Nepal Health Res Counc*, 21(58).
- Rahman, M.N. *et al.*, 2023, Obstetrical ballooning and avulsion of levator ani muscle after vaginal delivery: a literature review', *Bali Medical Journal*, 12(2).
- RISKESDAS. 2013. Riset Kesehatan Dasar Badan Penelitian & Pengembangan Kesehatan. Kementrian Kesehatan RI
- Schwertner-Tiepelmann, N., Thakar, R., Sultan, A.H., & Tunn, R., 2012. Obstetric levator ani muscle injuries : current status. *Ultrasound Obs. Gynecol* 39: 372–383. doi:10.1002/uog.11080
- Seo, J.T., & Kim, J.M., 2006. Voiding Dysfunction. *J. Urol.* 175: 1769–1772. doi:10.1016/S0022-5347(05)00993-6
- Serrano, S. *et al.*, 2022, Levator ani muscle avulsion in patients with pelvic floor dysfunction – Does it help in understanding pelvic organ prolapse?, *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 279, pp. 140–145. Available at: <https://doi.org/10.1016/j.ejogrb.2022.09.033>.
- Silva-Lindsey, A., Mojica-Pabellón, J.G. and Goenaga-Jimenez, M.A. (2020) 'Contraction monitor for high risk pregnancies', in L.L. Grewe, E.P. Blasch, and I. Kadar (eds) *Signal Processing, Sensor/Information Fusion, and Target Recognition XXIX. Signal Processing, Sensor/Information Fusion, and Target Recognition XXIX*, Online Only, United States: SPIE, p. 42. Available at: <https://doi.org/10.1117/12.2559153>.
- Steensma, A., 2009. Ultrasound Imaging of the Pelvic Floor: linking anatomical findings with clinical symptoms.
- Van de Waarsenburg, M., Verberne, E., van der Vaart, C. and Withagen, M., 2018. Recovery of puborectalis muscle after vaginal delivery: an ultrasound study. *Ultrasound in Obstetrics & Gynecology*, 52(3), pp.390-395.

WHO, 2008. *WHO policy on non-recruitment of smokers or other tobacco users. World Health Organ.* 6–7.

Wilson, D., Dornan, J., Milsom, I., & Freeman, R., 2014. UR-CHOICE: can we provide mothers-to-be with information about the risk of future pelvic floor dysfunction? *Int. Urogynecol. J.* 25: 1449–1452. doi:10.1007/s00192-014-2376-z

Youssef, A. *et al.*, 2023, The role of the maternal pelvic floor in vaginal delivery, *American Journal of Obstetrics and Gynecology*, p. S0002937823000133. Available at: <https://doi.org/10.1016/j.ajog.2022.12.323>.

Youssef, A., Montaguti, E., Dodaro, M., Kamel, R., Rizzo, N. and Pilu, G., 2019. Levator ani muscle coactivation at term is associated with longer second stage of labor in nulliparous women. *Ultrasound in Obstetrics & Gynecology*, 53(5), pp.686-692.

Zhuang, R.R., Song, Y.F., Chen, Z.Q., Ma, M., Huang, H.J., Chen, J.H., *et al.*, 2011. Levator avulsion using a tomographic ultrasound and magnetic resonance – based model. *Am. J. Obstet. Gynecol.* doi:10.1016/j.ajog.2011.03.052