

## DAFTAR PUSTAKA

- Adrie C, Azoulay E, Francois A, et al. OutcomeRea Study Group. Influence of gender on the outcome of severe sepsis: a reappraisal. *Chest* 2007;132:1786-93. doi:10.1378/chest.07-0420. pmid:17890473.
- American College of Chest Physicians/Society of Critical Care Medicine Consensus Conference: definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med* 1992;20:864-74. doi:10.1097/00003246-199206000-00025 pmid:1597042.
- Angele MK, Pratschke S, Hubbard WJ, Chaudry IH. Gender differences in sepsis: cardiovascular and immunological aspects. *Virulence* 2014;5:12-9. doi:10.4161/viru.26982. pmid:24193307.
- Alobaidi R, Basu RK, Goldstein SL, Bagshaw SM. Sepsis-associated acute kidney injury. *Semin Nephrol* 2015;35:2-11. doi:10.1016/j.semnephrol.2015.01.002. pmid:25795495.
- Beale R, Reinhart K, Brunkhorst FM, et al. PROGRESS Advisory Board. Promoting Global Research Excellence in Severe Sepsis (PROGRESS): lessons from an international sepsis registry. *Infection* 2009;37:222-32. doi:10.1007/s15010-008-8203-z. pmid:19404580.
- Boomer JS, To K, Chang KC, et al. Immunosuppression in patients who die of sepsis and multiple organ failure. *JAMA* 2011;306:2594-605. doi:10.1001/jama.2011.1829. pmid:22187279.
- Bone RC, Sprung CL, Sibbald WJ. Definitions for sepsis and organ failure. *Crit Care Med* 1992;20:724-6. doi:10.1097/00003246-199206000-00002 pmid:1597021.
- Brun-Buisson C, Doyon F, Carlet J, et al. French ICU Group for Severe Sepsis. Incidence, risk factors, and outcome of severe sepsis and septic shock in adults. A multicenter prospective study in intensive care units. *JAMA* 1995;274:968-74. doi:10.1001/jama.1995.03530120060042. pmid:7674528.
- Chandana, Athmananda., Mahendrappa, K., B., Praveen, Unki., Hemanth, Kumar., Madhu, P., Kerudi., K., T., Babu. (2023). Predictive value of CRP and albumin ratio in neonatal sepsis. *International Journal of Contemporary Pediatrics*, doi: 10.18203/2349-3291.ijcp20230431
- Chandra, J., Armengol de la Hoz, M. A., Lee, G., Lee, A., Thorat, P., Elbers, P., Lee, H. C., Munger, J. S., Celi, L. A., & Kaufman, D. A. 2022. A novel Vascular Leak Index identifies sepsis patients with a higher risk for in-hospital death and fluid accumulation. *Critical Care*, 26(1). <https://doi.org/10.1186/s13054-022-03968-4>
- De Backer D, Ortiz JA, Salgado D. Coupling microcirculation to systemic hemodynamics. *Curr Opin Crit Care* 2010;16:250-4. doi:10.1097/MCC.0b013e3283383621. pmid:20179590.
- Deutschman CS, Tracey KJ. Sepsis: current dogma and new perspectives. *Immunity* 2014;40:463-75. doi:10.1016/j.immuni.2014.04.001. pmid:24745331.
- Deretic V, Saitoh T, Akira S. Autophagy in infection, inflammation and immunity. *Nat Rev Immunol* 2013;13:722-37. doi:10.1038/nri3532. pmid:24064518.

- Doig CJ, Sutherland LR, Sandham JD, Fick GH, Verhoef M, Meddings JB. Increased intestinal permeability is associated with the development of multiple organ dysfunction syndrome in critically ill ICU patients. *Am J Respir Crit Care Med* 1998;158:444-51. doi:10.1164/ajrccm.158.2.9710092. pmid:9700119.
- Dombrovskiy VY, Martin AA, Sunderram J, Paz HL. Occurrence and outcomes of sepsis: influence of race. *Crit Care Med* 2007;35:763-8. doi:10.1097/01.CCM.0000256726.80998.BF. pmid:17255870.
- Drewry AM, Samra N, Skrupky LP, Fuller BM, Compton SM, Hotchkiss RS. Persistent lymphopenia after diagnosis of sepsis predicts mortality. *Shock* 2014;42:383-91. doi:10.1097/SHK.0000000000000234. pmid:25051284.
- Ferwerda B, Alonso S, Banahan K, et al. Functional and genetic evidence that the Mal/TIRAP allele variant 180L has been selected by providing protection against septic shock. *Proc Natl Acad Sci U S A* 2009;106:10272-7. doi:10.1073/pnas.0811273106. pmid:19509334.
- Flythe, J. E., Curhan, G. C., & Brunelli, S. M. (2015). Disentangling the ultrafiltration rate-mortality association: The respective roles of session length and weight gain. *Clinical Journal of the American Society of Nephrology*, 8(7), 1151-1161.
- Fullerton JN, O'Brien AJ, Gilroy DW. Pathways mediating resolution of inflammation: when enough is too much. *J Pathol* 2013;231:8-20. doi:10.1002/path.4232. pmid:23794437.
- Fisher CJ Jr, Agosti JM, Opal SM, et al. The Soluble TNF Receptor Sepsis Study Group. Treatment of septic shock with the tumor necrosis factor receptor:Fc fusion protein. *N Engl J Med* 1996;334:1697-702. doi:10.1056/NEJM199606273342603. pmid:8637514.
- Fink MP. Intestinal epithelial hyperpermeability: update on the pathogenesis of gut mucosal barrier dysfunction in critical illness. *Curr Opin Crit Care* 2003;9:143-51. doi:10.1097/00075198-200304000-00011 pmid:12657978.
- Garcia-Alvarez M, Marik P, Bellomo R. Sepsis-associated hyperlactatemia. *Crit Care* 2014;18:503. doi:10.1186/s13054-014-0503-3. pmid:25394679.
- Harrison DA, Welch CA, Eddleston JM. The epidemiology of severe sepsis in England, Wales and Northern Ireland, 1996 to 2004: secondary analysis of a high quality CRP/Albumin database, the ICNARC Case Mix Programme Database. *Crit Care* 2006;10:R42. doi:10.1186/cc4854. pmid:16542492.
- Hartl WH, Jauch K-W. Metabolic self-destruction in critically ill patients: origins, mechanisms and therapeutic principles. *Nutrition* 2014;30:261-7. doi:10.1016/j.nut.2013.07.019. pmid:24369911.
- Hassoun HT, Kone BC, Mercer DW, Moody FG, Weisbrodt NW, Moore FA. Post-injury multiple organ failure: the role of the gut. *Shock* 2001;15:1-10. doi:10.1097/00024382-200115010-00001 pmid:11198350.
- Kashyap, R., Sherani, K. M., Dutt, T., Gnanapandithan, K., Sagar, M., Vallabhajosyula, S., Vakil, A. P., & Surani, S. Current Utility of Sequential Organ Failure Assessment Score: A Literature Review and Future Directions. *The Open Respiratory Medicine Journal*, (2021)15(1), 1-6. <https://doi.org/10.2174/1874306402115010001>.

- Kaukonen K-M, Bailey M, Suzuki S, Pilcher D, Bellomo R. Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand, 2000-2012. *JAMA* 2014;311:1308-16. doi:10.1001/jama.2014.2637. pmid:24638143.
- Lambden, S., Laterre, P. F., Levy, M. M., & Francois, B. The SOFA score-development, utility and challenges of accurate assessment in CRP/Albuminnical trials. *Critical care (London, England)*, (2019) 23(1),374. <https://doi.org/10.1186/s13054-019-2663-7>
- Lee, H. J., Ko, B. S., Ryoo, S. M., Han, E., Suh, G. J., Choi, S. H., Chung, S. P., Lim, T. H., Kim, W. Y., Kwon, W. Y., Hwang, S. Y., Jo, Y. H., Shin, J., Shin, T. G., & Kim, K. Correction: Modified cardiovascular SOFA score in sepsis: development and internal and external validation. *BMC Medicine*,(2021). 20(1). <https://doi.org/10.1186/s12916-022-02694-6>
- Levi M. Pathogenesis and treatment of disseminated intravascular coagulation in the septic patient. *J Crit Care* 2001;16:167-77. doi:10.1053/jcrc.2001.30666. pmid:11815902.
- Levy BD, Serhan CN. Resolution of acute inflammation in the lung. *Annu Rev Physiol* 2014;76:467-92. doi:10.1146/annurev-physiol-021113 170408. pmid:24313723.
- Martin GS, Mannino DM, Eaton S, Moss M. The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 2003;348:1546-54. doi:10.1056/NEJMoa022139. pmid:12700374.
- Martin GS, Mannino DM, Moss M. The effect of age on the development and outcome of adult sepsis. *Crit Care Med* 2006;34:15-21. doi:10.1097/01.CCM.0000194535.82812.BA. pmid:16374151.
- Matthay MA, Ware LB, Zimmerman GA. The acute respiratory distress syndrome. *J CRP/Albuminn Invest* 2012;122:2731-40. doi:10.1172/ JCI60331. pmid:22850883.
- Mayr FB, Yende S, Linde-Zwirble WT, et al. Infection rate and acute organ dysfunction risk as explanations for racial differences in severe sepsis. *JAMA* 2010;303:2495-503. doi:10.1001/ jama.2010.851. pmid:20571016.
- Martinod K, Wagner DD. Thrombosis: tangled up in NETs. *Blood* 2014;123:2768-76. doi:10.1182/blood-2013-10- 463646. pmid:24366358.
- Moine P, Abraham E. Immunomodulation and sepsis: impact of the pathogen. *Shock* 2004;22:297-308. doi:10.1097/01. shk.0000140663.80530.73 pmid:15377883.
- Nehring, S. M. Role of C-reactive Protein as an Indicator for Determining the Outcome of Sepsis. (2019). *Indian Journal of Critical Care Medicine*, 23(1), 11–14. <https://doi.org/10.5005/jp-journals-10071-23105>.
- Obaro SK, Monteil MA, Henderson DC. The pneumococcal problem. *BMJ* 1996;312:1521-5. doi:10.1136/bmj.312.7045.1521 pmid:8646147.
- Otto GP, Sossdorf M, Claus RA, et al. The late phase of sepsis is characterized by an increased microbiological burden and death rate. *Crit Care* 2011;15:R183. doi:10.1186/cc10332. pmid:21798063.

- Parker MM, Shelhamer JH, Bacharach SL, et al. Profound but reversible myocardial depression in patients with septic shock. *Ann Intern Med* 1984;100:483-90. doi:10.7326/0003-4819-100-4-483 pmid:6703540.
- Rabuel C, Mebazaa A. Septic shock: a heart story since the 1960s. *Intensive Care Med* 2006;32:799-807. doi:10.1007/s00134-006-0142- 5. pmid:16570145.
- Rivers E, Nguyen B, Havstad S, et al. Early Goal-Directed Therapy Collaborative Group. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 2001;345:1368-77. doi:10.1056/ NEJMoa010307. pmid:11794169.
- Schramm P, Klein KU, Falkenberg L, et al. Impaired cerebrovascular autoregulation in patients with severe sepsis and sepsis- associated delirium. *Crit Care* 2012;16:R181. doi:10.1186/ cc11665. pmid:23036135.
- Shapiro NI, Howell MD, Talmor D, et al. Serum lactate as a predictor of mortality in emergency department patients with infection. *Ann Emerg Med* 2005;45:524-8. doi:10.1016/j. annemergmed.2004.12.006. pmid:15855951.
- Sopiyudin, D.M. (2013). Besar Sampel dan Cara Pengambilan Sampel. Jakarta : Salemba Medika.
- Park JE, Chung KS, Song JH, et al. The C-Reactive Protein/Albumin Ratio as a Predictor of Mortality in Critically Ill Patients. *Journal of CRP/Albuminnical Medicine* 2018;7:333-42. doi:10.3390/jcm7100333
- Prowle JR, Bellomo R. Sepsis-associated acute kidney injury: macrohemodynamic and microhemodynamic alterations in the renal circulation. *Semin Nephrol* 2015;35:64-74. doi:10.1016/j. semnephrol.2015.01.007. pmid:25795500.
- Ulloa L, Tracey KJ. The “cytokine profile”: a code for sepsis. *Trends Mol Med* 2005;11:56-63. doi:10.1016/j. molmed.2004.12.007.pmid:15694867.
- Takasu O, Gaut JP, Watanabe E, et al. Mechanisms of cardiac and renal dysfunction in patients dying of sepsis. *Am J Respir Crit Care Med* 2013;187:509-17. doi:10.1164/rccm.201211-1983OC. pmid:23348975.
- Tracey KJ, Beutler B, Lowry SF, et al. Shock and tissue injury induced by recombinant human cachectin. *Science* 1986;234:470-4. doi:10.1126/science.3764421 pmid:3764421.
- Ramdeen, S., Ferrell, B., Bonk, C., Schubel, L., Littlejohn, R., Capan, M., Arnold, R., & Miller, K. The Available Criteria for Different Sepsis Scoring Systems in the Emergency Department-A Retrospective Assessment. Open access emergency medicine OAEM (2021)13, 91–96. <https://doi.org/10.2147>
- Schroder K, Tschopp J. The inflammasomes. *Cell* 2010;140:821-32. doi:10.1016/j.cell.2010.01.040. pmid:20303873.
- Singer M. The role of mitochondrial dysfunction in sepsis-induced multi-organ failure. *Virulence* 2014;5:66-72. doi:10.4161/ viru.26907. pmid:24185508
- Vincent J-L, Opal SM, Marshall JC, Tracey KJ. Sepsis definitions: time for change. *Lancet* 2013;381:774-5. doi:10.1016/S0140- 6736(12)61815-7. pmid:23472921.
- Walley KR, Francis GA, Opal SM, et al. The central role of proprotein convertase subtilisin/kexin type 9 in septic pathogen lipid transport and clearance. *Am J Respir Crit Care Med* 2015;192:1275-86. doi:10.1164/ rccm.201505-0876CI. pmid:26252194.

- Wan L, Bagshaw SM, Langenberg C, Saotome T, May C, Bellomo R. Pathophysiology of septic acute kidney injury: what do we really know? *Crit Care Med* 2008;36(Suppl):S198-203. doi:10.1097/CCM.0b013e318168ccd5. pmid:18382194.
- Wang, B., Chen, G., Cao, Y., Xue, J., Li, J., & Wu, Y. 2015. Correlation of lactate/albumin ratio level to organ failure and mortality in severe sepsis and septic shock. *Journal of Critical Care*, 30(2), 271–275. <https://doi.org/10.1016/j.jcrc.2014.10.030>
- Wolinsky H. A proposal linking clearance of circulating lipoproteins to tissue metabolic activity as a basis for understanding atherogenesis. *Circ Res* 1980;47:301-11. doi:10.1161/01.RES.47.3.301 pmid:6996862.
- Yin, M., Si, L., Qin, W., Li, C., Zhang, J., Yang, H., Han, H., Zhang, F., Ding, S., Zhou, M., Wu, D., Chen, X., & Wang, H. 2016. Predictive Value of Serum Albumin Level for the Prognosis of Severe Sepsis Without Exogenous Human Albumin Administration: A Prospective Cohort Study. *Journal of Intensive Care Medicine*, 33(12), 687–694. <https://doi.org/10.1177/0885066616685300>.
- Yongkai, Li., Dandan, Li., Xin, Yuan., Haireti, Nazila., Liu, Yang., Ran, Xu., Xiao-Cheng, Liu., Xin, Li., Shu-Min, Jiang., Saimaiti, Xialaibaitigu., Jianzhong, Yang. (2023). [Predictive value of early lactate/albumin ratio in the prognosis of sepsis].. Chinese critical care medicine, doi: 10.3760/cma.j.cn121430-20220130-00108