



DAFTAR PUSTAKA

- Al-Mufti, F. *et al.* (2018) ‘Clinical and radiographic predictors of intracerebral hemorrhage outcome’, *Interventional Neurology*, 7(1–2), pp. 118–136. doi:10.1159/000484571.
- Alba, A.C. *et al.* (2017) ‘Discrimination and calibration of clinical prediction models’, *JAMA*, 318(14), p. 1377. doi:10.1001/jama.2017.12126.
- Alimohammadi, E. *et al.* (2022) ‘Baseline Neutrophil–lymphocyte ratio can be associated with hematoma expansion in patients with intracerebral hemorrhage: A retrospective observational study’, *BMC Neuroscience*, 23(1). doi:10.1186/s12868-022-00705-z.
- An, S.J., Kim, T.J. and Yoon, B.-W. (2017) ‘Epidemiology, risk factors, and clinical features of intracerebral hemorrhage: An update’, *Journal of Stroke*, 19(1), pp. 3–10. doi:10.5853/jos.2016.00864.
- Awad, I.A. *et al.* (2019) ‘Surgical performance determines functional outcome benefit in the minimally invasive surgery plus recombinant tissue plasminogen activator for intracerebral hemorrhage evacuation (MISTIE) procedure’, *Neurosurgery*, 84(6), pp. 1157–1168. doi:10.1093/neuros/nyz077.
- Badan Penelitian Dan Pengembangan Kesehatan, Departemen Kesehatan Republik Indonesia. (2008) Riset Kesehatan Dasar.
- Beckman, J.A. and Creager, M.A. (2016) ‘Vascular complications of diabetes’, *Circulation Research*, 118(11), pp. 1771–1785. doi:10.1161/circresaha.115.306884.
- Bhattathiri, P.S. *et al.* (2006) ‘Intraventricular hemorrhage and hydrocephalus after spontaneous intracerebral hemorrhage: Results from the stSICH trial’, *Brain Edema XIII*, pp. 65–68. doi:10.1007/3-211-30714-1_16.
- Broderick, J.P. *et al.* (1993) ‘Volume of intracerebral hemorrhage. A powerful and easy-to-use predictor of 30-day mortality.’, *Stroke*, 24(7), pp. 987–993. doi:10.1161/01.str.24.7.987.
- Buonacera, A. *et al.* (2022) ‘Neutrophil to lymphocyte ratio: An emerging marker of the relationships between the immune system and diseases’, *International Journal of Molecular Sciences*, 23(7), p. 3636. doi:10.3390/ijms23073636
- Chan, E. *et al.* (2015) ‘Significance of intraventricular hemorrhage in acute intracerebral hemorrhage’, *Stroke*, 46(3), pp. 653–658. doi:10.1161/strokeaha.114.008470.



Chen, C.-J. *et al.* (2020) ‘Intracranial pressure monitoring in patients with spontaneous intracerebral hemorrhage’, *Journal of Neurosurgery*, 132(6), pp. 1854–1864. doi:10.3171/2019.3.jns19545.

Chen, W. *et al.* (2019) ‘The predictive role of postoperative neutrophil to lymphocyte ratio for 30-day mortality after intracerebral hematoma evacuation’, *World Neurosurgery*, 134. doi:10.1016/j.wneu.2019.10.154.

Cheung, R.T. and Zou, L.-Y. (2003) ‘Use of the original, modified, or new intracerebral hemorrhage score to predict mortality and morbidity after intracerebral hemorrhage’, *Stroke*, 34(7), pp. 1717–1722. doi:10.1161/01.str.0000078657.22835.b9.

Clarke, J.L. *et al.* (2004) ‘External validation of the SICH score’, *Neurocritical Care*, 1(1), pp. 53–60. doi:10.1385/ncc:1:1:53.

Dahlan, M.S. (2010) *Besar Sampel dan Cara Pengambilan Sampel*. 3rd edn. Jakarta: Salemba Medika.

de Oliveira Manoel, A.L. (2020) ‘Surgery for spontaneous intracerebral hemorrhage’, *Critical Care*, 24(1). doi:10.1186/s13054-020-2749-2.

Derex, L., Rheims, S. and Peter-Derex, L. (2021) ‘Seizures and epilepsy after intracerebral hemorrhage: An update’, *Journal of Neurology*, 268(7), pp. 2605–2615. doi:10.1007/s00415-021-10439-3.

Dicpinigaitis, A.J. *et al.* (2023) ‘Increase in ruptured cerebral arteriovenous malformations and mortality in the United States: Unintended consequences of the Aruba Trial?’, *Stroke: Vascular and Interventional Neurology*, 3(1). doi:10.1161/svin.122.000442.

Dowlatzahi, D. *et al.* (2011) ‘Defining hematoma expansion in intracerebral hemorrhage: Relationship with patient outcomes’, *Neurology*, 76(14), pp. 1238–1244. doi:10.1212/wnl.0b013e3182143317.

Fakiri, M.O. *et al.* (2020) ‘Reliability of the intracerebral hemorrhage score for predicting outcome in patients with intracerebral hemorrhage using oral anticoagulants’, *European Journal of Neurology*, 27(10), pp. 2006–2013. doi:10.1111/ene.14336.

Fang, H. *et al.* (2013) ‘Toll-like receptor 4 signaling in intracerebral hemorrhage-induced inflammation and injury’, *Journal of Neuroinflammation*, 10(1). doi:10.1186/1742-2094-10-27.

Feigin, V.L. *et al.* (2021) ‘Global, regional, and national burden of stroke and its risk factors, 1990–2019: A systematic analysis for the global burden of



disease study 2019', *The Lancet Neurology*, 20(10), pp. 795–820. doi:10.1016/s1474-4422(21)00252-0.

Fiebach, J.B. et al. (2004) 'Stroke magnetic resonance imaging is accurate in hyperacute intracerebral hemorrhage', *Stroke*, 35(2), pp. 502–506. doi:10.1161/01.str.0000114203.75678.88.

Forget, P. et al. (2017) 'What is the normal value of the neutrophil-to-lymphocyte ratio?', *BMC Research Notes*, 10(1). doi:10.1186/s13104-016-2335-5.

Gautier, S. et al. (2009) 'Neutrophils contribute to intracerebral haemorrhages after treatment with recombinant tissue plasminogen activator following cerebral ischaemia', *British Journal of Pharmacology*, 156(4), pp. 673–679. doi:10.1111/j.1476-5381.2009.00068.x.

Greenberg, S.M. and Charidimou, A. (2018) 'Diagnosis of cerebral amyloid angiopathy', *Stroke*, 49(2), pp. 491–497. doi:10.1161/strokeaha.117.016990.

Gregson, B.A. et al. (2012) 'Individual patient data subgroup meta-analysis of surgery for spontaneous supratentorial intracerebral hemorrhage', *Stroke*, 43(6), pp. 1496–1504. doi:10.1161/strokeaha.111.640284.

Gross, B.A. and Du, R. (2017) 'Hemorrhage from cerebral cavernous malformations: A systematic pooled analysis', *Journal of Neurosurgery*, 126(4), pp. 1079–1087. doi:10.3171/2016.3.jns152419.

Hallan, D.R., Simion, C. and Rizk, E. (2022) 'Early versus late tracheostomy in spontaneous intracerebral hemorrhage', *Cureus* [Preprint]. doi:10.7759/cureus.24059.

Hanley, D.F. et al. (2017) 'Thrombolytic removal of intraventricular haemorrhage in treatment of severe stroke: Results of the randomised, multicentre, multiregion, placebo-controlled clear III trial', *The Lancet*, 389(10069), pp. 603–611. doi:10.1016/s0140-6736(16)32410-2.

Hanley, D.F. et al. (2019) 'Efficacy and safety of minimally invasive surgery with thrombolysis in intracerebral haemorrhage evacuation (Mistie III): A randomised, controlled, open-label, Blinded Endpoint Phase 3 trial', *The Lancet*, 393(10175), pp. 1021–1032. doi:10.1016/s0140-6736(19)30195-3.

Hemphill, J.C. et al. (2001) 'The SICH score', *Stroke*, 32(4), pp. 891–897. doi:10.1161/01.str.32.4.891.

Herrick, D.B. et al. (2014) 'Determinants of external ventricular drain placement and associated outcomes in patients with spontaneous intraventricular



hemorrhage', *Neurocritical Care*, 21(3), pp. 426–434. doi:10.1007/s12028-014-9959-x.

Hua, Y. et al. (2007) 'Brain injury after intracerebral hemorrhage', *Stroke*, 38(2), pp. 759–762. doi:10.1161/01.str.0000247868.97078.10.

Hwang, B.Y. et al. (2012) 'Evaluation of intraventricular hemorrhage assessment methods for predicting outcome following intracerebral hemorrhage', *Journal of Neurosurgery*, 116(1), pp. 185–192. doi:10.3171/2011.9.jns10850.

Ironside, N. et al. (2019) 'Perihematomal edema after spontaneous intracerebral hemorrhage', *Stroke*, 50(6), pp. 1626–1633. doi:10.1161/strokeaha.119.024965.

Ji, R. et al. (2014) 'Risk score to predict hospital-acquired pneumonia after spontaneous intracerebral hemorrhage', *Stroke*, 45(9), pp. 2620–2628. doi:10.1161/strokeaha.114.005023.

Karimy, J.K., Reeves, B.C. and Kahle, K.T. (2020) 'Targeting TLR4-dependent inflammation in post-hemorrhagic brain injury', *Expert Opinion on Therapeutic Targets*, 24(6), pp. 525–533. doi:10.1080/14728222.2020.1752182.

Keep, R.F., Hua, Y. and Xi, G. (2012) 'Intracerebral Haemorrhage: Mechanisms of injury and therapeutic targets', *The Lancet Neurology*, 11(8), pp. 720–731. doi:10.1016/s1474-4422(12)70104-7.

Kidwell, C.S. (2004) 'Comparison of MRI and CT for detection of acute intracerebral hemorrhage', *JAMA*, 292(15), p. 1823. doi:10.1001/jama.292.15.1823.

Koivunen, R.-J. et al. (2014) 'Predictors of early mortality in young adults after intracerebral hemorrhage', *Stroke*, 45(8), pp. 2454–2456. doi:10.1161/strokeaha.114.006020.

Lattanzi, S. et al. (2018) 'Neutrophil-to-lymphocyte ratio improves outcome prediction of acute intracerebral hemorrhage', *Journal of the Neurological Sciences*, 387, pp. 98–102. doi:10.1016/j.jns.2018.01.038.

Levine, J.M. et al. (2007) 'Early edema in warfarin-related intracerebral hemorrhage', *Neurocritical Care*, 7(1), pp. 58–63. doi:10.1007/s12028-007-0039-3.



Li, Q. *et al.* (2020) ‘Ultra-early blood pressure reduction attenuates hematoma growth and improves outcome in intracerebral hemorrhage’, *Annals of Neurology*, 88(2), pp. 388–395. doi:10.1002/ana.25793.

Linn, J. *et al.* (2010) ‘Prevalence of superficial siderosis in patients with cerebral amyloid angiopathy’, *Neurology*, 74(17), pp. 1346–1350. doi:10.1212/wnl.0b013e3181dad605.

Liu, S. *et al.* (2019) ‘Neutrophil–lymphocyte ratio predicts the outcome of intracerebral hemorrhage’, *Journal of Medicine*, 98(26). doi:10.1097/md.00000000000016211.

Lord, A.S. *et al.* (2014) ‘Infection after intracerebral hemorrhage’, *Stroke*, 45(12), pp. 3535–3542. doi:10.1161/strokeaha.114.006435.

Luo, S. *et al.* (2022) ‘The clinical value of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, and D-dimer-to-fibrinogen ratio for predicting pneumonia and poor outcomes in patients with acute intracerebral hemorrhage’, *Frontiers in Immunology*, 13. doi:10.3389/fimmu.2022.1037255.

Luzzi, S. *et al.* (2019) ‘Indication, timing, and surgical treatment of spontaneous intracerebral hemorrhage: Systematic Review and proposal of a management algorithm’, *World Neurosurgery*, 124. doi:10.1016/j.wneu.2019.01.016.

Magid-Bernstein, J. *et al.* (2022) ‘Cerebral hemorrhage: Pathophysiology, treatment, and future directions’, *Circulation Research*, 130(8), pp. 1204–1229. doi:10.1161/circresaha.121.319949.

Manning, L. *et al.* (2014) ‘Blood pressure variability and outcome after acute intracerebral haemorrhage: A post-hoc analysis of Interact2, a randomised controlled trial’, *The Lancet Neurology*, 13(4), pp. 364–373. doi:10.1016/s1474-4422(14)70018-3.

Meisel, C. *et al.* (2005) ‘Central nervous system injury-induced immune deficiency syndrome’, *Nature Reviews Neuroscience*, 6(10), pp. 775–786. doi:10.1038/nrn1765.

Mendelow, A.D. *et al.* (2013) ‘Early surgery versus initial conservative treatment in patients with spontaneous supratentorial lobar intracerebral haematomas (STICH II): A randomised trial’, *The Lancet*, 382(9890), pp. 397–408. doi:10.1016/s0140-6736(13)60986-1.

Mendelow, D. *et al.* (2005) ‘Early surgery versus initial conservative treatment in patients with spontaneous supratentorial intracerebral haematomas in the International Surgical Trial in Intracerebral Haemorrhage (STSICH): a



randomised trial', *Lancet*, 365(9457), pp. 387–397.
doi:[https://doi.org/10.1016/S0140-6736\(05\)17826-X](https://doi.org/10.1016/S0140-6736(05)17826-X).

Messé, S.R. et al. (2009) 'Prophylactic antiepileptic drug use is associated with poor outcome following SICH', *Neurocritical Care*, 11(1), pp. 38–44.
doi:[10.1007/s12028-009-9207-y](https://doi.org/10.1007/s12028-009-9207-y).

Miyagi, T. et al. (2015) 'Reduced estimated glomerular filtration rate affects outcomes 3 months after intracerebral hemorrhage: The Stroke Acute Management with urgent risk-factor assessment and improvement-intracerebral hemorrhage study', *Journal of Stroke and Cerebrovascular Diseases*, 24(1), pp. 176–182.
doi:[10.1016/j.jstrokecerebrovasdis.2014.08.015](https://doi.org/10.1016/j.jstrokecerebrovasdis.2014.08.015).

Moxon-Emre, I. and Schlichter, L.C. (2011) 'Neutrophil depletion reduces blood-brain barrier breakdown, axon injury, and inflammation after intracerebral hemorrhage', *Journal of Neuropathology & Experimental Neurology*, 70(3), pp. 218–235. doi:[10.1097/nen.0b013e31820d94a5](https://doi.org/10.1097/nen.0b013e31820d94a5).

Murthy, S.B. et al. (2017) 'Permanent CSF shunting after intraventricular hemorrhage in the clear III trial', *Neurology*, 89(4), pp. 355–362.
doi:[10.1212/wnl.0000000000004155](https://doi.org/10.1212/wnl.0000000000004155).

Nag, C. et al. (2012) 'Prediction of clinical outcome in acute hemorrhagic stroke from a single CT Scan on admission', *North American Journal of Medical Sciences*, 4(10), p. 463. doi:[10.4103/1947-2714.101986](https://doi.org/10.4103/1947-2714.101986).

Nagatsuna, T. et al. (2005) 'Systemic administration of Argatroban reduces secondary brain damage in a rat model of intracerebral hemorrhage: Histopathological assessment', *Cerebrovascular Diseases*, 19(3), pp. 192–200. doi:[10.1159/000083466](https://doi.org/10.1159/000083466).

Park, Y.K. et al. (2012) 'The relationship between metabolic syndrome (METS) and spontaneous intracerebral hemorrhage (SICH)', *Neurological Sciences*, 34(9), pp. 1523–1528. doi:[10.1007/s10072-012-1272-x](https://doi.org/10.1007/s10072-012-1272-x).

Pollock, B.E. et al. (1996) 'Factors that predict the bleeding risk of cerebral arteriovenous malformations', *Stroke*, 27(1), pp. 1–6.
doi:[10.1161/01.str.27.1.1](https://doi.org/10.1161/01.str.27.1.1).

Poon, M.T.C., Bell, S.M. and Al-Shahi Salman, R. (2015) 'Epidemiology of Intracerebral Haemorrhage', *New Insights in Intracerebral Hemorrhage*, pp. 1–12. doi:[10.1159/000437109](https://doi.org/10.1159/000437109).



Qureshi AI, Mendelow AD, Hanley DF. Intracerebral haemorrhage. *The Lancet* [Internet]. 2009 May 1;373(9675):1632–44. Available from: [https://doi.org/10.1016/s0140-6736\(09\)60371-8](https://doi.org/10.1016/s0140-6736(09)60371-8)

Radu, R.A. et al. (2021) ‘Neutrophil-to-lymphocyte ratio as an independent predictor of in-hospital mortality in patients with acute intracerebral hemorrhage’, *Medicina*, 57(6), p. 622. doi:10.3390/medicina57060622.

Roh, D.J. et al. (2019) ‘Low hemoglobin and hematoma expansion after intracerebral hemorrhage’, *Neurology*, 93(4). doi:10.1212/wnl.0000000000007820.

Rutledge, W.C. et al. (2014) ‘Hemorrhage rates and risk factors in the natural history course of brain arteriovenous malformations’, *Translational Stroke Research*, 5(5), pp. 538–542. doi:10.1007/s12975-014-0351-0.

Sacco, R.L. et al. (2013) ‘An updated definition of stroke for the 21st Century’, *Stroke*, 44(7), pp. 2064–2089. doi:10.1161/str.0b013e318296aeca.

Safatli, D. et al. (2016) ‘Predictors of 30-day mortality in patients with spontaneous primary intracerebral hemorrhage’, *Surgical Neurology International*, 7(19), p. 510. doi:10.4103/2152-7806.187493.

Salaun, E. et al. (2018) ‘Intracranial haemorrhage in infective endocarditis’, *Archives of Cardiovascular Diseases*, 111(12), pp. 712–721. doi:10.1016/j.acvd.2018.03.009.

Samarasekera, N. et al. (2015) ‘Influence of intracerebral hemorrhage location on incidence, characteristics, and outcome’, *Stroke*, 46(2), pp. 361–368. doi:10.1161/strokeaha.114.007953.

Sastroasmoro, S., Ismael, S. and Madiyono, B. (2014) ‘Perkiraan Besar Sampel’, in *Dasar-Dasar Metodologi Penelitian Klinis*. 5th edn. Jakarta, DKI Jakarta: Sagung Seto, pp. 352–387.

Saxena, A. et al. (2016) ‘Prognostic significance of hyperglycemia in acute intracerebral hemorrhage’, *Stroke*, 47(3), pp. 682–688. doi:10.1161/strokeaha.115.011627.

Schlunk, F. and Greenberg, S.M. (2015) ‘The pathophysiology of intracerebral hemorrhage formation and expansion’, *Translational Stroke Research*, 6(4), pp. 257–263. doi:10.1007/s12975-015-0410-1.

Seiffge, D.J. et al. (2019) ‘Meta-analysis of haematoma volume, haematoma expansion and mortality in intracerebral haemorrhage associated with oral



anticoagulant use', *Journal of Neurology*, 266(12), pp. 3126–3135.
doi:10.1007/s00415-019-09536-1.

Shebl, E. and Gulick, P.G. (2023) *Nosocomial pneumonia - statpearls - NCBI bookshelf*. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK535441/> (Accessed: 04 December 2023).

Shi, M. et al. (2022) 'Prognostic role of the Neutrophil-to-lymphocyte ratio in intracerebral hemorrhage: A systematic review and meta-analysis', *Frontiers in Neuroscience*, 16. doi:10.3389/fnins.2022.825859.

Shinohara, Mitsuru et al. (2016) 'Impact of sex and APOE4 on cerebral amyloid angiopathy in alzheimer's disease', *Acta Neuropathologica*, 132(2), pp. 225–234. doi:10.1007/s00401-016-1580-y.

Shoamanesh, A. and Selim, M. (2022) 'Use of lipid-lowering drugs after intracerebral hemorrhage', *Stroke*, 53(7), pp. 2161–2170. doi:10.1161/strokeaha.122.036889.

Song, M. et al. (2021) 'Neutrophil-to-lymphocyte ratio and mortality in the United States general population', *Scientific Reports*, 11(1). doi:10.1038/s41598-020-79431-7.

Sun, S. et al. (2019) 'The effect of mannitol in the early stage of supratentorial hypertensive intracerebral hemorrhage: A systematic review and meta-analysis', *World Neurosurgery*, 124, pp. 386–396. doi:10.1016/j.wneu.2018.11.249.

Sutherland, G.R. and Auer, R.N. (2006) 'Primary intracerebral hemorrhage', *Journal of Clinical Neuroscience*, 13(5), pp. 511–517. doi:10.1016/j.jocn.2004.12.012.

Tran, Q.K. et al. (2021) 'Preventing seizure occurrence following spontaneous intracerebral haemorrhage: A systematic review and meta-analysis of seizure prophylaxis', *Seizure*, 87, pp. 46–55. doi:10.1016/j.seizure.2021.02.029.

U.M., P. et al. (2019) 'Validation of SICH and SICH-GS scores in an Indian cohort: Impact of medical and Surgical Management', *Journal of Stroke and Cerebrovascular Diseases*, 28(8), pp. 2213–2220. doi:10.1016/j.jstrokecerebrovasdis.2019.05.003.

Viswanathan, A. and Greenberg, S.M. (2011) 'Cerebral amyloid angiopathy in the elderly', *Annals of Neurology*, 70(6), pp. 871–880. doi:10.1002/ana.22516.



- Wang, J. and Doré, S. (2006) ‘Inflammation after intracerebral hemorrhage’, *Journal of Cerebral Blood Flow & Metabolism*, 27(5), pp. 894–908. doi:10.1038/sj.jcbfm.9600403.
- Wei, J. et al. (2020) ‘Perihematomal brain tissue iron concentration measurement by MRI in patients with intracerebral hemorrhage’, *CNS Neuroscience & Therapeutics*, 26(9), pp. 896–901. doi:10.1111/cns.13395.
- Yang, H. et al. (2021) ‘A retrospective study on risk factors and disease burden for hospital-acquired pneumonia caused by multi-drug-resistant bacteria in patients with intracranial cerebral hemorrhage’, *Neurological Sciences*, 43(4), pp. 2461–2467. doi:10.1007/s10072-021-05721-1.
- Zacharia, B.E. et al. (2012) ‘Predictors of long-term shunt-dependent hydrocephalus in patients with intracerebral hemorrhage requiring emergency cerebrospinal fluid diversion’, *Neurosurgical Focus*, 32(4). doi:10.3171/2012.2.focus11372.
- Zahuranec, D.B. et al. (2014) ‘Intracerebral hemorrhage mortality is not changing despite declining incidence’, *Neurology*, 82(24), pp. 2180–2186. doi:10.1212/wnl.0000000000000519.
- Zhao, X. et al. (2009) ‘Neuroprotective role of haptoglobin after intracerebral hemorrhage’, *The Journal of Neuroscience*, 29(50), pp. 15819–15827. doi:10.1523/jneurosci.3776-09.2009.
- Zhao, Y. et al. (2022) ‘The predictive value of neutrophil to lymphocyte ratio on 30-day outcomes in spontaneous intracerebral hemorrhage patients after surgical treatment: A retrospective analysis of 128 patients’, *Frontiers in Neurology*, 13. doi:10.3389/fneur.2022.963397.