

DAFTAR PUSTAKA

- Altun, I. and Yüksel, K.Z. (2017) ‘Lumbar herniated disc: spontaneous regression’, *The Korean Journal of Pain*, 30(1), p. 44. Available at: <https://doi.org/10.3344/KJP.2017.30.1.44>.
- Aoki, Y. *et al.* (2012) ‘Evaluation of Nonspecific Low Back Pain Using a New Detailed Visual Analogue Scale for Patients in Motion, Standing, and Sitting: Characterizing Nonspecific Low Back Pain in Elderly Patients’, *Pain Research and Treatment*, 2012. Available at: <https://doi.org/10.1155/2012/680496>.
- Betteridge, C., Mobbs, R.J. and Ho, D. (2021) ‘Proposed objective scoring algorithm for walking performance, based on relevant gait metrics: the Simplified Mobility Score (SMoS™)-observational study’, *Journal of orthopaedic surgery and research*, 16(1), p. 419. Available at: <https://doi.org/10.1186/S13018-021-02546-8>.
- Bienstock, D.M. *et al.* (2022) ‘Accelerometry Data Delineate Phases of Recovery and Supplement Patient-Reported Outcome Measures Following Lumbar Laminectomy’, *World neurosurgery*, 160, pp. e608–e615. Available at: <https://doi.org/10.1016/J.WNEU.2022.01.097>.
- Cao, J. *et al.* (2019) ‘Percutaneous endoscopic lumbar discectomy for lumbar disc herniation as day surgery – short-term clinical results of 235 consecutive cases’, *Medicine*, 98(49). Available at: <https://doi.org/10.1097/MD.00000000000018064>.
- Chiu, C.C. *et al.* (2015) ‘The probability of spontaneous regression of lumbar herniated disc: a systematic review’, *Clinical rehabilitation*, 29(2), pp. 184–195. Available at: <https://doi.org/10.1177/0269215514540919>.
- Cicco, F.L. De and Willhuber, G.O.C. (2023) ‘Nucleus Pulposus Herniation’, *StatPearls* [Preprint]. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK542307/> (Accessed: 8 September 2024).
- Dydyk, A.M., Massa, R.N. and Mesfin, F.B. (2023) ‘Disc Herniation’, *StatPearls* [Preprint]. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK441822/> (Accessed: 8 September 2024).
- Ehrler, F., Weber, C. and Lovis, C. (2016) ‘Influence of Pedometer Position on Pedometer Accuracy at Various Walking Speeds: A Comparative Study’, *Journal of medical Internet research*, 18(10). Available at: <https://doi.org/10.2196/JMIR.5916>.
- Eun, S.S., Chachan, S. and Lee, S.H. (2018) ‘Interlaminar Percutaneous Endoscopic Lumbar Discectomy: Rotate and Retract Technique’, *World*

- Neurosurgery*, 118, pp. 188–192. Available at: <https://doi.org/10.1016/J.WNEU.2018.07.083>.
- Frost, B.A., Camarero-Espinosa, S. and Johan Foster, E. (2019) ‘Materials for the Spine: Anatomy, Problems, and Solutions’, *Materials*, 12(2). Available at: <https://doi.org/10.3390/MA12020253>.
- Gebhart, G.F. *et al.* (2013) *Encyclopedia of pain, vol. 1-7, Encyclopedia of Pain*. Edited by G.F. Gebhart and R.F. Schmidt. Berlin, Heidelberg: Springer Publishing Company. Available at: <https://vbn.aau.dk/en/publications/encyclopedia-of-pain-vol-1-7> (Accessed: 8 September 2024).
- Gilmore, S.J. *et al.* (2019) ‘Predictors of substantial improvement in physical function six months after lumbar surgery: is early post-operative walking important? A prospective cohort study’, *BMC musculoskeletal disorders*, 20(1). Available at: <https://doi.org/10.1186/S12891-019-2806-7>.
- Hashimoto, Y. *et al.* (2018) ‘Objectively Measured Physical Activity and Low Back Pain in Japanese Men’, *Journal of physical activity & health*, 15(6), pp. 417–422. Available at: <https://doi.org/10.1123/JPAH.2017-0085>.
- Haugen, A.J. *et al.* (2011) ‘Estimates of success in patients with sciatica due to lumbar disc herniation depend upon outcome measure’, *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*, 20(10), pp. 1669–1675. Available at: <https://doi.org/10.1007/S00586-011-1809-3>.
- Hirachan, M.P. *et al.* (2017) ‘Clinical Outcome of Percutaneous Endoscopic Lumbar Surgery (PELS) in Treatment of Lumbar Disc Herniation’, *Open Journal of Orthopedics*, 7(4), pp. 99–109. Available at: <https://doi.org/10.4236/OJO.2017.74012>.
- Inoue, Y. *et al.* (2003) ‘Is laparoscopic colorectal surgery less invasive than classical open surgery? Quantitation of physical activity using an accelerometer to assess postoperative convalescence’, *Surgical endoscopy*, 17(8), pp. 1269–1273. Available at: <https://doi.org/10.1007/S00464-002-8876-9>.
- Kambin, P. and Bracer, M.D. (1987) ‘Percutaneous posterolateral discectomy. Anatomy and mechanism.’, *Clinical Orthopaedics and Related Research*, NA;(223), p. 145–154. Available at: <https://doi.org/10.1097/00003086-198710000-00016>.
- Kambin, P. and Gellman, H. (1983) ‘Percutaneous Lateral Discectomy of the Lumbar Spine: A Preliminary Report’, *Clinical Orthopaedics and Related Research*, 174(NA;), p. 113–121. Available at: <https://doi.org/10.1097/00003086-198304000-00015>.

- Kim, D.H. *et al.* (2019) ‘The Usefulness of a Wearable Device in Daily Physical Activity Monitoring for the Hospitalized Patients Undergoing Lumbar Surgery’, *Journal of Korean Neurosurgical Society*, 62(5), p. 561. Available at: <https://doi.org/10.3340/JKNS.2018.0131>.
- Knight, S.R. *et al.* (2021) ‘Mobile devices and wearable technology for measuring patient outcomes after surgery: a systematic review’, *NPJ digital medicine*, 4(1). Available at: <https://doi.org/10.1038/S41746-021-00525-1>.
- Lee, M.K. *et al.* (2023) ‘Nuts and Bolts of Patient-Reported Outcomes in Orthopaedics’, *The Journal of arthroplasty*, 38(4), pp. 616–621. Available at: <https://doi.org/10.1016/J.ARTH.2022.11.011>.
- Lee, S.-H. *et al.* (2013) ‘Percutaneous Endoscopic Lumbar Discectomy for L5–S1 Disc Herniation: Transforaminal versus Interlaminar Approach’, *Pain Physician Journal*, 16(6), pp. 547–556.
- Mark Erwin, W. *et al.* (2014) ‘The cellular and molecular biology of the intervertebral disc: A clinician’s primer’, *The Journal of the Canadian Chiropractic Association*, 58(3), p. 246. Available at: </pmc/articles/PMC4139773/> (Accessed: 8 September 2024).
- Master, H. *et al.* (2021) ‘How Many Steps Per Day During the Early Postoperative Period are Associated With Patient-Reported Outcomes of Disability, Pain, and Opioid Use After Lumbar Spine Surgery?’, *Archives of physical medicine and rehabilitation*, 102(10), pp. 1873–1879. Available at: <https://doi.org/10.1016/J.APMR.2021.06.002>.
- Le Masurier, G.C. and Tudor-locke, C. (2003) ‘Comparison of pedometer and accelerometer accuracy under controlled conditions’, *Medicine and science in sports and exercise*, 35(5), pp. 867–871. Available at: <https://doi.org/10.1249/01.MSS.0000064996.63632.10>.
- Mobbs, R.J., Mobbs, R.R. and Choy, W.J. (2019) ‘Proposed objective scoring algorithm for assessment and intervention recovery following surgery for lumbar spinal stenosis based on relevant gait metrics from wearable devices: the Gait Posture index (GPi)’, *Journal of spine surgery (Hong Kong)*, 5(3), pp. 300–309. Available at: <https://doi.org/10.21037/JSS.2019.09.06>.
- Monticone, M. *et al.* (2012) ‘Responsiveness of the Oswestry Disability Index and the Roland Morris Disability Questionnaire in Italian subjects with sub-acute and chronic low back pain’, *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*, 21(1), pp. 122–129. Available at: <https://doi.org/10.1007/S00586-011-1959-3>.
- Morino, S. *et al.* (2016) ‘The relationship between the daily step counts and low back pain during pregnancy’, *Clinical and Experimental Obstetrics and*

- Gynecology*, 43(2), pp. 192–197. Available at: <https://doi.org/10.12891/CEOG2078.2016/PDF>.
- Natarajan, P. *et al.* (2023) ‘Continuous data capture of gait and mobility metrics using wearable devices for postoperative monitoring in common elective orthopaedic procedures of the hip, knee, and spine: a scoping review’, *Journal of orthopaedic surgery and research*, 18(1). Available at: <https://doi.org/10.1186/S13018-023-04303-5>.
- Pan, M. *et al.* (2020) ‘Percutaneous Endoscopic Lumbar Discectomy: Indications and Complications’, *Pain Physician Journal*, 23(1), pp. 49–56.
- Qin, R. *et al.* (2018) ‘Percutaneous Endoscopic Lumbar Discectomy Versus Posterior Open Lumbar Microdiscectomy for the Treatment of Symptomatic Lumbar Disc Herniation: A Systemic Review and Meta-Analysis’, *World neurosurgery*, 120, pp. 352–362. Available at: <https://doi.org/10.1016/J.WNEU.2018.08.236>.
- Ruan, W. *et al.* (2016) ‘Comparison of percutaneous endoscopic lumbar discectomy versus open lumbar microdiscectomy for lumbar disc herniation: A meta-analysis’, *International journal of surgery (London, England)*, 31, pp. 86–92. Available at: <https://doi.org/10.1016/J.IJSU.2016.05.061>.
- Sánchez-Margallo, J.A. *et al.* (2021) ‘Wearable Technology for Assessment and Surgical Assistance in Minimally Invasive Surgery’, *Advances in Minimally Invasive Surgery* [Preprint]. Available at: <https://doi.org/10.5772/INTECHOPEN.100617>.
- Sassack, B. and Carrier, J.D. (2023) ‘Anatomy, Back, Lumbar Spine’, *StatPearls* [Preprint]. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK557616/> (Accessed: 8 September 2024).
- Scheer, J.K. *et al.* (2017) ‘Initial Experience With Real-Time Continuous Physical Activity Monitoring in Patients Undergoing Spine Surgery’, *Clinical spine surgery*, 30(10), pp. E1434–E1443. Available at: <https://doi.org/10.1097/BSD.0000000000000521>.
- Schulte, T.L. *et al.* (2010) ‘Step activity monitoring in lumbar stenosis patients undergoing decompressive surgery’, *European Spine Journal*, 19(11), p. 1855. Available at: <https://doi.org/10.1007/S00586-010-1324-Y>.
- Sesay, M. *et al.* (2015) ‘Responses of heart rate variability to acute pain after minor spinal surgery: optimal thresholds and correlation with the numeric rating scale’, *Journal of neurosurgical anesthesiology*, 27(2), pp. 148–154. Available at: <https://doi.org/10.1097/ANA.000000000000102>.
- Stienen, M.N. *et al.* (2020) ‘Objective activity tracking in spine surgery: a prospective feasibility study with a low-cost consumer grade wearable accelerometer’, *Scientific Reports 2020 10:1*, 10(1), pp. 1–11. Available at: <https://doi.org/10.1038/s41598-020-61893-4>.

- Tang, M. *et al.* (2023) 'Wearable technology and measurement of outcomes from lumbar spondylolisthesis surgery', *Seminars in Spine Surgery*, 35(3), pp. 101047. Available at: <https://doi.org/10.1016/J.SEMSS.2023.101047>.
- Telfeian, A.E. *et al.* (2016) 'A brief history of endoscopic spine surgery', *Neurosurgical focus*, 40(2), pp. 1–5. Available at: <https://doi.org/10.3171/2015.11.FOCUS15429>.
- Teplick, J.G. and Haskin, M.E. (2019) 'Spontaneous regression of herniated nucleus pulposus: The clinical findings of 76 patients', *Medicine*, 98(8), pp. 331–335. Available at: <https://doi.org/10.1097/MD.00000000000014667>.
- Thompson, J.C.. and Netter, F.H.. (2015) 'Netter's concise orthopaedic anatomy'. Available at: https://books.google.com/books/about/Netter_s_Concise_Orthopaedic_Anatomy.html?id=MeAyLgEACAAJ (Accessed: 8 September 2024).
- Turner, L.J. *et al.* (2012) 'Reliability of pedometers to measure step counts in patients with chronic respiratory disease', *Journal of cardiopulmonary rehabilitation and prevention*, 32(5), pp. 284–291. Available at: <https://doi.org/10.1097/HCR.0B013E31825C49F2>.
- Vianin, M. (2008) 'Psychometric properties and clinical usefulness of the Oswestry Disability Index', *Journal of chiropractic medicine*, 7(4), pp. 161–163. Available at: <https://doi.org/10.1016/J.JCM.2008.07.001>.
- Wu, X. *et al.* (2019) 'Comparison of Clinical Outcomes of Two-Level PELD and Foraminoplasty PELD for Highly Migrated Disc Herniations: A Comparative Study', *BioMed Research International*, 2019. Available at: <https://doi.org/10.1155/2019/9681424>.
- Zhang, C. *et al.* (2021) 'A Postoperative Phenomenon of Percutaneous Endoscopic Lumbar Discectomy: Rebound Pain'. Available at: <https://doi.org/10.1111/os.13088>.