

DAFTAR PUSTAKA

- Adwan L, Al-Sadi T, Shawakha S, Al-Shami NA. (2023). Clinical outcomes of COVID-19 in hemodialysis patients. *Front Med (Lausanne)*. 13;10:1281594. doi: 10.3389/fmed.2023.1281594. PMID: 38020138; PMCID: PMC10679386.
- Ahmadi, Sahar., Seyedeh, Zahra., Ranjbaran, Mehdi. (2024). Survival Rate of Hemodialysis Patients and Its Predictors: A Multicentric Retrospective Cohort Study. *Jundishapur Journal Of Chronic Disease Care*.
<https://doi.org/10.5812/jjcdc-142669>.
- Bazeley J, Bieber B, Li Y, Morgenstern H, de Sequera P. (2018). C-reactive protein and prediction of 1-year mortality in prevalent hemodialysis patients. *Clin J Am Soc Nephrol*. (10):2452-61. doi: 10.2215/CJN.00710111.
- Benzing, T. and Salant, D. (2021). ‘Insights into Glomerular Filtration and Albuminuria’, *N. Engl. J. Med*, pp. 1437–1446.
- Bouillanne, O. *et al.* (2005). *Geriatric Nutritional Risk Index: a new index for evaluating at-risk elderly medical patients 1-3*.
- Buzby, G., Knox, L. and Crosby, L. (2008) ‘Study protocol: a randomized clinical trial of total parenteral nutrition in malnourished surgical patients’, *Am J Clin Nutr*, 47.
- Carrero JJ, Stenvinkel P, Cuppari L, Ikizler TA, Kalantar-Zadeh K, Kaysen G. (2013). Etiology of the protein-energy wasting syndrome in chronic kidney disease: a consensus statement from the international society of renal nutrition and metabolism (ISRNM). *J Ren Nutr* 23:77–90.
- Chen Y, Liang X, Zheng S. (2018). Association of Body Fat Mass and Fat Distribution With the Incidence of Hypertension in a Population-Based Chinese Cohort: A 22-Year Follow-Up. *J Am Heart Assoc*. DOI: 10.1161/JAHA.117.007153

- Cho, A., Hong, Y.S., Park, H.C. *et al.* (2022). Geriatric nutritional risk index is associated with retinopathy in patients with type 2 diabetes. *Sci Rep* **12**, 11746. <https://doi.org/10.1038/s41598-022-15463-5>
- Cockwell, P. and Fisher, L.A. (2020). ‘The global burden of chronic kidney disease’, *The Lancet*, pp. 662–664.
- Dad T, Weiner DE. (2015). Stroke and Chronic Kidney Disease: Epidemiology, Pathogenesis, and Management Across Kidney Disease Stages. *Semin Nephrol.* 35(4):311-22. doi: 10.1016/j.semnephrol.2015.06.003. PMID: 26355250; PMCID: PMC4567691.
- Davenport, A. (2014). ‘Chronic Kidney Failure. Kidney Transplantation—Principles and Practice’, pp. 39–53.
- Fang Y, Gong AY, Haller ST, Dworkin LD, Liu Z, Gong R. (2020). The ageing kidney: Molecular mechanisms and clinical implications. *Ageing Res Rev.* 2020 Nov;63:101151. doi: 10.1016/j.arr.2020.101151. Epub Aug 22. PMID: 32835891; PMCID: PMC7595250.
- Feneberg, R., Schaefer, F. and Veldhuis, J. (2003). ‘Neuroendocrine adaptations in renal disease’, *Pediatr Nephrol* [Preprint].
- Fiaccadori E, Sabatino A, Barazzoni R, Carrero JJ, Cupisti A, De Waele E. (2021). ESPEN guideline on clinical nutrition in hospitalized patients with acute or chronic kidney disease. *Clin Nutr.* 2021 Apr;40(4):1644-1668. doi: 10.1016/j.clnu.2021.01.028. Epub. PMID: 33640205.
- Hamirudin, A., Charlton, K, Walton, K. (2016) ‘Outcomes related to nutrition screening in community living older adults: a systematic literature review.’, *Arch Gerontol Geriatri* [Preprint].
- Harmilah. (2020). *Asuhan Keperawatan Pada Pasien dengan Gangguan Sistem Perkemihan*. Pustaka Baru Press.
- He, J. (2022). Creatinine-to-body weight ratio is a predictor of incident diabetes: a population-based retrospective cohort study. *Diabetol Metab Syndr* **14**, 7. <https://doi.org/10.1186/s13098-021-00776-8>

- Ikizler TA, Cano NJ, Franch H, Fouque D, Himmelfarb J, Kalantar-Zadeh K. (2013). International Society of Renal Nutrition and Metabolism. Prevention and treatment of protein energy wasting in chronic kidney disease patients: a consensus statement by the International Society of Renal Nutrition and Metabolism. *Kidney Int.* 84(6):1096-107. doi: 10.1038/ki.2013.147. Epub 2013 May 22. PMID: 23698226.
- Inker, L., Astor, B, Fox, C. (2014) ‘KDOQI us commentary on the 2012 KDIGO clinical practice guideline for the evaluation and management of CKD’, *Am J Kidney Dis*, 63(713–35).
- Juliana Rodrigues, Lilian Cuppari, Katrina L Campbell, Carla Maria Avesani. (2017). Nutritional assessment of elderly patients on dialysis: pitfalls and potentials for practice, *Nephrology Dialysis Transplantation*, Volume 32, Issue 11, Pages 1780–1789, <https://doi.org/10.1093/ndt/gfw471>
- Kalantar, Z. Fouque, K. (2014) ‘Outcome research, nutrition, and reverse epidemiology in maintenance dialysis patients’, *J. Ren. Nutr*, 14, pp. 64–71.
- Kanno, Y., Kanda, E. and Kato, A. (2021) ‘Methods and nutritional interventions to improve the nutritional status of dialysis patients in JAPAN’, *A narrative review. Nutrients* [Preprint].
- Kasiske, B. (2013) ‘Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease’, *Official Journal of The International Society of Nephrology*, 3.
- Kementerian Kesehatan. (2020). *Buku Kesehatan Lansia*. Kementerian Kesehatan Republik Indonesia.
- Kim DH, Lee YK, Park HC, Kim BY, Lee M, Kim GO, Kim J, Cho A. (2023). The Geriatric Nutritional Risk Index Is an Optimal Evaluation Parameter for Predicting Mortality in Adult All Ages Hemodialysis Patients: A Korean Population-Based Study. *Nutrients*. 1;15(17):3831. doi: 10.3390/nu15173831. PMID: 37686863; PMCID: PMC10490009.

- Kobayashi I, Ishimura E, Kato Y, Okuno S, Yamamoto T, Yamakawa T. (2010). Geriatric Nutritional Risk Index, a simplified nutritional screening index, is a significant predictor of mortality in chronic dialysis patients. *Nephrol Dial Transplant*. (10):3361-5. doi: 10.1093/ndt/gfq211. Epub 2010 Apr 16. PMID: 20400447.
- Komatsu M, Okazaki M, Tsuchiya K, Kawaguchi H, Nitta K. (2015). Geriatric Nutritional Risk Index Is a Simple Predictor of Mortality in Chronic Hemodialysis Patients. *Blood Purif*. 39(4):281-7. doi: 10.1159/000381798.
- Koppe L, Pillon NJ, Vella RE, Croze ML, Pelletier CC *et al.*, (2013). p-Cresyl sulfate promotes insulin resistance associated with CKD. *J Am Soc Nephrol*. (1):88-99. doi: 10.1681/ASN.2012050503. PMID: 23274953; PMCID: PMC3537215.
- Kovesdy CP. (2022). Epidemiology of chronic kidney disease: an update 2022. *Kidney Int Suppl*. 2022 Apr;12(1):7-11. doi: 10.1016/j.kisu.2021.11.003. Epub 2022 Mar 18. PMID: 35529086; PMCID: PMC9073222.
- Krzywińska-Siemaszko R, Mossakowska M, Skalska A, Klich-Rączka A, Tobis S, Szybalska A. (2015). Social and economic correlates of malnutrition in Polish elderly population: the results of PolSenior study. *J Nutr Health Aging*. 19(4):397-402. doi: 10.1007/s12603-014-0572-7. PMID: 25809803.
- Kusnandar and Budy, V. (2023) *Persentase Penduduk Lansia menurut Provinsi, <https://databoks.katadata.co.id/datapublish/2022/05/31/wilayah-mana-yang-punya-porsi-penduduk-lansia-terbesar>*.
- Larbi A, Cabreiro F, Zelba H, Marthandan S, Combet E. (2010). Reduced oxygen tension results in reduced human T cell proliferation and increased intracellular oxidative damage and susceptibility to apoptosis upon activation. *Free Radic Biol Med*. 1;48(1):26-34. doi: 10.1016/j.freeradbiomed.2009.09.025. Epub 2009 Sep 28. PMID: 19796677.
- Li Y, Shen J, Hou X, Su Y, Jiao Y, Wang J, Liu H, Fu Z. (2023). Geriatric nutritional risk index predicts all-cause mortality in the oldest-old patients with acute

coronary syndrome: A 10-year cohort study. *Front Nutr.* Mar 7;10:1129978. doi: 10.3389/fnut.2023.1129978. PMID: 36960208; PMCID: PMC10027908.

Marlies Noordzij, Kitty J. Jager. (2014). Increased mortality early after dialysis initiation: a universal phenomenon. *Kidney International*: Volume 85: 12-14. <https://doi.org/10.1038/ki.2013.316>.

Maseda A, Diego-Diez C, Lorenzo-López L, López-López R, Regueiro-Folgueira L, Millán-Calenti JC. (2018). Quality of life, functional impairment and social factors as determinants of nutritional status in older adults: The VERISAÚDE study. *Clin Nutr.* 37(3):993-999. doi: 10.1016/j.clnu.2017.04.009. Epub 2017 Apr 15. PMID: 28456537.

Matsukuma Y, Tanaka S, Taniguchi M, Nakano T, Masutani K, Hirakata. (2019). Association of geriatric nutritional risk index with infection-related mortality in patients undergoing hemodialysis: The Q-Cohort Study. *Clin Nutr.* 38(1):279-287. doi: 10.1016/j.clnu.2018.01.019. Epub 2018 Feb 15. PMID: 29398335.

Matthew AK, Melamed ML, Bauer C, Raff AC, Hostetter TH. (2013). Effects of oral sodium bicarbonate in patients with CKD. *Clin J Am Soc Nephrol.* 8(5):714-20. doi: 10.2215/CJN.08340812. Epub 2013 Feb 7. PMID: 23393105; PMCID: PMC3641618.

McDougall KA, Larkin JW, Wingard RL, Jiao Y, Rosen S *et al.*, (2018). Depressive affect in incident hemodialysis patients. *Clin Kidney J.* 11(1):123-129. doi: 10.1093/ckj/sfx054.

Msaad R, Essadik R, Mohtadi K, Meftah H, Lebrazi H, *et al.*, (2019). Predictors of mortality in hemodialysis patients. *Pan Afr Med J.* 28;33:61. doi: 10.11604/pamj.2019.33.61.18083. PMID: 31448023; PMCID: PMC6689835.

Muhayati, S. (2018). *Konseling Lansia: Upaya Lanjut Usia dalam Membangun Kemandirian Hidup dan Penerimaan Diri Terhadap Kesiapan Memasuki Masa Lanjut.*

- Munawirah Munawirah, Masrul Masrul, Rose Dinda Martini. (2017). Hubungan Beberapa Faktor Risiko dengan Malnutrisi pada Usia Lanjut di Nagari Sijunjung Kecamatan Sijunjung. *Jurnal Kesehatan Andalas* Vol 6, No 2.
- Myung J, Choi JH, Yi JH, Kim I. (2020). Cancer incidence according to the National Health Information Database in Korean patients with end-stage renal disease receiving hemodialysis. *Korean J Intern Med*. 2020 Sep;35(5):1210-1219. doi: 10.3904/kjim.2018.400. Epub Feb 20. PMID: 32066223; PMCID: PMC7487292.
- Nakagawa N, Maruyama K, Hasebe N. (2021). Utility of Geriatric Nutritional Risk Index in Patients with Chronic Kidney Disease: A Mini-Review. *Nutrients*. doi: 10.3390/nu13113688. PMID: 34835944; PMCID: PMC8624060.
- National Kidney Foundation. (2016). KDOQI Clinical Practice Guideline for Hemodialysis Adequacy. *Am J Kidney Dis*. 66(5):884-930. doi: 10.1053/j.ajkd.2015.07.015.
- Nur Adilah, Nur Islami, Mohd Fahmi Teng, Mazuin Kamarul. (2019). Geriatric Nutrition Risk Index is comparable to the mini nutritional assessment for assessing nutritional status in elderly hospitalized patients. *Clinical Nutrition*. Vol. 29: 77-85.
- O'Sullivan, E.D., Hughes, J. and Ferenbach, D.A. (2017) 'Renal Aging: Causes and Consequences', *J. Am. Soc. Nephrol*, pp. 407–420.
- Park JH, Kim SB, Shin HS *et al*. (2012). Geriatric nutritional risk index may be a significant predictor of mortality in Korean hemodialysis patients: a single center study. *Ther Apher Dial* 16:121–126.
- Panichi V, Cupisti A, Rosati A, Di Giorgio A, Scatena A, Menconi O. (2014). Geriatric nutritional risk index is a strong predictor of mortality in hemodialysis patients: data from the Riscavid cohort. *J Nephrol*. Apr;27(2):193-201. doi: 10.1007/s40620-013-0033-0. Epub 2014 Jan 16. PMID: 24430765.
- Potter, P. and Perry, A. (2005) *Buku Ajar Fundamental Keperawatan : Konsep, Proses, dan Praktik*. EGC.

- Raphael KL, Zhang Y, Ying J, Greene T. (2014). Prevalence of and risk factors for reduced serum bicarbonate in chronic kidney disease. *Nephrology* (Carlton). Oct;19(10):648-54. doi: 10.1111/nep.12315. PMID: 25066359.
- Rastad, H., Ejtahed, HS., Shafiee, G. (2021). The risk factors associated with COVID-19-Related death among patients with end-stage renal disease. *BMC Nephrol.* <https://doi.org/10.1186/s12882-020-02221>.
- Riskesdas. (2018). *Laporan hasil riset kesehatan dasar (riskesdas), from <https://www.litbang.kemkes.go.id/laporan-riset-kesehatan-dasar-riskesdas/>*.
- Robinson BM, Zhang J, Morgenstern H, Bradbury BD, Ng LJ, McCullough KP,. (2014). Worldwide, mortality risk is high soon after initiation of hemodialysis. *Kidney Int.* 85(1):158-65. doi: 10.1038/ki.2013.252. Epub 2013 Jun 26. PMID: 23802192; PMCID: PMC3877739.
- Saeed Fahad, Susana Arrigain, Jesse D. Schold, Joseph V. Nally Jr, Sankar Dass Navaneethan. (2019). What are the Risk Factors for One-Year Mortality in Older Patients with Chronic Kidney Disease? An Analysis of the Cleveland Clinic CKD Registry. *Nephron.* 141 (2): 98–104. <https://doi.org/10.1159/000494298>
- Serón-Arbeloa C, Labarta-Monzón L, Puzo-Foncillas J, Mallor-Bonet T, Lafita-López A *et al.*, (2022). Malnutrition Screening and Assessment. *Nutrients.* 9;14(12):2392. doi: 10.3390/nu14122392. PMID: 35745121; PMCID: PMC9228435.
- Setiati, I. (2014) *Buku Ajar Ilmu Penyakit Dalam*. Jakarta: Internal Publishing.
- Smeltzer, S.C. and Bare, B.G. (2012) *Buku Ajar Keperawatan medical bedah Brunner & Suddarth*.
- Sun DF, Zheng Z, Tummala P, Oh J, Schaefer F, Rabkin R. (2004). Chronic uremia attenuates growth hormone-induced signal transduction in skeletal muscle. *J Am Soc Nephrol.* (10):2630-6. doi: 10.1097/01.ASN.0000139492.36400.6C. PMID: 15466267.
- Takahashi H, Ito Y, Ishii H, Aoyama T, Kamoi D, *et al.*, (2014). Geriatric nutritional risk index accurately predicts cardiovascular mortality in incident hemodialysis

- patients. *J Cardiol*. 64(1):32-6. doi: 10.1016/j.jjcc.2013.10.018. Epub 2013 Dec 22. PMID: 24365385.
- Tanaka, A, Inaguma, D., Shinjo, H. (2017). Relationship between mortality and Geriatric Nutritional Risk Index (GNRI) at the time of dialysis initiation: a prospective multicenter cohort study. *Ren Replace Ther* . <https://doi.org/10.1186/s41100-017-0108-9>.
- Tollitt, J., Odudu, A., Flanagan, E. (2019). Impact of prior stroke on major clinical outcome in chronic kidney disease: the Salford kidney cohort study. *BMC Nephrol* **20**, 432. <https://doi.org/10.1186/s12882-019-1614-5>
- Vallet M, Metzger M, Haymann JP, Flamant M, Gauci C, Thervet E. (2015).. Urinary ammonia and long-term outcomes in chronic kidney disease. *Kidney Int*. 2015 Jul;88(1):137-45. doi: 10.1038/ki.2015.52. Epub. PMID: 25760321.
- Varleta P, Tagle R. (2019). A feared combination: Hypertension and chronic kidney disease. *J Clin Hypertens (Greenwich)*. 21(1):102-104. doi: 10.1111/jch.13432. Epub 2018 Nov 14. PMID: 30427110; PMCID: PMC8030333.
- Vaziri ND, Yuan J, Rahimi A, Ni Z, Said H, Subramanian VS. (2012). Disintegration of colonic epithelial tight junction in uremia: a likely cause of CKD-associated inflammation. *Nephrol Dial Transplant*. 27(7):2686-93. doi: 10.1093/ndt/gfr624. Epub 2011 Nov 29. PMID: 22131233; PMCID: PMC3616758.
- Wang, Renjie MDa; Liang, Yuxiang MDa; Jiang, Jiaojiao MDa. (2024). Association between the geriatric nutritional risk index and clinical outcomes among peritoneal dialysis patients: A meta-analysis. *Medicine* 103(18):p e38048. | DOI: 10.1097/MD.00000000000038048.
- Watanabe H, Miyamoto Y, Honda D, Tanaka H, Wu Q, Endo M. (2013). p-Cresyl sulfate causes renal tubular cell damage by inducing oxidative stress by activation of NADPH oxidase. *Kidney Int*. 83(4):582-92. doi: 10.1038/ki.2012.448. Epub 2013 Jan 16. PMID: 23325087.
- Wong J, Piceno YM, DeSantis TZ, Pahl M, Andersen GL, Vaziri ND. (2014). Expansion of urease- and uricase-containing, indole- and p-cresol-forming and

- contraction of short-chain fatty acid-producing intestinal microbiota in ESRD. *Am J Nephrol.* 39(3):230-237. doi: 10.1159/000360010. Epub 2014 Mar 8. PMID: 24643131; PMCID: PMC4049264.
- Xiong J, Wang M, Zhang Y, Nie L, He T, Wang Y. (2018). Association of Geriatric Nutritional Risk Index with Mortality in Hemodialysis Patients: A Meta-Analysis of Cohort Studies. *Kidney Blood Press Res.* 43(6):1878-1889. doi: 10.1159/000495999. Epub 2018 Dec 14. PMID: 30566933.
- Xiong, J., Wang, M., Wang, J., Nie L, He. (2019). Geriatric nutrition risk index is associated with renal progression, cardiovascular events and all-cause mortality in chronic kidney disease. *J Nephrol.* Vol 33.
- Xu JJ, Jia SD, Zhu P, Jiang L, Jiang P *et al.*, (2021). Does Prior Stroke Predict Long-Term Recurrent Stroke After Percutaneous Coronary Intervention? Five-Year Results From a Large Cohort Study. *Front Neurol.* Nov 2;12:740136. doi: 10.3389/fneur.2021.740136. PMID: 34795628; PMCID: PMC8593040.
- Yamada K, Furuya R, Takita T, Maruyama Y, Yamaguchi Y, Ohkawa S. (2008). Simplified nutritional screening tools for patients on maintenance hemodialysis. *Am J Clin Nutr.* 87(1):106-13. doi: 10.1093/ajcn/87.1.106. PMID: 18175743.
- Yin, H., Price, F. and Rudnicki, M. (2013) 'Satellite cells and the muscle stem cell niche', *Physiol Rev* [Preprint].
- Yu Fan, Lian He, Yongjing Zhou and Changfeng Man. (2021). Predictive Value of Geriatric Nutritional Risk Index in Patients With Coronary Artery Disease: A Meta-Analysis. *Sec. Clinical Nutrition Volume 8*
- Zha, Y. and Qian, Q. (2017) 'Protein nutrition and malnutrition in CKD and ESRD', *Nutrients.* MDPI AG. Available at: <https://doi.org/10.3390/nu9030208>.