

REFERENCES

- Agati, G., Azzarello, E., Pollastri, S., & Tattini, M. (2012). Flavonoids as antioxidants in plants: location and functional significance. *Plant science*, *196*, 67-76.
- Agunloye, O. M., Oboh, G., Ademiluyi, A. O., Ademosun, A. O., Akindahunsi, A. A., Oyagbemi, A. A., Omobowale, T. O., Ajibade, T. O., & Adedapo, A. A. (2019). Cardio-protective and antioxidant properties of caffeic acid and chlorogenic acid: Mechanistic role of angiotensin converting enzyme, cholinesterase and arginase activities in cyclosporine induced hypertensive rats. *Biomedicine & Pharmacotherapy*, *109*, 450-458.
- Al-Rajhi, A. M., & Ghany, T. M. A. (2023). In vitro repress of breast cancer by bio-product of edible *Pleurotus ostreatus* loaded with chitosan nanoparticles. *Applied Biological Chemistry*, *66*(1), 33.
<https://doi.org/https://doi.org/10.1186/s13765-023-00788-0>
- Al Mamari, H. H. (2021). *Phenolic compounds: Classification, chemistry, and updated techniques of analysis and synthesis*. IntechOpen.
- Alam, N., Yoon, K. N., Lee, K. R., Kim, H. Y., Shin, P. G., Cheong, J. C., Yoo, Y. B., Shim, M. J., Lee, M. W., & Lee, T. S. (2011). Assessment of antioxidant and phenolic compound concentrations as well as xanthine oxidase and tyrosinase inhibitory properties of different extracts of *Pleurotus citrinopileatus* fruiting bodies. *Mycobiology*, *39*(1), 12-19.
<https://doi.org/https://doi.org/10.4489/MYCO.2011.39.1.012>

- Alara, O. R., Abdurahman, N. H., Ukaegbu, C. I., & Azhari, N. H. (2018). Vernonia cinerea leaves as the source of phenolic compounds, antioxidants, and anti-diabetic activity using microwave-assisted extraction technique. *Industrial Crops and Products*, 122, 533-544. <https://doi.org/https://doi.org/10.1016/j.indcrop.2018.06.034>
- Aliaño-González, M. J., Barea-Sepúlveda, M., Espada-Bellido, E., Ferreiro-González, M., López-Castillo, J. G., Palma, M., Barbero, G. F., & Carrera, C. (2022). Ultrasound-assisted extraction of total phenolic compounds and antioxidant activity in mushrooms. *Agronomy*, 12(8), 1812. <https://doi.org/https://doi.org/10.3390/agronomy12081812>
- Amirullah, N. A., Abidin, N. Z., Abdullah, N., & Manickam, S. (2021). Application of ultrasound towards improving the composition of phenolic compounds and enhancing in vitro bioactivities of *Pleurotus pulmonarius* (Fr.) QuéL extracts. *Biocatalysis and Agricultural Biotechnology*, 31, 101881. <https://doi.org/https://doi.org/10.1016/j.bcab.2020.101881>
- Azwanida, N. (2015). A review on the extraction methods use in medicinal plants, principle, strength and limitation. *Med aromat plants*, 4(196), 2167-0412.
- Bach, F., Zielinski, A. A. F., Helm, C. V., Maciel, G. M., Pedro, A. C., Stafussa, A. P., Ávila, S., & Haminiuk, C. W. I. (2019). Bio compounds of edible mushrooms: In vitro antioxidant and antimicrobial activities. *LWT*, 107, 214-220. <https://doi.org/https://doi.org/10.1016/j.lwt.2019.03.017>
- Berger, P. D., Maurer, R. E., Celli, G. B., Berger, P. D., Maurer, R. E., & Celli, G. B. (2018). Literature on Experimental Design and Discussion of Some

Topics Not Covered in the Text. *Experimental Design: With Application in Management, Engineering, and the Sciences.*, 611-623.

Bhattacharya, M., Srivastav, P., & Mishra, H. (2014). Optimization of microwave-convective drying of Oyster mushrooms (*Pleurotus ostreatus*) using response-surface methodology. *International Food Research Journal*, 21(4), 1575-1581.

Boonsong, S., Klaypradit, W., & Wilaipun, P. (2016). Antioxidant activities of extracts from five edible mushrooms using different extractants. *Agriculture and Natural Resources*, 50(2), 89-97.
<https://doi.org/http://dx.doi.org/10.1016/j.anres.2015.07.002>

Cateni, F., Gargano, M. L., Procida, G., Venturella, G., Cirlincione, F., & Ferraro, V. (2021). Mycochemicals in wild and cultivated mushrooms: nutrition and health. *Phytochemistry Reviews*, 1-45.

Çayan, F., Tel-Çayan, G., Deveci, E., & Duru, M. E. (2021). HPLC–DAD characterization of phenolic profile and in vitro antioxidant, anticholinesterase, and antidiabetic activities of five mushroom species from Turkey. *3 Biotech*, 11(6), 273.
<https://doi.org/https://doi.org/10.1007/s13205-021-02819-3>

Chan, C.-H., Yusoff, R., Ngoh, G.-C., & Kung, F. W.-L. (2011). Microwave-assisted extractions of active ingredients from plants. *Journal of chromatography A*, 1218(37), 6213-6225.
<https://doi.org/https://doi.org/10.1016/j.chroma.2011.07.040>

- Chemat, F., & Khan, M. K. (2011). Applications of ultrasound in food technology: processing, preservation and extraction. *Ultrasonics Sonochemistry*, *18*(4), 813-835.
- Chu, M., Khan, R. D., Zhou, Y., Agar, O. T., Barrow, C. J., Dunshea, F. R., & Suleria, H. A. (2023). LC-ESI-QTOF-MS/MS characterization of phenolic compounds in common commercial mushrooms and their potential antioxidant activities. *Processes*, *11*(6), 1711. <https://doi.org/https://doi.org/10.3390/pr11061711>
- Contato, A. G., Inácio, F. D., de Araújo, C. A. V., Brugnari, T., Maciel, G. M., Haminiuk, C. W. I., Bracht, A., Peralta, R. M., & de Souza, C. G. M. (2020). Comparison between the aqueous extracts of mycelium and basidioma of the edible mushroom *Pleurotus pulmonarius*: Chemical composition and antioxidant analysis. *Journal of Food Measurement and Characterization*, *14*, 830-837. <https://doi.org/https://doi.org/10.1007/s11694-019-00331-0>
- Corrêa, R. C. G., Brugnari, T., Bracht, A., Peralta, R. M., & Ferreira, I. C. (2016). Biotechnological, nutritional and therapeutic uses of *Pleurotus* spp.(Oyster mushroom) related with its chemical composition: A review on the past decade findings. *Trends in Food Science & Technology*, *50*, 103-117.
- Cotas, J., Leandro, A., Monteiro, P., Pacheco, D., Figueirinha, A., Gonçalves, A. M., da Silva, G. J., & Pereira, L. (2020). Seaweed phenolics: From extraction to applications. *Marine drugs*, *18*(8), 384.
- Dahmoune, F., Spigno, G., Moussi, K., Remini, H., Cherbal, A., & Madani, K. (2014). *Pistacia lentiscus* leaves as a source of phenolic compounds: Microwave-assisted extraction optimized and compared with ultrasound-

assisted and conventional solvent extraction. *Industrial Crops and Products*, 61, 31-40.

Dai, J., & Mumper, R. J. (2010). Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*, 15(10), 7313-7352.

<https://doi.org/https://doi.org/10.3390/molecules15107313>

De Bruijn, J., Loyola, C., Aqueveque, P., Cañumir, J., Cortéz, M., & France, A. (2009). Antioxidant properties of extracts obtained from *Grifola gargar* mushrooms. *Micologia Aplicada International*, 21(1), 11-18.

Deepalakshmi, K., & Sankaran, M. (2014). *Pleurotus ostreatus*: an oyster mushroom with nutritional and medicinal properties. *Journal of Biochemical Technology*, 5(2), 718-726.

Deo, S., Janghel, A., Raut, P., Bhosle, D., Verma, C., Kumar, S. S., Agrawal, M., Amit, N., Sharma, M., & Giri, T. (2015). Emerging microwave assisted extraction (MAE) techniques as an innovative green technologies for the effective extraction of the active phytopharmaceuticals. *Research Journal of Pharmacy and Technology*, 8(5), 655-666.

Desai, M. A., & Parikh, J. (2012). Microwave assisted extraction of essential oil from *Cymbopogon flexuosus* (Steud.) Wats.: a parametric and comparative study. *Separation Science and Technology*, 47(13), 1963-1970.

Devi, P. V., Islam, J., Narzary, P., Sharma, D., & Sultana, F. (2024). Bioactive compounds, nutraceutical values and its application in food product development of oyster mushroom. *Journal of Future Foods*, 4(4), 335-342.

<https://doi.org/https://doi.org/10.1016/j.jfutfo.2023.11.005>

- Dludla, P. V., Nkambule, B. B., Jack, B., Mkandla, Z., Mutize, T., Silvestri, S., Orlando, P., Tiano, L., Louw, J., & Mazibuko-Mbeje, S. E. (2018). Inflammation and oxidative stress in an obese state and the protective effects of gallic acid. *Nutrients*, *11*(1), 23.
- Duraisamy, R., Shuge, T., Worku, B., Berekete, A. K., & Ramasamy, K. M. (2020). Extraction, screening and spectral characterization of tannins from acacia xanthophloea (Fever Tree) Bark. *Research Journal of Textile and Leather*, *1*, 1-10. <https://doi.org/https://doi.org/10.46590/rjtl.2020.010101>
- Effiong, M. E., Bella-Omunagbe, M., Afolabi, I. S., & Chinedu, S. N. (2024). In silico evaluation of potential breast cancer receptor antagonists from GC-MS and HPLC identified compounds in *Pleurotus ostreatus* extracts. *RSC advances*, *14*(33), 23744-23771. <https://doi.org/https://doi.org/10.1039/d4ra03832k>
- Eskilsson, C. S., & Björklund, E. (2000). Analytical-scale microwave-assisted extraction. *Journal of chromatography A*, *902*(1), 227-250. [https://doi.org/https://doi.org/10.1016/S0021-9673\(00\)00921-3](https://doi.org/https://doi.org/10.1016/S0021-9673(00)00921-3)
- Espada-Bellido, E., Ferreiro-González, M., Carrera, C., Palma, M., Álvarez, J. A., F. Barbero, G., & Ayuso, J. (2019). Extraction of antioxidants from blackberry (*Rubus ulmifolius* L.): comparison between ultrasound-and microwave-assisted extraction techniques. *Agronomy*, *9*(11), 745. <https://doi.org/http://dx.doi.org/10.3390/agronomy9110745>
- Fernandes, Â., Barros, L., Martins, A., Herbert, P., & Ferreira, I. C. (2015). Nutritional characterisation of *Pleurotus ostreatus* (Jacq. ex Fr.) P. Kumm.

produced using paper scraps as substrate. *Food Chemistry*, 169, 396-400.

<https://doi.org/https://doi.org/10.1016/j.foodchem.2014.08.027>

Fernández, P., Vázquez, C., Lorenzo, R., Carro, A., Álvarez, I., & Cabarcos, P.

(2010). Experimental design for optimization of microwave-assisted extraction of benzodiazepines in human plasma. *Analytical and Bioanalytical Chemistry*, 397, 677-685.

<https://doi.org/https://doi.org/10.1007/s00216-010-3572-4>

Ferreira, S. C., Bruns, R., Ferreira, H. S., Matos, G. D., David, J., Brandão, G., da

Silva, E. P., Portugal, L., Dos Reis, P., & Souza, A. (2007). Box-Behnken design: An alternative for the optimization of analytical methods. *Analytica chimica acta*, 597(2), 179-186.

Flores, G. A., Girometta, C. E., Cusumano, G., Angelini, P., Tirillini, B., Ianni, F.,

Blasi, F., Cossignani, L., Pellegrino, R. M., & Emiliani, C. (2022). Untargeted metabolomics used to describe the chemical composition, antioxidant and antimicrobial effects of extracts from *Pleurotus* spp. mycelium grown in different culture media. *Antibiotics*, 11(11), 1468.

<https://doi.org/https://doi.org/10.3390/antibiotics11111468>

Fogarasi, M., Socaciu, M.-I., Sălăgean, C.-D., Ranga, F., Fărcaș, A. C., Socaci, S.

A., Socaciu, C., Țibulcă, D., Fogarasi, S., & Semeniuc, C. A. (2021). Comparison of different extraction solvents for characterization of antioxidant potential and polyphenolic composition in *Boletus edulis* and *Cantharellus cibarius* mushrooms from Romania. *Molecules*, 26(24), 7508.

<https://doi.org/https://doi.org/10.3390/molecules26247508>

- Fraga, C. G., Croft, K. D., Kennedy, D. O., & Tomás-Barberán, F. A. (2019). The effects of polyphenols and other bioactives on human health. *Food & function*, *10*(2), 514-528.
- Gao, M., Song, B.-Z., & Liu, C.-Z. (2006). Dynamic microwave-assisted extraction of flavonoids from *Saussurea medusa* Maxim cultured cells. *Biochemical Engineering Journal*, *32*(2), 79-83.
- Garcia, K., Garcia, C. J., Bustillos, R., & Dulay, R. M. (2020). Mycelial biomass, antioxidant, and myco-actives of mycelia of abalone mushroom *Pleurotus cystidiosus* in liquid culture. *J Appl Biol Biotechnol*, *8*(2), 94-97. <https://doi.org/http://dx.doi.org/10.7324/JABB.2020.80215>
- Gąsecka, M., Mleczek, M., Siwulski, M., & Niedzielski, P. (2016). Phenolic composition and antioxidant properties of *Pleurotus ostreatus* and *Pleurotus eryngii* enriched with selenium and zinc. *European Food Research and Technology*, *242*, 723-732. <https://doi.org/https://doi.org/10.1007/s00217-015-2580-1>
- Golak-Siwulska, I., Kałużewicz, A., Spizewski, T., Siwulski, M., & Sobieralski, K. (2018). Bioactive compounds and medicinal properties of Oyster mushrooms (*Pleurotus* sp.). *Folia Horticulturae*, *30*(2), 191-201. <https://doi.org/https://doi.org/10.2478/fhort-2018-0012>
- Gong, J., Zhou, S., & Yang, S. (2019). Vanillic acid suppresses HIF-1 α expression via inhibition of mTOR/p70S6K/4E-BP1 and Raf/MEK/ERK pathways in human colon cancer HCT116 cells. *International journal of molecular sciences*, *20*(3), 465.

Harmita, H. (2004). Petunjuk pelaksanaan validasi metode dan Cara Perhitungannya. *Majalah Ilmu Kefarmasian*, 1(3), 1.

Hip SengYim, H. S., Fook YeeChye, F. Y., Swee KhengHo, S. K., & Chun WaiHo, C. W. (2009). Phenolic profiles of selected edible wild mushrooms as affected by extraction solvent, time and temperature. *As. J. Food Ag-Ind*, 2, 392-401.

Huie, C. W. (2002). A review of modern sample-preparation techniques for the extraction and analysis of medicinal plants. *Analytical and Bioanalytical Chemistry*, 373, 23-30.

Kumar, K., Mehra, R., Guiné, R. P., Lima, M. J., Kumar, N., Kaushik, R., Ahmed, N., Yadav, A. N., & Kumar, H. (2021). Edible mushrooms: A comprehensive review on bioactive compounds with health benefits and processing aspects. *Foods*, 10(12), 2996.
<https://doi.org/https://doi.org/10.3390/foods10122996>

Lattanzio, V. (2013). Phenolic compounds: introduction 50. *Nat. Prod*, 1543-1580.

Li, H., Deng, Z., Wu, T., Liu, R., Loewen, S., & Tsao, R. (2012). Microwave-assisted extraction of phenolics with maximal antioxidant activities in tomatoes. *Food Chemistry*, 130(4), 928-936.

Liazid, A., Palma, M., Brigui, J., & Barroso, C. G. (2007). Investigation on phenolic compounds stability during microwave-assisted extraction. *Journal of chromatography A*, 1140(1-2), 29-34.
<https://doi.org/https://doi.org/10.1016/j.chroma.2006.11.040>

- Mandal, V., Mohan, Y., & Hemalatha, S. (2007). Microwave assisted extraction— an innovative and promising extraction tool for medicinal plant research. *Pharmacognosy Reviews*, 1(1), 7-18.
- Matkovits, A., Fodor, M., & Jókai, Z. (2024). Analysis of Polyphenol Patterns of *Pleurotus ostreatus* Cultivars by UHPLC-ESI-MS/MS; Application of FT-NIR and Chemometric Methods, Classification Options. *Chemosensors*, 12(2), 19. <https://doi.org/https://doi.org/10.3390/chemosensors12020019>
- Mattila, P., Könkö, K., Eurola, M., Pihlava, J.-M., Astola, J., Vahteristo, L., Hietaniemi, V., Kumpulainen, J., Valtonen, M., & Piironen, V. (2001). Contents of vitamins, mineral elements, and some phenolic compounds in cultivated mushrooms. *Journal of Agricultural and Food Chemistry*, 49(5), 2343-2348. <https://doi.org/https://doi.org/10.1021/jf001525d>
- Mazzutti, S., Ferreira, S. R., Riehl, C. A., Smania Jr, A., Smania, F. A., & Martínez, J. (2012). Supercritical fluid extraction of *Agaricus brasiliensis*: antioxidant and antimicrobial activities. *The Journal of Supercritical Fluids*, 70, 48-56. <https://doi.org/https://doi.org/10.1016/j.supflu.2012.06.010>
- Medini, F., Fellah, H., Ksouri, R., & Abdelly, C. (2014). Total phenolic, flavonoid and tannin contents and antioxidant and antimicrobial activities of organic extracts of shoots of the plant *Limonium delicatulum*. *Journal of Taibah University for Science*, 8(3), 216-224. <https://doi.org/https://doi.org/10.1016/j.jtusci.2014.01.003>
- Mondal, A., & Bennett, L. L. (2016). Resveratrol enhances the efficacy of sorafenib mediated apoptosis in human breast cancer MCF7 cells through ROS, cell

cycle inhibition, caspase 3 and PARP cleavage. *Biomedicine & Pharmacotherapy*, 84, 1906-1914.

Muszynska, B., Sulkowska-Ziaja, K., & Ekiert, H. (2013). Phenolic acids in selected edible basidiomycota species: *Armillaria mellea*, *Boletus badius*, *Boletus edulis*, *Cantharellus cibarius*, *Lactarius deliciosus* and *Pleurotus ostreatus*. *Acta Scientiarum Polonorum. Hortorum Cultus*, 12(4), 107-116.

Nelson, S. O., & Datta, A. K. (2001). Dielectric properties of food materials and electric field interactions. In *Handbook of microwave technology for food application* (pp. 69-114). CRC Press.

Nicolás-García, M., Perucini-Avedaño, M., Jiménez-Martínez, C., Perea-Flores, M. d. J., Gómez-Patiño, M. B., Arrieta-Báez, D., & Dávila-Ortiz, G. (2021). Bean phenolic compound changes during processing: Chemical interactions and identification. *Journal of Food Science*, 86(3), 643-655.

Nongthombam, J., Kumar, A., Ladli, B., Madhushekhar, M., & Patidar, S. (2021). A review on study of growth and cultivation of oyster mushroom. *Plant Cell Biotechnology and Molecular Biology*, 22(5&6), 55-65.

Nour, A. H., Oluwaseun, A. R., Nour, A. H., Omer, M. S., & Ahmed, N. (2021). Microwave-assisted extraction of bioactive compounds. *Microwave heating. Electromagnetic fields causing thermal and non-thermal effects*, 1-31.

Nurkhasanah, A., Fardad, T., Carrera, C., Setyaningsih, W., & Palma, M. (2023). Ultrasound-Assisted Anthocyanins Extraction from Pigmented Corn: Optimization Using Response Surface Methodology. *Methods and Protocols*, 6(4), 69. <https://doi.org/https://doi.org/10.3390/mps6040069>

Oh, H.-I., Lim, T.-S., Lee, G.-D., & Kim, H.-K. (2006). Establishment of microwave-assisted extraction conditions for antioxidative extracts from *Pleurotus eryngii*. *Journal of The Korean Society of Food Science and Nutrition*, 35(1), 1-6.

<https://doi.org/https://doi.org/10.3746/jkfn.2006.35.1.001>

Osorio-Tobón, J. F. (2020). Recent advances and comparisons of conventional and alternative extraction techniques of phenolic compounds. *Journal of Food Science and Technology*, 57, 4299-4315.

<https://doi.org/https://doi.org/10.1007/s13197-020-04433-2>

Özyürek, M., Bener, M., Güçlü, K., & Apak, R. (2014). Antioxidant/antiradical properties of microwave-assisted extracts of three wild edible mushrooms.

Food Chemistry, 157, 323-331.

<https://doi.org/https://doi.org/10.1016/j.foodchem.2014.02.053>

Pereira, A. G., Cruz, L., Cassani, L., Chamorro, F., Lourenço-Lopes, C., Freitas, V., Otero, P., Fraga-Corral, M., Prieto, M. A., & Simal-Gandara, J. (2023). Comparative Study of Microwave-Assisted Extraction and Ultrasound-Assisted Extraction Techniques (MAE vs. UAE) for the Optimized Production of Enriched Extracts in Phenolic Compounds of *Camellia japonica* var *Eugenia* de Montijo. *Engineering Proceedings*, 37(1), 124.

<https://doi.org/https://doi.org/10.3390/ECP2023-14615>

Périno-Issartier, S., Abert-Vian, M., & Chemat, F. (2011). Solvent free microwave-assisted extraction of antioxidants from sea buckthorn (*Hippophae rhamnoides*) food by-products. *Food and Bioprocess Technology*, 4, 1020-1028.

- Pietrofesa, R. A., Velalopoulou, A., Arguiri, E., Menges, C. W., Testa, J. R., Hwang, W.-T., Albelda, S. M., & Christofidou-Solomidou, M. (2016). Flaxseed lignans enriched in secoisolariciresinol diglucoside prevent acute asbestos-induced peritoneal inflammation in mice. *Carcinogenesis*, *37*(2), 177-187.
- Proestos, C., Boziaris, I., Nychas, G.-J., & Komaitis, M. (2006). Analysis of flavonoids and phenolic acids in Greek aromatic plants: Investigation of their antioxidant capacity and antimicrobial activity. *Food Chemistry*, *95*(4), 664-671.
- Radzki, W., Tutaj, K., Skrzypczak, K., Michalak-Majewska, M., & Gustaw, W. (2023). Ethanolic Extracts of Six Cultivated Mushrooms as a Source of Bioactive Compounds. *Applied Sciences*, *14*(1), 66. <https://doi.org/https://doi.org/10.3390/app14010066>
- Rahman, M. M., Rahaman, M. S., Islam, M. R., Rahman, F., Mithi, F. M., Alqahtani, T., Almikhlaifi, M. A., Alghamdi, S. Q., Alruwaili, A. S., & Hossain, M. S. (2021). Role of phenolic compounds in human disease: current knowledge and future prospects. *Molecules*, *27*(1), 233. <https://doi.org/https://doi.org/10.3390/molecules27010233>
- Rasul, M. G. (2018). Conventional extraction methods use in medicinal plants, their advantages and disadvantages. *International Journal of Basic Sciences and Applied Computing*, *2*, 10-14.
- Reis, F. S., Barros, L., Martins, A., & Ferreira, I. C. (2012). Chemical composition and nutritional value of the most widely appreciated cultivated mushrooms:

An inter-species comparative study. *Food and Chemical Toxicology*, 50(2), 191-197.

Reis, F. S., Martins, A., Barros, L., & Ferreira, I. C. (2012). Antioxidant properties and phenolic profile of the most widely appreciated cultivated mushrooms: A comparative study between in vivo and in vitro samples. *Food and Chemical Toxicology*, 50(5), 1201-1207.
<https://doi.org/https://doi.org/10.1016/j.fct.2012.02.013>

Reungoat, V., Chadni, M., & Ioannou, I. (2021). Response Surface Methodology Applied to the Optimization of Phenolic Compound Extraction from Brassica. *IntechOpen*, 10.
<https://doi.org/https://doi.org/10.5772/intechopen.97655>.

Roselló-Soto, E., Parniakov, O., Deng, Q., Patras, A., Koubaa, M., Grimi, N., Boussetta, N., Tiwari, B. K., Vorobiev, E., & Lebovka, N. (2016). Application of non-conventional extraction methods: Toward a sustainable and green production of valuable compounds from mushrooms. *Food Engineering Reviews*, 8, 214-234.

Ruan, G.-H., & Li, G.-K. (2007). The study on the chromatographic fingerprint of *Fructus xanthii* by microwave assisted extraction coupled with GC-MS. *Journal of Chromatography B*, 850(1-2), 241-248.

Setyaningsih, W., Guamán-Balcázar, M. d. C., Oktaviani, N. M. D., & Palma, M. (2023). Response surface methodology optimization for analytical microwave-assisted extraction of resveratrol from functional marmalade and cookies. *Foods*, 12(2), 233.
<https://doi.org/https://doi.org/10.3390/foods12020233>

- Shams, K. A., Abdel-Azim, N. S., Saleh, I. A., Hegazy, M.-E. F., El-Missiry, M. M., Hammouda, F. M., Bohouth, E., & Tahrir, E. (2015). Green technology: Economically and environmentally innovative methods for extraction of medicinal & aromatic plants (MAP) in Egypt. *J. Chem. Pharm. Res*, 7(5), 1050-1074.
- Smeriglio, A., Barreca, D., Bellocco, E., & Trombetta, D. (2017). Proanthocyanidins and hydrolysable tannins: occurrence, dietary intake and pharmacological effects. *British journal of pharmacology*, 174(11), 1244-1262.
- Spigno, G., & De Faveri, D. M. (2009). Microwave-assisted extraction of tea phenols: A phenomenological study. *Journal of food engineering*, 93(2), 210-217.
- Taofiq, O., Rodrigues, F., Barros, L., Barreiro, M. F., Ferreira, I. C., & Oliveira, M. B. P. (2019). Mushroom ethanolic extracts as cosmeceuticals ingredients: Safety and ex vivo skin permeation studies. *Food and Chemical Toxicology*, 127, 228-236. <https://doi.org/https://doi.org/10.1016/j.fct.2019.03.045>
- Tatke, P., & Jaiswal, Y. (2011). An overview of microwave assisted extraction and its applications in herbal drug research.
- Tejedor-Calvo, E., García-Barreda, S., Sánchez, S., Morte, A., Siles-Sánchez, M. d. I. N., Soler-Rivas, C., Santoyo, S., & Marco, P. (2022). Application of pressurized liquid extractions to obtain bioactive compounds from *Tuber aestivum* and *Terfezia clavaryi*. *Foods*, 11(3), 298. <https://doi.org/https://doi.org/10.3390/foods11030298>

- Terigar, B., Balasubramanian, S., Boldor, D., Xu, Z., Lima, M., & Sabliov, C. (2010). Continuous microwave-assisted isoflavone extraction system: design and performance evaluation. *Bioresource technology*, *101*(7), 2466-2471.
- Thompson, M., Ellison, S. L., & Wood, R. (2002). Harmonized guidelines for single-laboratory validation of methods of analysis (IUPAC Technical Report). *Pure and applied chemistry*, *74*(5), 835-855.
- Törös, G. H., El-Ramady, H., & Prokisch, J. (2022). Edible mushroom of *Pleurotus* spp.: a case study of oyster mushroom (*Pleurotus ostreatus* L.). *Environment, Biodiversity and Soil Security*, *6*(2022), 51-59.
- Tsai, H.-Y., Ho, C.-T., & Chen, Y.-K. (2017). Biological actions and molecular effects of resveratrol, pterostilbene, and 3'-hydroxypterostilbene. *Journal of food and drug analysis*, *25*(1), 134-147.
- Ucar, T. M., & Karadag, A. (2019). The effects of vacuum and freeze-drying on the physicochemical properties and in vitro digestibility of phenolics in oyster mushroom (*Pleurotus ostreatus*). *Journal of Food Measurement and Characterization*, *13*, 2298-2309.
<https://doi.org/https://doi.org/10.1007/s11694-019-00149-w>
- Veggi, P. C., Santos, D. T., Fabiano-Tixier, A.-S., Bourvellec, C. I., Meireles, M. A. A., & Chemat, F. (2013). Ultrasound-assisted extraction of polyphenols from jatoba (*Hymenaea courbaril* L. var *stilbocarpa*) bark.
- Wang, L., & Weller, C. L. (2006). Recent advances in extraction of nutraceuticals from plants. *Trends in Food Science & Technology*, *17*(6), 300-312.

- Xia, N., Daiber, A., Förstermann, U., & Li, H. (2017). Antioxidant effects of resveratrol in the cardiovascular system. *British journal of pharmacology*, *174*(12), 1633-1646.
- Yasir, M., Sultana, B., & Amicucci, M. (2016). Biological activities of phenolic compounds extracted from Amaranthaceae plants and their LC/ESI-MS/MS profiling. *Journal of Functional Foods*, *26*, 645-656.
- Zhang, Q.-W., Lin, L.-G., & Ye, W.-C. (2018). Techniques for extraction and isolation of natural products: A comprehensive review. *Chinese medicine*, *13*, 1-26.
- Zhang, Y., Cai, P., Cheng, G., & Zhang, Y. (2022). A brief review of phenolic compounds identified from plants: Their extraction, analysis, and biological activity. *Natural Product Communications*, *17*(1), 1934578X211069721.
- Zhang, Z., Lv, G., Pan, H., & Fan, L. (2012). Optimisation of the microwave-assisted extraction process for six phenolic compounds in *Agaricus blazei murrill*. *International Journal of Food Science & Technology*, *47*(1), 24-31. <https://doi.org/https://doi.org/10.1111/j.1365-2621.2011.02802.x>
- Zin, M. M., Anucha, C. B., & Bánvölgyi, S. (2020). Recovery of phytochemicals via electromagnetic irradiation (microwave-assisted-extraction): Betalain and phenolic compounds in perspective. *Foods*, *9*(7), 918.