



DAFTAR PUSTAKA

- Aarti, P. D., Tanaka, R., & Tanaka, A. (2006). Effects of oxidative stress on chlorophyll biosynthesis in cucumber (*Cucumis sativus*) cotyledons. *Physiologia Plantarum*, 128(1), 186–197. <https://doi.org/10.1111/j.1399-3054.2006.00720.x>
- Albro, P. W., Corbett, J. T., & Schroeder, J. L. (1986). Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on lipid peroxidation in microsomal systems in vitro. *Chemico-Biological Interactions*, 57(3), 301–313. [https://doi.org/10.1016/0009-2797\(86\)90005-0](https://doi.org/10.1016/0009-2797(86)90005-0)
- Alexander, A. G. (1966). Oxidizing Enzymes of Sugarcane: Peroxidase. *The Journal of Agriculture of the University of Puerto Rico*, 50(1 SE-Articles), 36–52. <https://doi.org/10.46429/jaupr.v50i1.3440>
- Alfatemi, S. M. H., Sharifi-Rad, J. S., Rad, M. S., Mohsenzadeh, S., & da Silva, J. A. T. (2015). Chemical composition, antioxidant activity and in vitro antibacterial activity of Achillea wilhelmsii C. Koch essential oil on methicillin-susceptible and methicillin-resistant *Staphylococcus aureus* spp. *3 Biotech*, 5(1), 39–44. <https://doi.org/10.1007/s13205-014-0197-x>
- Almarie, A. A. A. (2020). *Roles of Terpenoids in Essential Oils and Its Potential as Natural Weed Killers: Recent Developments*. IntechOpen. <https://doi.org/10.5772/intechopen.91322>
- Ameena, M., Geethakumari, V. L., & George, S. (2014). Allelopathic influence of purple nutsedge (*Cyperus rotundus L.*) root exudates on germination and growth of important field crops. *Ann. Agric. Res. New Series*, 35(4), 362–366. <https://epubs.icar.org.in/index.php/AAR/article/view/46894>
- Ameena, M., Geethakumaril, V. L., & George, S. (2015). Allelopathic effects of root exudates of purple nutsedge (*Cyperus rotundus L.*) on growth of field crops. *Journal Crop and Weed*, 11(Special Issue), 142–145.
- Anaya, A. L. (1999). Allelopathy as a tool in the management of biotic resources in agroecosystems. In *Critical Reviews in Plant Sciences* (Vol. 18, Issue 6). <https://doi.org/10.1080/0735268991309450>
- Anh, L. H., Van Quan, N., Tuan Nghia, L., & Dang Xuan, T. (2021). Phenolic allelochemicals: Achievements, limitations, and prospective approaches in weed management. *Weed Biology and Management*, 21(2), 37–67. <https://doi.org/10.1111/WBM.12230>
- Araújo, C. A., Morgado, C. S. A., Gomes, A. K. C., Gomes, A. C. C., & Simas, N. K. (2021). Asteraceae family: A review of its allelopathic potential and the case of *Acmella oleracea* and *Sphagneticola trilobata*. *Rodriguesia*, 72. <https://doi.org/10.1590/2175-7860202172137>
- Arnon, D. I. (1949). Copper Enzymes in Isolated Chloroplasts. Polyphenoloxidase in Beta Vulgaris. *Plant Physiology*, 24(1), 1–15. <https://doi.org/10.1104/PP.24.1.1>
- Arruda, D. C., D'Alexandri, F. L., Katzin, A. M., & Uliana, S. R. B. (2005). Antileishmanial activity of the terpene nerolidol. *Antimicrobial Agents*



- and *Chemotherapy*, 49(5), 1679–1687.
<https://doi.org/10.1128/AAC.49.5.1679-1687.2005>
- Awouafack, M. D., Tane, P., Kuete, V., & Eloff, J. N. (2013). Sesquiterpenes from the Medicinal Plants of Africa. In *Medicinal Plant Research in Africa: Pharmacology and Chemistry* (pp. 33–103). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-405927-6.00002-3>
- Azizan, K. A., Baharum, S. N., & Noor, N. M. (2012). Metabolic profiling of *Lactococcus lactis* under different culture conditions. *Molecules*, 17(7), 8022–8036. <https://doi.org/10.3390/molecules17078022>
- Azizan, K. A., Ghani, N. H. A., & Nawawi, M. F. (2015). GC-MS based metabolomics and multivariate statistical analysis of *Wedelia trilobata* extracts for the identification of potential phytochemical properties. *POJ*, 8(6), 537–543.
- Azizan, K. A., Ibrahim, S., Ghani, N. H. A., & Nawawi, M. F. (2016). LC-MS Based Metabolomics Analysis to Identify Potential Allelochemicals in *Wedelia trilobata*. *Nat. Prod*, 10, 788–793. www.acgpubs.org/RNP
- Bajwa, A. A. (2014). Sustainable weed management in conservation agriculture. *Crop Protection*, 65, 105–113. <https://doi.org/10.1016/J.CROP.2014.07.014>
- Balekar, N., Nakpheng, T., & Srichana, T. (2014). *Wedelia trilobata L.*: A Phytochemical and Pharmacological Review. In *Chiang Mai J. Sci* (Vol. 41, Issue 3). <http://epg.science.cmu.ac.th/ejournal/>
- Batish, D. R., Singh, H. P., Setia, N., Kaur, S., & Kohli, R. K. (2006). 2-Benzoxazolinone (BOA) induced oxidative stress, lipid peroxidation and changes in some antioxidant enzyme activities in mung bean (*Phaseolus aureus*). *Plant Physiology and Biochemistry*, 44(11–12), 819–827. <https://doi.org/10.1016/j.plaphy.2006.10.014>
- Bendixen, L. E., & Nandihalli, U. B. (1987). Worldwide Distribution of Purple and Yellow Nutsedge (*Cyperus rotundus* and *C. esculentus*). *Technology*, 1(1), 61–65. <https://www.jstor.org/stable/3986985>
- Benvenuti, S., Cioni, P. L., Flamini, G., & Pardossi, A. (2017). Weeds for weed control: Asteraceae essential oils as natural herbicides. *Weed Research*, 57(5), 342–353. <https://doi.org/10.1111/wre.12266>
- Bharati, A., Kar, M., & Sabat, S. C. (2012). Artemisinin inhibits chloroplast electron transport activity: mode of action. *PloS One*, 7(6). <https://doi.org/10.1371/journal.pone.0038942>
- Bhattacharya, B., & Gupta, K. (1981). Steroid Hormone Effects on Growth and Apical Dominance of Sunflower. *Phytochemistry*, 20(5), 989–991.
- Bianchi, F., Careri, M., Mangia, A., & Musci, M. (2007). Retention indices in the analysis of food aroma volatile compounds in temperature-programmed gas chromatography: Database creation and evaluation of precision and robustness. *Journal of Separation Science*, 30(4), 563–572. <https://doi.org/10.1002/jssc.200600393>
- Bohm, W. (1979). Root Parameters and Their Measurement. In *Methods of Studying Root Systems* (Vol. 33, pp. 126–138). https://doi.org/https://doi.org/10.1007/978-3-642-67282-8_12



- Candido, L. P., Varela, R. M., Torres, A., Molinillo, J. M. G., Gualtieri, S. C. J., & Macías, F. A. (2016). Evaluation of the Allelopathic Potential of Leaf, Stem, and Root Extracts of *Ocotea pulchella* Nees et Mart. *Chemistry and Biodiversity*, 1058–1067. <https://doi.org/10.1002/cbdv.201500501>
- Cantillo-Ciau, Z., Moo-Puc, R., Quijano, L., & Freile-Pelegrín, Y. (2010). The tropical brown alga *Lobophora variegata*: A source of antiprotozoal compounds. *Marine Drugs*, 8(4), 1292–1304. <https://doi.org/10.3390/md8041292>
- Carvalho, M. H. C. de. (2008). Drought stress and reactive oxygen species. *Plant Signaling & Behavior*, 3, 156–5. www.landesbioscience.com/journals/psb/article/5537
- Chaïb, S., Pistevos, J. C. A., Bertrand, C., & Bonnard, I. (2021a). Allelopathy and allelochemicals from microalgae: An innovative source for bio-herbicidal compounds and biocontrol research. *Algal Research*, 54, 102213. <https://doi.org/10.1016/J.AL GAL.2021.102213>
- Chaïb, S., Pistevos, J. C. A., Bertrand, C., & Bonnard, I. (2021b). Allelopathy and allelochemicals from microalgae: An innovative source for bio-herbicidal compounds and biocontrol research. In *Algal Research* (Vol. 54). Elsevier B.V. <https://doi.org/10.1016/j.algal.2021.102213>
- Chaimovitsh, D., Rogovoy Stelmakh, O., Altshuler, O., Belausov, E., Abu-Abied, M., Rubin, B., Sadot, E., & Dudai, N. (2012). The relative effect of citral on mitotic microtubules in wheat roots and BY2 cells. *Plant Biology*, 14(2), 354–364. <https://doi.org/10.1111/j.1438-8677.2011.00511.x>
- Chand, M., Lal, R., Khippal, A., & Singh, S. (20110). Integrated weed management in sugarcane ratoon. *Indian Journal of Sugarcane Technology*, 25(1 & 2), 17–19. <https://www.researchgate.net/publication/327532027>
- Cheema, Z. A., Farooq, M., & Wahid, A. (2013). Allelopathy: Current trends and future applications. In *Allelopathy: Current Trends and Future Applications*. Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-642-30595-5/COVER>
- Chen', P. K., Polatnick, M., & Leather, G. (1991). Comparative Study on Artemisinin, 2,4-D, and Glyphosate. *J. Agric. Food Chem*, 39, 991–994.
- Choudhuri, S., & Chanderbhan, R. F. (2016). The biology of nutrients: Genetic and molecular principles. In *Nutraceuticals: Efficacy, Safety and Toxicity* (pp. 209–225). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-802147-7.00017-6>
- Daley, S. L., & Hassell, R. L. (2014). Fatty Alcohol Application to Control Meristematic Regrowth in Bottle Gourd and Interspecific Hybrid Squash Rootstocks Used for Grafting Watermelon. *HORTSCIENCE*, 49(3), 260–264. <https://doi.org/https://doi.org/10.21273/HORTSCI.49.3.260>
- DellaGreca, M., Zarrelli, A., Fergola, P., Cerasuolo, M., Pollio, A., & Pinto, G. (2010). Fatty acids released by *Chlorella vulgaris* and Their Role in Interference with *Pseudokirchneriella subcapitata*: Experiments and



- Modelling. *Journal of Chemical Ecology*, 36(3), 339–349. <https://doi.org/10.1007/s10886-010-9753-y>
- Dobrescu, A., Giuffrida, M. V., & Tsafaris, S. A. (2017). Leveraging multiple datasets for deep leaf counting. *Proceedings - 2017 IEEE International Conference on Computer Vision Workshops, ICCVW 2017, 2018-January*, 2072–2079. <https://doi.org/10.1109/ICCVW.2017.243>
- Domonkos, I., Kis, M., Gombos, Z., & Ughy, B. (2013). Carotenoids, versatile components of oxygenic photosynthesis. In *Progress in Lipid Research* (Vol. 52, Issue 4, pp. 539–561). <https://doi.org/10.1016/j.plipres.2013.07.001>
- Doorandishan, M., Gholami, M., Ebrahimi, P., & Jassbi, A. R. (2021). Spathulenol as the most abundant component of essential oil of moluccella aucheri (Boiss.) scheen. *Natural Volatiles and Essential Oils*, 8(2), 37–41. <https://doi.org/10.37929/nveo.817562>
- Dorning, M., & Cipollini, D. (2006). Leaf and root extracts of the invasive shrub, *Lonicera maackii*, inhibit seed germination of three herbs with no autotoxic effects. *Plant Ecology*, 184(2), 287–296. <https://doi.org/10.1007/s11258-005-9073-4>
- El Sawi, S. A., Ibrahim, M. E., El-Rokiek, K. G., & El-Din, S. A. S. (2019). Allelopathic potential of essential oils isolated from peels of three citrus species. *Annals of Agricultural Sciences*, 64(1), 89–94. <https://doi.org/10.1016/j.aoas.2019.04.003>
- El-Amier, Y. A., & Abdullah, T. J. (2014). Allelopathic Effect of Four Wild Species on Germination and Seedling Growth of *Echinocloa crusgalli* (L.) P. Beauv. Article in *International Journal of Advanced Research*, 2(9), 287–294.
- El-Darier, S. M., Abdelaziz, H. A., & Zein El-Dien, M. H. (2014). Effect of soil type on the allelotoxic activity of *Medicago sativa* L. residues in *Vicia faba* L. agroecosystems. *Journal of Taibah University for Science*, 8(2), 84–89. <https://doi.org/10.1016/j.jtusci.2014.01.001>
- El-Rokiek, Kowthar, G., Rafat, R. E., Nadia K.M., & Salah A.A. (2010). The Allelopathic Effect of Mango Leaves on the Growth and Propagative Capacity of Purple Nutsedge (*Cyperus rotundus* L.). *Journal of American Science*, 6(9), 151–159. <http://www.americanscience.orgeditor@americanscience.org>
- Entzeroth, M., Mead, D. J., Pa-Iterson, G. M. L., & Moore, R. E. (1985). A HERBICIDAL FATTY ACID PRODUCED BY LYZVGBYA AESTUARZZ (Vol. 24, Issue 12).
- Ferdous, J., Mamun, M., & Sarafat Ali, Md. (2017). Impact of Environmental and Stress Factors on the Photosynthetic Capabilities of Plants. *Universal Journal of Agricultural Research*, 5(2), 113–118. <https://doi.org/10.13189/ujar.2017.050205>
- Fischer, N. H., & Quijano, L. (1985). Allelopathic Agents from Common Weeds. In *The Chemistry of Allelopathy* (pp. 133–147). American Chemical Society. <https://doi.org/10.1021/bk-1985-0268.ch009>



- Fischer, N. H., Williamson, G. B., Tanrisever, N., Pena, A. D. L., Weidenhamer, J. D., Jordan, E. D., & Richardson, D. R. (1989). Allelopathic Actions in the Florida Scrub Community. In *BIOLOGIA PLANTARUM (PRAHA)* (Vol. 31, Issue 6).
- Foyer C.H., N. G. (2005). Redox Homeostasis and Antioxidant Signaling: A Metabolic Interface between Stress Perception and Physiological Responses. *Plant Cell*, 17(7), 1866–1875. www.plantcell.org
- Fu, Y. H., Quan, W. X., Li, C. C., Qian, C. Y., Tang, F. H., & Chen, X. J. (2019). Allelopathic effects of phenolic acids on seedling growth and photosynthesis in rhododendron delavayi franch. *Photosynthetica*, 57(2), 377–387. <https://doi.org/10.32615/ps.2019.045>
- Gharde, Y., Singh, P. K., Dubey, R. P., & Gupta, P. K. (2018). Assessment of yield and economic losses in agriculture due to weeds in India. *Crop Protection*, 107, 12–18. <https://doi.org/10.1016/j.croppro.2018.01.007>
- Gill, S. S., Nafees, K. A., Naser, A. A., & Narendra Tuteja. (2011). Amelioration of Cadmium Stress in Crop Plants by Nutrient Management: Morphological, Physiological and Biochemical Aspects. *Plant Stress*, 5(1), 1–15. <https://api.semanticscholar.org/CorpusID:40009775>
- Gill, S. S., & Tuteja, N. (2010). Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry*, 48(12), 909–930. <https://doi.org/10.1016/J.PLAPHY.2010.08.016>
- Gómez-Sagasti, M. T., López-Pozo, M., Artetxe, U., Becerril, J. M., Hernández, A., García-Plazaola, J. I., & Esteban, R. (2023). Carotenoids and their derivatives: A “Swiss Army knife-like” multifunctional tool for fine-tuning plant-environment interactions. In *Environmental and Experimental Botany* (Vol. 207). Elsevier B.V. <https://doi.org/10.1016/j.envexpbot.2023.105229>
- Grassmann, J., & Elstner, E. F. (2003). *ESSENTIAL OILS Contents Properties and Uses Isolation and Production Properties and Uses*.
- Hambalko, J., Gajdoš, P., Nicaud, J. M., Ledesma-Amaro, R., Tupec, M., Pichová, I., & Čertík, M. (2021). Production of Long Chain Fatty Alcohols Found in Bumblebee Pheromones by *Yarrowia lipolytica*. *Frontiers in Bioengineering and Biotechnology*, 8. <https://doi.org/10.3389/fbioe.2020.593419>
- Harborne, J. Barry., & Tomas-Barberan, F. A. . (1991). *Ecological chemistry and biochemistry of plant terpenoids: International symposium : Papers*. Clarendon.
- Harris, R. W. (1992). ROOT-SHOOT RATIOS. In *Journal of Arboriculture* (Vol. 18, Issue 1).
- Hasan, M., Ahmad-Hamdani, M. S., Rosli, A. M., & Hamdan, H. (2021). Bioherbicides: An eco-friendly tool for sustainable weed management. In *Plants* (Vol. 10, Issue 6). MDPI AG. <https://doi.org/10.3390/plants10061212>



- Hasanuzzaman, M., Bhuyan, M. H. M. B., Zulfiqar, F., Raza, A., Mohsin, S. M., Al Mahmud, J., Fujita, M., & Fotopoulos, V. (2020). Reactive oxygen species and antioxidant defense in plants under abiotic stress: Revisiting the crucial role of a universal defense regulator. In *Antioxidants* (Vol. 9, Issue 8, pp. 1–52). MDPI. <https://doi.org/10.3390/antiox9080681>
- Hazrati, H., Saharkhiz, M. J., Niakousari, M., & Moein, M. (2017). Natural herbicide activity of Satureja hortensis L. essential oil nanoemulsion on the seed germination and morphophysiological features of two important weed species. *Ecotoxicology and Environmental Safety*, 142, 423–430. <https://doi.org/10.1016/j.ecoenv.2017.04.041>
- He, J., Reddy, G. V. P., Liu, M., & Shi, P. (2020). A general formula for calculating surface area of the similarly shaped leaves: Evidence from six Magnoliaceae species. *Global Ecology and Conservation*, 23. <https://doi.org/10.1016/j.gecco.2020.e01129>
- Heftmann, E. (1975). Functions of Steroid in Plants. In *Phytochemistry* (Vol. 14). Pergamon Press. Printed in England.
- Hidayatullah, N., Hoesain, M., Hasjim, D. S., Agroteknologi, S., Pertanian, F., & Jember, U. (2019). Effect of Bamboo Leaves Extract Concentration (*Dendrocalamus sasper L.*) as Bioherbicide on Nutsedge (*Cyperus rotundus L.*). *International Journal Of Science, Engineering, And Information Technology*, 03(02).
- Hoagland, R. E., Boyette, C. D., Weaver, M. A., & Abbas, H. K. (2007). Bioherbicides: Research and risks. *Toxin Reviews*, 26(4), 313–342. <https://doi.org/10.1080/15569540701603991>
- Horowitz, M. (1992). Mechanisms of establishment and spreading of *Cyperus rotundus*-the worst weed of warm regions. *Proceedings of the First International Weed Control Congress*.
- Hosni, K., Hassen, I., Sebei, H., & Casabianca, H. (2013). Secondary metabolites from *Chrysanthemum coronarium* (Garland) flowerheads: Chemical composition and biological activities. *Industrial Crops and Products*, 44, 263–271. <https://doi.org/10.1016/J.INDCROP.2012.11.033>
- Housley, T. L., & Peterson, D. M. (1982). Oat Stem Vascular Size in Relation to Kernel Number and Weight. I. Controlled Environment 1. *Crop Science*, 22(2), 259–263. <https://doi.org/10.2135/cropsci1982.0011183x002200020014x>
- Hunt, R. (2016). Growth Analysis, Individual Plants. In *Encyclopedia of Applied Plant Sciences* (Vol. 1, pp. 421–429). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-394807-6.00226-4>
- Hussain, I., Singh, N. B., Singh, A., & Singh, H. (2017). Allelopathic potential of sesame plant leachate against *Cyperus rotundus L.* *Annals of Agrarian Science*, 15(1), 141–147. <https://doi.org/10.1016/j.aasci.2016.10.003>
- Ikawa, M. (2004). Algal polyunsaturated fatty acids and effects on plankton ecology Algal polyunsaturated fatty acids and effects on plankton



ecology and other organisms and other organisms Algal polyunsaturated fatty acids and effects on plankton ecology and other organisms. In *UNH Center Freshwat. Biol. Res* (Vol. 6, Issue 2). <https://scholars.unh.edu/cfb/10>

Ikawa, M., Hartshorne, T., Caron, L.-A., Iannitelli, R. C., Barbero, L. J., & Wegener, K. (1984). Inhibition of Growth of the Green Alga Chlorella pyrenoidosa by Unsaturated Fatty Acids. *J Am Oil Chem Soc*, 61, 1877–1878. [https://doi.org/https://doi.org/10.1007/BF02540823](https://doi.org/10.1007/BF02540823)

Jabran, K., Mahajan, G., Sardana, V., & Chauhan, B. S. (2015). Allelopathy for weed control in agricultural systems. *Crop Protection*, 72, 57–65. <https://doi.org/10.1016/J.CROP.2015.03.004>

Jasim, I., Alwattar, M., & Yaqub, H. (2023). Terpenoids as Natural Allelopathic Compounds in Plants. *Rafidain Journal of Science*, 32(4), 106–116. <https://doi.org/10.33899/rjs.2023.181275>

Kaur, I., & Sharma, R. (2016). Allelopathic Effect of Ageratum Conyzoides on Chlorophyll Content in The Leaves of Mungbean. *International Journal of Recent Scientific Research*, 7(9), 13296–13298. <http://www.recentscientific.com>

Kaur, S., Singh, H. P., Mittal, S., Batish, D. R., & Kohli, R. K. (2010). *Phytotoxic effects of volatile oil from Artemisia scoparia against weeds and its possible use as a bioherbicide*. 32(1), 54–61. <https://doi.org/https://doi.org/10.1016/j.indcrop.2010.03.007>

Khursheed, A., Rather, M. A., Jain, V., Wani, A. R., Rasool, S., Nazir, R., Malik, N. A., & Majid, S. A. (2022). Plant based natural products as potential ecofriendly and safer biopesticides: A comprehensive overview of their advantages over conventional pesticides, limitations and regulatory aspects. *Microbial Pathogenesis*, 173. <https://doi.org/10.1016/j.micpath.2022.105854>

Kiri, I. Z. (2023). A Review on Plant Root Architecture and Methods for Measuring Root Growth Parameters. *Dutse Journal of Pure and Applied Sciences*, 9(1a), 57–66. <https://doi.org/10.4314/dujopas.v9i1a.6>

Kokpol, U., Chavasiri, W., Chittawong, V., Bruce, M., Cunningham, G. N., & Miles, D. H. (1993). Long Chain Aliphatic Alcohols and Saturated Carboxylic Acids from Heartwood of Rhizophora Apiculata. *Phytochemistry*, 33(5), 1129–1131.

Kostina-Bednarz, M., Płonka, J., & Barchanska, H. (2023). Allelopathy as a source of bioherbicides: challenges and prospects for sustainable agriculture. *Reviews in Environmental Science and Biotechnology*, 22(2), 471–504. <https://doi.org/10.1007/s11157-023-09656-1>

Kpoviessi, D. S. S., Gdaguidi, F. A., Gbenou, J. D., Accrombessi, G. C., Haddad, M., Moudachirou, M., & Quetin-Leclercq, J. (2006). Allelopathic effects on cowpea (*Vigna unguiculata* (L.) Walp) plant and cytotoxic activities of sterols and triterpene isolated from *Justicia anselliana* (NEES) T. Anders. *Electronic Journal of Natural Substances*, 1, 12–19. <http://ejns.univ-lyon1.fr>



- Lawal, O. A., & Oyedele, A. O. (2009). molecules Chemical Composition of the Essential Oils of *Cyperus rotundus L.* from South Africa. *Molecules*, 14, 2909–2917. <https://doi.org/10.3390/molecules14082909>
- Li, D., Liang, Z., Guo, M., Zhou, J., Yang, X., & Xu, J. (2012). Study on the chemical composition and extraction technology optimization of essential oil from *Wedelia trilobata* (L.) Hitchc. *African Journal of Biotechnology*, 11(20). <https://doi.org/10.5897/ajb11.3198>
- Lichtenthaler, H. K. (1987). [34] Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methods in Enzymology*, 148(C), 350–382. [https://doi.org/10.1016/0076-6879\(87\)48036-1](https://doi.org/10.1016/0076-6879(87)48036-1)
- Lobón, N. C., De La Cruz, I. F., & Gallego, J. C. A. (2019). Autotoxicity of diterpenes present in leaves of *cistus ladanifer* L. *Plants*, 8(2). <https://doi.org/10.3390/plants8020027>
- Long, A., Zhang, J., Yang, L. T., Ye, X., Lai, N. W., Tan, L. L., Lin, D., & Chen, L. S. (2017). Effects of low pH on photosynthesis, related physiological parameters, and nutrient profiles of citrus. *Frontiers in Plant Science*, 8. <https://doi.org/10.3389/fpls.2017.00185>
- Macias, F. A., Simonet, A. M., & Galindo, J. C. G. (1997). BIOACTIVE STEROIDS AND TRITERPENES FROM *Melilotus messanensis* AND THEIR ALLELOPATHIC POTENTIAL1. *Journal of Chemical Ecology*, 23(7).
- Mahan, J. R., Dotray, P. A., & Light, G. G. (2004). Thermal dependence of enzyme function and inhibition; implications for, herbicide efficacy and tolerance. *PHYSIOLOGIA PLANTARUM*, 120(2), 187–195. <https://doi.org/10.1111/j.0031-9317.2004.0255.x>
- Maoka, T. (2020). Carotenoids as natural functional pigments. In *Journal of Natural Medicines* (Vol. 74, Issue 1). Springer. <https://doi.org/10.1007/s11418-019-01364-x>
- Matloob, A., Shareef, M. N., Khaliq, A., & Farooq, M. (2010). Quantification of allelopathic potential of different crop residues for the purple nutsedge suppression Zahid Ata Cheema. *Pak J Weed Sci*, 16(1), 1–12. <https://www.researchgate.net/publication/303242850>
- Mérillon, J.-M., & Ramawat, K. G. (2020). *Co-Evolution of Secondary Metabolites* (J.-M. Mérillon & K. G. Ramawat, Eds.). Springer Nature Switzerland AG. <https://doi.org/https://doi.org/10.1007/978-3-319-96397-6>
- Messiha, N. K., Ahmed, S. A., El-Rokiek, K. G., Dawood, M. G., & El-Masry, R. R. (2013). The physiological influence of allelochemicals in two Brassicaceae plant seeds on the growth and propagative capacity of *Cyperus rotundus* and *Zea mays* L. *World Applied Sciences Journal*, 26(9), 1142–1149. <https://doi.org/10.5829/idosi.wasj.2013.26.09.13548>
- Miao, C., Yang, S., Xu, J., Wang, H., Zhang, Y., Cui, J., Zhang, H., Jin, H., Lu, P., He, L., Yu, J., Zhou, Q., & Ding, X. (2023). Effects of Light Intensity on Growth and Quality of Lettuce and Spinach Cultivars in a Plant Factory. *Plants*, 12(18). <https://doi.org/10.3390/plants12183337>



- Miller, G., Suzuki, N., Ciftci-Yilmaz, S., & Mittler, R. (2010). Reactive oxygen species homeostasis and signalling during drought and salinity stresses. *Plant, Cell and Environment*, 33(4), 453–467. <https://doi.org/10.1111/j.1365-3040.2009.02041.x>
- Miller, G. W., Sasse, J. M., Lovelace, J., & Rowan, K. S. (1984a). Effects of Podolactone-Type Inhibitors and Abscisic Acid on Chlorophyll Biosynthesis in Barley Leaves. *Plant and Cell Physiology*, 25(4), 635–642. <https://doi.org/https://doi.org/10.1093/oxfordjournals.pcp.a076754>
- Miller, G. W., Sasse, J. M., Lovelace, J., & Rowan, K. S. (1984b). Effects of Podolactone-Type Inhibitors and Abscisic Acid on Chlorophyll Biosynthesis in Barley Leaves. In *Plant & Cell Physiol* (Vol. 25, Issue 4).
- Mittler, R. (2002). Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci*, 7, 405. [https://doi.org/doi:10.1016/s1360-1385\(02\)02312-9](https://doi.org/doi:10.1016/s1360-1385(02)02312-9)
- Molisch, H. (1938). Der Einfluss einer Pflanze auf die Andere, Allelopathie (Pengaruh suatu tumbuhan terhadap tumbuhan lain, alelopati). In *Nature* (Vol. 141). <https://doi.org/10.1038/141493a0>
- Molisch, H. (2001). *The influence of one plant on another: allelopathy*. (S. S. Narwal, Ed.). Scientific Publisheris (India). <https://www.researchgate.net/publication/322466455>
- Morales-Payan, J. P., Stall, W. M., Shilling, D. G., Charudattan, R., Dusky, J. A., & Bewick, T. A. (2003). Above- and Belowground Interference of Purple and Yellow Nutsedge (*Cyperus* spp.) with Tomato. *Weed Science*, 51(2), 181–185. [https://doi.org/doi:10.1614/0043-1745\(2003\)051\[0181:AABIOP\]2.0.CO;2](https://doi.org/doi:10.1614/0043-1745(2003)051[0181:AABIOP]2.0.CO;2)
- Muller-Rober', B., Sonnewald, U., & Willmitzer, L. (1992). Inhibition of the ADP-glucose pyrophosphorylase in transgenic potatoes leads to sugar-storing tubers and influences tuber formation and expression of tuber storage protein genes. In *The EMBO Journal* (Vol. 1, Issue 4).
- Nie, C., Luo, S., Zeng, R., Mo, M., Li, H., & Lin, C. (2005). Allelopathic potential of *Wedelia trilobata* L.: effects on germination, growth and physiological parameters of rice. *The Regional Institute* . http://www.regional.org.au/au/allelopathy/2005/2/3/2293_chengrongn.htm?print=1
- Nievola, C. C., Carvalho, C. P., Carvalho, V., & Rodrigues, E. (2017). Rapid responses of plants to temperature changes. *Temperature*, 4(4), 371–405. <https://doi.org/10.1080/23328940.2017.1377812>
- Ninkuu, V., Zhang, L., Yan, J., Fu, Z., Yang, T., & Zeng, H. (2021). Biochemistry of terpenes and recent advances in plant protection. In *International Journal of Molecular Sciences* (Vol. 22, Issue 11). MDPI. <https://doi.org/10.3390/ijms22115710>
- Nooshahi, H. A., Khan, A. H., Nooshahi, U. F., Hussain, M., Javed, T., Zafar, M., Batool, M., Ahmed, U., Liu, K., Harrison, M. T., Saud, S., Fahad, S., & Shu, S. (2022). Biosynthetic pathways of triterpenoids and strategies



- to improve their Biosynthetic Efficiency. In *Plant Growth Regulation* (Vol. 97, Issue 3, pp. 439–454). Springer Science and Business Media B.V. <https://doi.org/10.1007/s10725-022-00818-9>
- Nugroho, I. F., Sugito, Y., & Widaryanto, E. (2018). Pengaruh Kerapatan Gulma Teki (*Cyperus rotundus L.*) terhadap Tanaman Buncis Tegak (*Phaseolus vulgaris L.*). *Jurnal Produksi Tanaman*, 6(4), 649–654.
- Oerke, E. C. (2006). Crop losses to pests. *Journal of Agricultural Science*, 144(1), 31–43. <https://doi.org/10.1017/S0021859605005708>
- Ohkawa, M., & Nishikawa, Y. (1987). Plant Growth Inhibitory Activity of Fatty Acids and The Related Compounds by The Avena Coleoptile Test. *Plant Science*, 53(1), 35–38. [https://doi.org/10.1016/0168-9452\(87\)90175-0](https://doi.org/10.1016/0168-9452(87)90175-0)
- Pacanoski, Z., & Mehmeti, A. (2023). Herbicides weed management in changing environmental conditions. *Acta Agriculturae Slovenica*, 119(4). <https://doi.org/10.14720/aas.2023.119.4.12497>
- Parsons, W. T., Cuthbertson, E. G., & CSIRO. (2001). *Noxious weeds of Australia*. CSIRO Publishing. <https://nla.gov.au/nla.cat-vn2521594>
- Peerzada, A. M. (2017). Biology, agricultural impact, and management of *Cyperus rotundus L.*: the world's most tenacious weed. *Acta Physiologiae Plantarum*, 39(12). <https://doi.org/10.1007/s11738-017-2574-7>
- Penuelas, J., Ribas-Carbo, M., & Giles, L. (1996). Effects of Allelochemicals on Plant Respiration and Oxygen Isotope Fractionation by The Alternative Oxsidase. *Journal of Chemical Ecology*, 22(4).
- Perveen, S. (2018). Introductory Chapter: Terpenes and Terpenoids. In *Terpenes and Terpenoids*. IntechOpen. <https://doi.org/10.5772/intechopen.79683>
- Qi, Y., Wei, W., Chen, C., & Chen, L. (2019). Plant root-shoot biomass allocation over diverse biomes: A global synthesis. *Global Ecology and Conservation*, 18. <https://doi.org/10.1016/j.gecco.2019.e00606>
- Qian, H., Xu, J., Lu, T., Zhang, Q., Qu, Q., Yang, Z., & Pan, X. (2018). Responses of unicellular alga *Chlorella pyrenoidosa* to allelochemical linoleic acid. *Science of the Total Environment*, 625, 1415–1422. <https://doi.org/10.1016/j.scitotenv.2018.01.053>
- Qian, H., Xu, X., Chen, W., Jiang, H., Jin, Y., Liu, W., & Fu, Z. (2009). Allelochemical stress causes oxidative damage and inhibition of photosynthesis in *Chlorella vulgaris*. *Chemosphere*, 75(3), 368–375. <https://doi.org/10.1016/J.CHEMOSPHERE.2008.12.040>
- Rabbani, N., Bajwa, R., & Javaid, A. (2011). Interference of five problematic weed species with rice growth and yield. *African Journal of Biotechnology*, 10(10), 1854–1862. <https://doi.org/10.5897/AJB10.1866>
- Radhakrishnan, R., Alqarawi, A. A., & Abd Allah, E. F. (2018). Bioherbicides: Current knowledge on weed control mechanism. *Ecotoxicology and Environmental Safety*, 158, 131–138. <https://doi.org/10.1016/j.ecoenv.2018.04.018>



- Respatie, D. W., Yudono, P., Purwantoro, A., & Andi Trisyono, Y. (2019). The potential of *Cosmos sulphureus* Cav. extracts as a natural herbicides. *AIP Conference Proceedings*, 2202. <https://doi.org/10.1063/1.5141690>
- Rice, E. L. (1983). *Allelopathy* (2nd ed.). Elsevier Science, Academic Press.
- Sánchez-Moreiras, A. M., & Reigosa, M. J. (2005). Whole plant response of lettuce after root exposure to BOA (2(3H)-benzoxazolinone). *Journal of Chemical Ecology*, 31(11), 2689–2703. <https://doi.org/10.1007/s10886-005-7620-z>
- Santos, B. M., Morales-Payan, J. P., Stall, W. M., Bewick, T. A., & Shilling, D. G. (1997). Effects of Shading on the Growth of Nutsedges (*Cyperus* spp.). *Weed Science*, 45(5), 670–673. <https://www.jstor.org/stable/4045892>
- Sasamoto, H., Suzuki, S., Mardani-Korranı, H., Sasamoto, Y., & Fujii, Y. (2021). Allelopathic activities of three carotenoids, neoxanthin, crocin and β-carotene, assayed using protoplast co-culture method with digital image analysis. *Plant Biotechnology*, 38(1), 101–107. <https://doi.org/10.5511/plantbiotechnology.20.1211a>
- Sasse, J. M., Cerana, R., & Colombo, R. (1984). The effects of podolactone-type inhibitors on fusicoccin-induced growth and proton efflux. *Physiol. Plant*, 303–308.
- Sasse, J. M., Wardrop, J. J., Rowan, I., Aspinall, D., Coombe, G., Paleg, L. G., George Buta Sasse, I., Sasse, J. M., Wardrop, J. J., & Rowan, K. S. (1982). Some physiological effects of podolactone-type inhibitors. *Physiologia Plantarum*, 55(1). <https://doi.org/https://doi.org/10.1111/j.1399-3054.1982.tb00284.x>
- Schroeder, J., Thomas, S. H., & Murray, L. (1993). *Yellow and Purple Nutsedge and Chile Peppers Host Southern Root-Knot Nematode*. 41(1), 150–156. <https://about.jstor.org/terms>
- Scognamiglio, M., D'Abrosca, B., Esposito, A., Pacifico, S., Monaco, P., & Fiorentino, A. (2013). Plant growth inhibitors: Allelopathic role or phytotoxic effects? Focus on Mediterranean biomes. In *Phytochemistry Reviews* (Vol. 12, Issue 4, pp. 803–830). Kluwer Academic Publishers. <https://doi.org/10.1007/s11101-013-9281-9>
- Senthilkumar, M., Amaresan, N., & Sankaranarayanan, A. (2021). *Estimation of Malondialdehyde (MDA) by Thiobarbituric Acid (TBA) Assay*. 103–105. https://doi.org/10.1007/978-1-0716-1080-0_25
- Shahena, S., Rajan, M., Chandran, V., & Mathew, L. (2021). Allelopathic effect of *Wedelia trilobata* L., on the germination and growth of *Cicer arietinum*, *Vigna unguiculata*, and *Vigna radiata* seedlings. *Journal of Applied Biology & Biotechnology*, 9(2), 93–114. <https://doi.org/10.7324/JABB.2021.9209>
- Sharifi-Rad, J., Hoseini-Alfatemi, S. M., Sharifi-Rad, M., & Teixeira da Silva, J. A. (2015). Antibacterial, antioxidant, antifungal and anti-inflammatory activities of crude extract from *Nitraria schoberi* fruits. *3 Biotech*, 5(5), 677–684. <https://doi.org/10.1007/s13205-014-0266-1>



- Singh, H., Chawla, G., Rowe, J. W., Toda, J. K., RowE, W., Bower, C. L., & WAaNER, E. R. (1970). Waxes and Sterols od *Erythrina suberosa* Bark. *Proc. Ind. Acad. Sci.*, 9, 494.
- Singh, K. D., Mobolade, A. J., Bharali, R., Sahoo, D., & Rajashekhar, Y. (2021). Main plant volatiles as stored grain pest management approach: A review. *Journal of Agriculture and Food Research*, 4, 1–12. <https://doi.org/10.1016/j.jafr.2021.100127>
- Singh, N. B., Pandey, B. N., & Singh, A. (2009). Allelopathic effects of *cyperus rotundus* extract in vitro and ex vitro on banana. *Acta Physiologiae Plantarum*, 31(3), 633–638. <https://doi.org/10.1007/s11738-009-0274-7>
- Singh, S., Apoorva, Saha, P., Rai, N., Kumari, S., & Pandey-Rai, S. (2023). Unravelling triterpenoid biosynthesis in plants for applications in bioengineering and large-scale sustainable production. In *Industrial Crops and Products* (Vol. 199). Elsevier B.V. <https://doi.org/10.1016/j.indcrop.2023.116789>
- Singh, S. K., Singh Chhokar, R., Singh, R. G., & Ladha, J. K. (2009). Integrated weed management: a key to success for direct-seeded rice in the Indo-Gangetic Plains. In *Agricultural and Food Sciences, Environmental Science* (pp. 261–277). <https://www.researchgate.net/publication/285280264>
- Soledade, M., Pedras, C., & Zheng, Q. (2010). The Chemistry of *Arabidopsis thaliana*. In Hung-Wen (Ben) Liu & Lew Mander (Eds.), *Comprehensive Natural Products II* (pp. 1297–1315). Elsevier. <https://doi.org/https://doi.org/10.1016/B978-008045382-8.00090-3>
- Sozer, O., Komenda, J., Ughy, B., Domonkos, I., Laczk-Dobos, H., Malec, P., Gombos, Z., & Kis, M. (2010). Involvement of carotenoids in the synthesis and assembly of protein subunits of photosynthetic reaction centers of *synechocystis* sp. PCC 6803. *Plant and Cell Physiology*, 51(5), 823–835. <https://doi.org/10.1093/pcp/pcq031>
- Srivastava, R. K., Singh, A., & Shukla, S. V. (2013). Chemical Investigation and Pharmaceutical Action of *Cyperus rotundus* - A Review. *Journal of Biologically Active Products from Nature*, 3(3), 166–172. <https://doi.org/10.1080/22311866.2013.833381>
- Stoller, E. W., Sweet, R. D., & Sweet2, R. D. (1987). Biology and Life Cycle of Purple and Yellow Nutsedges (*Cyperus rotundus* and *C. esculentus*) Biology and Life Cycle of Purple and Yellow Nutsedges (*Cyperus rotundus* and *C. esculentus*). *Technology*, 1(1), 66–73. <https://www.jstor.org/stable/3986986>
- Sudhakar, P., Latha, P., & Reddy, P. V. (2016). Photosynthetic rates. In *Phenotyping Crop Plants for Physiological and Biochemical Traits* (pp. 33–39). Elsevier. <https://doi.org/10.1016/b978-0-12-804073-7.00004-1>
- Suttle, J. C., Olson, L. L., & Lulai, E. C. (2016). The Involvement of Gibberellins in 1,8-Cineole-Mediated Inhibition of Sprout Growth in Russet Burbank Tubers. *American Journal of Potato Research*, 93(1), 72–79. <https://doi.org/10.1007/s12230-015-9490-4>



- Talukder, M. R., Asaduzzaman, M., Ueno, M., Tanaka, H., & Asao, T. (2020). Alleviation of allelochemical stress-induced growth inhibition and oxidative damage in lettuce under closed hydroponics through electro-degradation. *Horticultural Science*, 47(1), 53–68. <https://doi.org/10.17221/32/2019-HORTSCI>
- Telfer, A., Bollman, K. M., & Poethig, R. S. (1997). Phase change and the regulation of trichome distribution in *Arabidopsis thaliana*. *Development*, 124(3), 645–654. <https://doi.org/https://doi.org/10.1242/dev.124.3.645>
- Thullen, R. J., & Keeley, P. E. (1979). Seed Production and Germination in *Cyperus esculentus* and *C. rotundus*. Source: *Weed Science*, 27(5), 502–505. <https://www.jstor.org/stable/4043107>
- Tiwari, V. K. ., & Mishra, B. B. . (2011). *Opportunity, challenge and scope of natural products in medicinal chemistry*. Research Signpost.
- t'Kindt, R., Morreel, K., Deforce, D., Boerjan, W., & Van Bocxlaer, J. (2009). Joint GC-MS and LC-MS platforms for comprehensive plant metabolomics: Repeatability and sample pre-treatment. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences*, 877(29), 3572–3580. <https://doi.org/10.1016/j.jchromb.2009.08.041>
- Toyomasu, T., Usui, M., Sugawara, C., Otomo, K., Hirose, Y., Miyao, A., Hirochika, H., Okada, K., Shimizu, T., Koga, J., Hasegawa, M., Chuba, M., Kawana, Y., Kuroda, M., Minami, E., Mitsuhashi, W., & Yamane, H. (2014). Reverse-genetic approach to verify physiological roles of rice phytoalexins: Characterization of a knockdown mutant of OsCPS4 phytoalexin biosynthetic gene in rice. *Physiologia Plantarum*, 150(1), 55–62. <https://doi.org/10.1111/ppl.12066>
- Traxler, C., Gaines, T. A., Küpper, A., Luemmen, P., & Dayan, F. E. (2023). The nexus between reactive oxygen species and the mechanism of action of herbicides. *Journal of Biological Chemistry*, 299(11). <https://doi.org/10.1016/j.jbc.2023.105267>
- Tripathy, B. C., & Oelmüller, R. (2012). Reactive oxygen species generation and signaling in plants. *Plant Signaling and Behavior*, 7(12), 1621–1633. <https://doi.org/10.4161/psb.22455>
- Tuor, F. A., & Froud-Williams, R. J. (2002). Influence of nitrogen on competition between purple nutsedge, maize and soybean. *International Journal of Pest Management*, 48(1), 73–79. <https://doi.org/10.1080/09670870110094378>
- Upchurch, R. P., Keaton, J. A., & Coble, H. D. (1969). Woody Plant Shoot Management and Response to Herbicidal Treatment. *Weed Science*, 17(2), 175–180. <https://doi.org/10.1017/s0043174500031258>
- Venkateshwarlu, G., Ravindra, V., & Challa, P. (2001). Mangiferin: An allelopathin from mango (*Mangifera indica L.*) leaves. *Allelopathy Journal*, 8(2), 221–224. <https://www.researchgate.net/publication/289141126>



- Verdeguer, M., Blázquez, M. A., & Boira, H. (2009). Phytotoxic effects of *Lantana camara*, *Eucalyptus camaldulensis* and *Eriocephalus africanus* essential oils in weeds of Mediterranean summer crops. *Biochemical Systematics and Ecology*, 37(4), 362–369. <https://doi.org/10.1016/j.bse.2009.06.003>
- Verdeguer, M., Sánchez-Moreiras, A. M., & Araniti, F. (2020). Phytotoxic effects and mechanism of action of essential oils and terpenoids. *Plants*, 9(11), 1–48. <https://doi.org/10.3390/plants9111571>
- Walter, A., & Schurr, U. (1999). The modular character of growth in *Nicotiana tabacum* plants under steady-state nutrition. *Journal of Experimental Botany*, 50(336), 1169–1177. <https://doi.org/https://doi.org/10.1093/jxb/50.336.1169>
- Wang, X., Singh, D., Marla, S., Morris, G., & Poland, J. (2018). Field-based high-throughput phenotyping of plant height in sorghum using different sensing technologies. *Plant Methods*, 14(1). <https://doi.org/10.1186/s13007-018-0324-5>
- Wenxiang, G., Shunshan, D., & Shiming, L. (1998). Ecological Characteristic of Terpenoids and Their Allelopathic Effects to Plants. *Journal of South China Agricultural University*, 4, 108–112. https://xuebao.scau.edu.cn/zr/hnny_zren/article/pdf/199804121
- Wheeler, J. J., Seltmann, H., & Motten, A. G. (1991). The Mode of Action of Fatty Alcohols on Leaf Tissue. *J Plant Growth Regul*, 10, 129–137. <https://doi.org/https://doi.org/10.1007/BF02279324>
- Wichert, R. A., Bozsa, R., Talbert, R. E., & Oliver, L. R. (1992). Temperature and Relative Humidity Effects on Diphenylether Herbicides. *Weed Technology*, 6(1), 19–24. <https://doi.org/10.1017/s0890037x00034230>
- Willis, R. J. (2007). *The History of Allelopathy* (1st ed.). Springer Dordrecht. <https://doi.org/10.1007/978-1-4020-4093-1>
- Wittlekens, H., Inz62, D., Van Montagu, M., & Van Camp, W. (1995). Catalases in plants. In *Molecular Breeding* (Vol. 1).
- Xu, M., Galhano, R., Wiemann, P., Bueno, E., Tiernan, M., Wu, W., Chung, I. M., Gershenson, J., Tudzynski, B., Sesma, A., & Peters, R. J. (2012). Genetic evidence for natural product-mediated plant-plant allelopathy in rice (*Oryza sativa*). *New Phytologist*, 193(3), 570–575. <https://doi.org/10.1111/j.1469-8137.2011.04005.x>
- Xu, Y., Wu, Y. G., Chen, Y., Zhang, J. F., Song, X. Q., Zhu, G. P., & Hu, X. W. (2015). Autotoxicity in *pogostemon cablin* and their allelochemicals. *Revista Brasileira de Farmacognosia*, 25(2), 117–123. <https://doi.org/10.1016/j.bjfp.2015.02.003>
- Yan, Z. Q., Wang, D. D., Ding, L., Cui, H. Y., Jin, H., Yang, X. Y., Yang, J. S., & Qin, B. (2015). Mechanism of artemisinin phytotoxicity action: Induction of reactive oxygen species and cell death in lettuce seedlings. *Plant Physiology and Biochemistry*, 88, 53–59. <https://doi.org/10.1016/j.plaphy.2015.01.010>
- You, C., Wang, H., Huang, Y., Xu, P., Wu, L., Yu, F., Zhong, X., Gao, J., Zhang, L., He, H., & Ke, J. (2021). Relationship Between the Vascular



- Bundle Structure of Panicle Branches and the Filling of Inferior Spikelets in Large-Panicle Japonica Rice. *Frontiers in Plant Science*, 12. <https://doi.org/10.3389/fpls.2021.774565>
- Zanuncio, A., Teodoro, P. E., Riberio, L. P., Correa, C. C. G., & Torres, M. O. e F. E. (2013). Allelopathy of green manures on *Cyperus rotundus*. *Revista de Ciências Agrárias*, 36(4), 441–446. <https://www.researchgate.net/publication/317475150>
- Zhang, S., Zhang, B., Dai, W., & Zhang, X. (2011). Oxidative damage and antioxidant responses in *Microcystis aeruginosa* exposed to the allelochemical berberine isolated from golden thread. *Journal of Plant Physiology*, 168(7), 639–643. <https://doi.org/10.1016/j.jplph.2010.10.005>
- Zhang, Z. H., Hu, B. Q., & Hu, G. (2013). Assessment of allelopathic potential of *Wedelia trilobata* on the germination, seedling growth and chlorophyll content of rape. *Advanced Materials Research*, 807–809, 719–722. <https://doi.org/10.4028/www.scientific.net/AMR.807-809.719>
- Zhou, B., Wang, J., Guo, Z., Tan, H., & Zhu, X. (2006). A simple colorimetric method for determination of hydrogen peroxide in plant tissues. *Plant Growth Regulation*, 49(2–3), 113–118. <https://doi.org/10.1007/S10725-006-9000-2/METRICS>
- Zhou, Z. L., Yin, W. Q., Yang, Y. M., He, C. H., Li, X. N., Zhou, C. P., & Guo, H. (2016). New Iridoid Glycosides with Antidepressant Activity Isolated from *Cyperus rotundus*. *Chemical & Pharmaceutical Bulletin*, 64(1), 73–77. <https://doi.org/10.1248/CPB.C15-00686>
- Zohaib, A., Anjum, S. A., Jabbar, A., Tabassum, T., Abbas, T., & Nazir, U. (2017). Allelopathic Effect of Leguminous Weeds on Rate, Synchronization and Time of Germination, and Biomass Partitioning in Rice. *Planta Daninha*, 35. <https://doi.org/10.1590/S0100-83582017350100032>