

REFERENCES

- Alston, J.M., Sambucci, O. (2019). Grapes in the World Economy. In: Cantu, D., Walker, M. (eds) *The Grape Genome. Compendium of Plant Genomes*. Springer, Cham. https://doi.org/10.1007/978-3-030-18601-2_1
- Arite, Tomotsugu, Kana Ishizaki, Ayaka Date, Kenichi Matsuda, Tetsuya Sasaki, & Akira Miwa. (2015). Analysis of sugars, organic acids and volatile components of grape ‘Ruby Roman’. 有手友嗣, 石崎佳奈, 伊達彩香, 松田賢一, 笹木哲也, & 三輪章志. (2015). ブドウ ‘ルビーロマン’の糖, 有機酸および揮発性成分の分析.
- Artés-Hernández, F., Aguayo, E., & Artés, F. (2004). Alternative atmosphere treatments for keeping quality of ‘Autumn seedless’ table grapes during long-term cold storage. *Postharvest biology and Technology*, 31(1), 59-67. [https://doi.org/10.1016/S0925-5214\(03\)00116-9](https://doi.org/10.1016/S0925-5214(03)00116-9)
- Asproudi, A., Petrozziello, M., Cavalletto, S., & Guidoni, S. (2016). Grape aroma precursors in cv. Nebbiolo as affected by vine microclimate. *Food chemistry*, 211, 947-956. <https://doi.org/10.1016/j.foodchem.2016.05.070>
- Baek, H. H., & Cadwallader, K. R. (1999). Contribution of free and glycosidically bound volatile compounds to the aroma of muscadine grape juice. *Journal of food Science*, 64(3), 441-444. <https://doi.org/10.1111/j.1365-2621.1999.tb15059.x>
- Bartley, I. M., Stoker, P. G., Martin, A. D., Hatfield, S. G., & Knee, M. (1985). Synthesis of aroma compounds by apples supplied with alcohols and methyl esters of fatty acids. *Journal of the Science of Food and Agriculture*, 36(7), 567-574. <https://doi.org/10.1002/jsfa.2740360708>
- Bhat, N. R., Desai, B. B., & Suleiman, M. K. (2010). Flavor in grapes: its characterization and commercial applications. *Handbook of Fruit and Vegetable Flavors*, 279-302. <https://doi.org/10.1002/9780470622834.ch17>
- Boss, P. K., & Davies, C. (2009). Molecular biology of anthocyanin accumulation in grape berries. In *Grapevine molecular physiology & biotechnology* (pp. 263-292). Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-90-481-2305-6_10
- Capone, S., Tufariello, M., & Siciliano, P. (2013). Analytical characterisation of Negroamaro red wines by “Aroma Wheels”. *Food Chemistry*, 141(3), 2906-2915. <https://doi.org/10.1016/j.foodchem.2013.05.105>
- Chang, E. H., Jeong, S. M., Hur, Y. Y., Koh, S. W., & Choi, I. M. (2015). Changes of volatile compounds in *Vitis labrusca* ‘Doonuri’ grapes during stages of

- fruit development and in wine. *Horticulture, Environment, and Biotechnology*, 56, 137-144. <https://doi.org/10.1007/s13580-015-0068-3>
- Cheng, G. T., Wang, X. Y., Li, Y. S., Qi, S. M., Wang, J., Zhao, P., ... & Liang, Y. (2021). Comparison of volatile components and aroma profiles between peel and flesh of tomatoes (*Solanum lycopersicum*). <https://doi.org/10.7506/spkx1002-6630-20200930-374>
- Choi, K. O., Hur, Y. Y., Park, S. J., Lee, D. H., Kim, S. J., & Im, D. (2022). Relationships between instrumental and sensory quality indices of shine muscat grapes with different harvesting times. *Foods*, 11(16), 2482. <https://doi.org/10.3390/foods11162482>
- Darriet, P., Thibon, C., Dubourdiou, D., Gerós, H., Chaves, M. M., & Delrot, S. (2012). Aroma and aroma precursors in grape berry. *The biochemistry of the grape berry*, 26, 111-136.
- Davies, C., & Böttcher, C. (2009). Hormonal control of grape berry ripening. In *Grapevine molecular physiology & biotechnology* (pp. 229-261). Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-90-481-2305-6_9
- de Ancos, B., Ibanez, E., Reglero, G., & Cano, M. P. (2000). Frozen storage effects on anthocyanins and volatile compounds of raspberry fruit. *Journal of agricultural and food chemistry*, 48(3), 873-879. <https://doi.org/10.1021/jf990747c>
- Deng, H., He, R., Long, M., Li, Y., Zheng, Y., Lin, L., ... & Xia, H. (2021). Comparison of the fruit volatile profiles of five muscadine grape cultivars (*Vitis rotundifolia* Michx.) using HS-SPME-GC/MS combined with multivariate statistical analysis. *Frontiers in plant science*, 12, 728891. <https://doi.org/10.3389/fpls.2021.728891>
- Doligez, A., Audiot, E., Baumes, R., & This, P. (2006). QTLs for muscat flavor and monoterpenic odorant content in grapevine (*Vitis vinifera* L.). *Molecular Breeding*, 18(2), 109-125. <https://doi.org/10.1007/s11032-006-9016-3>
- Dunlevy, J., Kalua, C., Keyzers, R., Boss, P. (2009). The Production of Flavour & Aroma Compounds in Grape Berries. In: Roubelakis-Angelakis, K.A. (eds) *Grapevine Molecular Physiology & Biotechnology*. Springer, Dordrecht. https://doi.org/10.1007/978-90-481-2305-6_11
- Dutra, M. D. C. P., de Souza, J. F., Viana, A. C., de Oliveira, D., Pereira, G. E., & dos Santos Lima, M. (2018). Rapid determination of the aromatic compounds methyl-anthranilate, 2'-aminoacetophenone and furaneol by GC-MS: Method validation and characterization of grape derivatives. *Food research international*, 107, 613-618. <https://doi.org/10.1016/j.foodres.2018.03.020>

- El Hadi, M. A. M., Zhang, F. J., Wu, F. F., Zhou, C. H., & Tao, J. (2013). Advances in fruit aroma volatile research. *Molecules*, 18(7), 8200-8229. <https://doi.org/10.3390/molecules18078200>
- Falchi, M., Bertelli, A., Lo Scalzo, R., Morassut, M., Morelli, R., Das, S., ... & Das, D. K. (2006). Comparison of cardioprotective abilities between the flesh and skin of grapes. *Journal of agricultural and food chemistry*, 54(18), 6613-6622. <https://doi.org/10.1021/jf061048k>
- Fan, W., Xu, Y., Jiang, W., & Li, J. (2010). Identification and quantification of impact aroma compounds in 4 nonfloral *Vitis vinifera* varieties grapes. *Journal of food science*, 75(1), S81-S88. <https://doi.org/10.1111/j.1750-3841.2009.01436.x>
- Fenoll, J., Manso, A., Hellín, P., Ruiz, L., & Flores, P. (2009). Changes in the aromatic composition of the *Vitis vinifera* grape Muscat Hamburg during ripening. *Food Chemistry*, 114(2), 420-428. <https://doi.org/10.1016/j.foodchem.2008.09.060>
- Fernández-Fernández, A. M., Iriando-DeHond, A., Nardin, T., Larcher, R., Dellacassa, E., Medrano-Fernandez, A., & Castillo, M. D. D. (2020). In Vitro bioaccessibility of extractable compounds from tannat grape skin possessing health promoting properties with potential to reduce the risk of diabetes. *Foods*, 9(11), 1575. <https://doi.org/10.3390/foods10030672>
- Fisher, W. B., & VanPeppen, J. F. (2000). Cyclohexanol and cyclohexanone. *Kirk-Othmer Encyclopedia of Chemical Technology*. <https://doi.org/10.1002/0471238961.0325031206091908.a01>
- Flamini, R., & Traldi, P. (2009). *Mass spectrometry in grape and wine chemistry*. John Wiley & Sons. DOI:10.1002/9780470552926
- Forney, C. F., Kalt, W., & Jordan, M. A. (2000). The composition of strawberry aroma is influenced by cultivar, maturity, and storage. *HortScience*, 35(6), 1022-1025. <https://doi.org/10.21273/HORTSCI.35.6.1022>
- Francis, I. L., & Newton, J. L. (2005). Determining wine aroma from compositional data. *Australian Journal of Grape and Wine Research*, 11(2), 114-126. <https://doi.org/10.1111/j.1755-0238.2005.tb00283.x>
- Franco, M., Peinado, R. A., Medina, M., & Moreno, J. (2004). Off-vine grape drying effect on volatile compounds and aromatic series in must from Pedro Ximénez grape variety. *Journal of Agricultural and Food Chemistry*, 52(12), 3905-3910. DOI: 10.1021/jf0354949
- García-Carpintero, E. G., Sánchez-Palomo, E., Gallego, M. G., & González-Viñas, M. A. (2011). Volatile and sensory characterization of red wines from cv. Moravia Agria minority grape variety cultivated in La Mancha region over five consecutive vintages. *Food Research International*, 44(5), 1549-1560. <https://doi.org/10.1016/j.foodres.2011.04.022>

- García, E., Chacón, J. L., Martínez, J., & Izquierdo, P. M. (2003). Changes in volatile compounds during ripening in grapes of Airén, Macabeo and Chardonnay white varieties grown in La Mancha region (Spain). *Food science and technology international*, 9(1), 33-41. <https://doi.org/10.1177/1082013203009001006>
- García-Lomillo, J., & González-SanJosé, M. L. (2017). Applications of wine pomace in the food industry: Approaches and functions. *Comprehensive reviews in food science and food safety*, 16(1), 3-22. <https://doi.org/10.1111/1541-4337.12238>
- Genovese, A., Lamorte, S. A., Gambuti, A., & Moio, L. (2013). Aroma of Aglianico and Uva di Troia grapes by aromatic series. *Food research international*, 53(1), 15-23. <https://doi.org/10.1016/j.foodres.2013.03.051>
- Genovese, A., Basile, B., Lamorte, S. A., Lisanti, M. T., Corrado, G., Lecce, L., ... & Gambuti, A. (2021). Influence of Berry Ripening Stages over Phenolics and Volatile Compounds in Aged Aglianico Wine. *Horticulturae*, 7(7), 184. <https://doi.org/10.1016/j.foodres.2013.03.051>
- Gomez, E., Martinez, A., & Laencina, J. (2015). Localization of free and bound aromatic compounds among skin, juice and pulp fractions some grape varieties. *VITIS-Journal of Grapevine Research*, 33(1), 1.
- González-Centeno, M. R., Rosselló, C., Simal, S., Garau, M. C., López, F., & Femenia, A. (2010). Physico-chemical properties of cell wall materials obtained from ten grape varieties and their byproducts: Grape pomaces and stems. *LWT - Food Science and Technology*, 43(10), 1580–1586. <https://doi.org/10.1016/j.lwt.2010.06.024>
- Gonçalves, G. A., Resende, N. S., Carvalho, E. E., de Resende, J. V., & Vilas Boas, E. V. D. B. (2018). Physicochemical and volatile profile alterations in pasteurized and frozen strawberry pulp during storage. *Journal of Food Processing and Preservation*, 42(1), e13317. <https://doi.org/10.1111/jfpp.13317>
- Gougoulis, N., & Mashev, N. (2008). Evaluation of polyphenols antioxidant activity of grape seeds (V. vinifera). *Oxidation Communications*, 31(1), 88.
- Gunata, Y. Z., Bayonove, C. L., Baumes, R. L., & Cordonnier, R. E. (1985). The aroma of grapes. Localisation and evolution of free and bound fractions of some grape aroma components cv Muscat during first development and maturation. *Journal of the Science of Food and Agriculture*, 36(9), 857-862. <https://doi.org/10.1002/jsfa.2740360915>
- Gu, X., Zhang, X., Wang, K., Lv, X., Li, R., & Ma, W. (2022). GC–MS Untargeted analysis of volatile compounds in four red grape Varieties (Vitis vinifera

- L. cv) at Different Maturity Stages near Harvest. *Foods*, 11(18), 2804. <https://doi.org/10.3390/foods11182804>
- Hirano, K., Shibahara, R., Asaoka, K., & Okamoto, G. (1998). Seasonal changes in monoterpenes of Muscat of Alexandria grape juice during berry development. *Journal of the Japanese Society for Horticultural Science*, 67(6), 907-911. <https://doi.org/10.2503/jjshs.67.907>
- Jayaprakasha, G. K., Kumar, C. K., Chaudhary, P., Murthy, K. C., & Patil, B. S. (2013). Identification of volatiles from Kumquats and their biological activities. In *Tropical and subtropical fruits: Flavors, color, and health benefits* (pp. 63-92). American Chemical Society. DOI: 10.1021/bk-2013-1129.ch004
- Ji ShuJuan, J. S., Bu QingZhuang, B. Q., Li JiangKuo, L. J., & Zhang Ping, Z. P. (2012). Effects of 1-MCP treatment on aroma components of Nanguo pear during shelf life after cold storage.
- Ju, Y. L., Liu, M., Zhao, H., Meng, J. F., & Fang, Y. L. (2016). Effect of exogenous abscisic acid and methyl jasmonate on anthocyanin composition, fatty acids, and volatile compounds of Cabernet Sauvignon (*Vitis vinifera* L.) grape berries. *Molecules*, 21(10), 1354. <https://doi.org/10.3390/molecules21101354>
- Kalua, C. M., & Boss, P. K. (2009). Evolution of volatile compounds during the development of Cabernet Sauvignon grapes (*Vitis vinifera* L.). *Journal of agricultural and food chemistry*, 57(9), 3818-3830. <https://doi.org/10.1021/jf803471n>
- Kaminaga, Y., Schnepf, J., Peel, G., Kish, C. M., Ben-Nissan, G., Weiss, D., ... & Dudareva, N. (2006). Plant phenylacetaldehyde synthase is a bifunctional homotetrameric enzyme that catalyzes phenylalanine decarboxylation and oxidation. *Journal of Biological Chemistry*, 281(33), 23357-23366. <https://doi.org/10.1074/jbc.M602708200>
- Kandylis, P. (2021). Grapes and their derivatives in functional foods. *Foods*, 10(3), 672. <https://doi.org/10.3390/foods10030672>
- Kang, S., Yan, H., Zhu, Y., Liu, X., Lv, H. P., Zhang, Y., ... & Lin, Z. (2019). Identification and quantification of key odorants in the world's four most famous black teas. *Food Research International*, 121, 73-83. <https://doi.org/10.1016/j.foodres.2019.03.009>
- Khalil, M. N., Fekry, M. I., & Farag, M. A. (2017). Metabolome based volatiles profiling in 13 date palm fruit varieties from Egypt via SPME GC-MS and chemometrics. *Food chemistry*, 217, 171-181. <https://doi.org/10.1016/j.foodchem.2016.08.089>
- Koyama, R., M. Demura, S. Noma, N. Hayashi, T. Haraguchi, E. Miyamoto, ... & Ueno, Daisuke. (2021). Identification of odorous substances in the violet

- spider *Trentepohlia aurea* (Linnaeus) Martius by sniffing gas chromatography. *Journal of Odour, Scent and Environment*, 52(4), 226-232. 小山玲音, 出村幹英, 野間誠司, 林信行, 原口智和, 宮本英揮, ... & 上野大介. (2021). スミレモ *Trentepohlia aurea* (Linnaeus) Martius のにおい嗅ぎガスクロマトグラフィーによるにおい物質の同定. におい・かおり環境学会誌, 52(4), 226-232. <https://doi.org/10.2171/jao.52.226>
- Khan, N., Fahad, S., Naushad, M., & Faisal, S. (2020). Grape production critical review in the world. Available at SSRN 3595842. <https://dx.doi.org/10.2139/ssrn.3595842>
- Kumazawa, K., & Masuda, H. (2002). Identification of potent odorants in different green tea varieties using flavor dilution technique. *Journal of Agricultural and Food Chemistry*, 50(20), 5660-5663. <https://doi.org/10.1021/jf020498j>
- Langos, D., Granvogl, M., & Schieberle, P. (2013). Characterization of the key aroma compounds in two Bavarian wheat beers by means of the sensomics approach. *Journal of agricultural and food chemistry*, 61(47), 11303-11311. <https://doi.org/10.1021/jf403912j>
- Li, X. L., Luan, K. A. N. G., HU, J. J., LI, X. F., & Xiang, S. H. E. N. (2008). Aroma volatile compound analysis of SPME headspace and extract samples from crabapple (*Malus* sp.) fruit using GC-MS. *Agricultural Sciences in China*, 7(12), 1451-1457. [https://doi.org/10.1016/S1671-2927\(08\)60402-2](https://doi.org/10.1016/S1671-2927(08)60402-2)
- Li, C., Xin, M., Li, L., He, X., Yi, P., Tang, Y., ... & Li, Z. (2021a). Characterization of the aromatic profile of purple passion fruit (*Passiflora edulis* Sims) during ripening by HS-SPME-GC/MS and RNA sequencing. *Food chemistry*, 355, 129685. <https://doi.org/10.1016/j.foodchem.2021.129685>
- Li, S., Chen, K., & Grierson, D. (2021b). Molecular and hormonal mechanisms regulating fleshy fruit ripening. *Cells*, 10(5), 1136. <https://doi.org/10.3390/cells10051136>
- Li, D., Zeng, Y., Ye, Z. K., Li, H. K., Li, Y. Z., Dong, B., ... & Zhong, H. N. (2023). Analysis of volatile organic compounds and potential odour compounds in food contact paperboard using headspace two-dimensional GC-QTOF-MS. *Food Additives & Contaminants: Part A*, 40(11), 1482-1493. <https://doi.org/10.1080/19440049.2023.2259029>
- Li, Y., He, L., Song, Y., Zhang, P., Chen, D., Guan, L., & Liu, S. (2023). Comprehensive study of volatile compounds and transcriptome data providing genes for grape aroma. *BMC Plant Biology*, 23(1), 171. <https://doi.org/10.1186/s12870-023-04191-1>

- Lin, J., Massonnet, M., & Cantu, D. (2019). The genetic basis of grape and wine aroma. *Horticulture Research*, 6. <https://doi.org/10.1038%2Fs41438-019-0163-1>
- Ma, C., Li, J., Chen, W., Wang, W., Qi, D., Pang, S., & Miao, A. (2018). Study of the aroma formation and transformation during the manufacturing process of oolong tea by solid-phase micro-extraction and gas chromatography–mass spectrometry combined with chemometrics. *Food Research International*, 108, 413-422. <https://doi.org/10.1016/j.foodres.2018.03.052>
- Mahattanatawee, K., Perez-Cacho, P. R., Davenport, T., & Rouseff, R. (2007). Comparison of three lychee cultivar odor profiles using gas chromatography-olfactometry and gas chromatography-sulfur detection. *Journal of agricultural and food chemistry*, 55(5), 1939–1944. <https://doi.org/10.1021/jf062925p>
- Matsumoto, H., & Ikoma, Y. (2016). Effect of postharvest temperature on the muscat flavor and aroma volatile content in the berries of “Shine Muscat”(Vitis labruscana Baily× V. vinifera L.). *Postharvest Biology and Technology*, 112, 256-265. <https://doi.org/10.1016/j.postharvbio.2015.09.004>
- Maoz, I., Rikanati, R. D., Schlesinger, D., Bar, E., Gonda, I., Levin, E., ... & Lewinsohn, E. (2018). Concealed ester formation and amino acid metabolism to volatile compounds in table grape (Vitis vinifera L.) berries. *Plant science*, 274, 223-230. <https://doi.org/10.1016/j.plantsci.2018.05.020>
- Maoz, I., Kaplunov, T., Raban, E., Dynkin, I., Degani, O., Lewinsohn, E., & Lichter, A. (2020). Insights into the chemosensory basis of flavor in table grapes. *Journal of the Science of Food and Agriculture*, 100(4), 1405-1417. <https://doi.org/10.1002/jsfa.10102>
- McCARTHY, A. I., PALMER, J. K., SHAW, C. P., & ANDERSON, E. E. (1963). Correlation of Gas Chromatographic Data with Flavor Profiles of Fresh Banana Fruit a. *Journal of Food Science*, 28(4), 379-384. <https://doi.org/10.1111/j.1365-2621.1963.tb00214.x>
- Mihaylova, D., Popova, A., & Dincheva, I. (2022). Pattern recognition of varieties of peach fruit and pulp from their volatile components and metabolic profile using HS-SPME-GC/MS combined with multivariable statistical analysis. *Plants*, 11(23), 3219. <https://doi.org/10.3390/plants11233219>
- Ministry of Agriculture, Forestry and Fisheries Public Relations Office. (2019). Special Feature 1 Grape. *Agriculture Forestry Fisheries*. 50(5), p.8.
- Modise, D. M. (2008). Does freezing and thawing affect the volatile profile of strawberry fruit (Fragaria× ananassa Duch.). *Postharvest Biology and*

- Technology*, 50(1), 25-30.
<https://doi.org/10.1016/j.postharvbio.2008.03.009>
- Moriyama, K., Kono, A., Matsuzaki, R., Azuma, A., Onoue, N., Sekozawa, Y., ... & Sugaya, S. (2024). Diversity of flavour characteristics of table grapes and their contributing volatile compounds analysed by the solvent-assisted flavour evaporation method. *Horticulture Research*, 11(4), uhae048. <https://doi.org/10.1093/hr/uhae048>
- Noguerol-Pato, R., González-Barreiro, C., Cancho-Grande, B., Santiago, J. L., Martínez, M. C., & Simal-Gándara, J. (2012). Aroma potential of Brancellao grapes from different cluster positions. *Food chemistry*, 132(1), 112-124. <https://doi.org/10.1016/j.foodchem.2011.10.042>
- Nouri, B., Mohtasebi, S. S., & Rafiee, S. (2020). Quality detection of pomegranate fruit infected with fungal disease. *International Journal of Food Properties*, 23(1), 9-21. <https://doi.org/10.1080/10942912.2019.1705851>
- Okamoto, G., Liao, K., Fushimi, T., & Hirano, K. (2001). Aromatic substances evolved from the whole berry, skin, and flesh of Muscat of Alexandria grapes. *岡山大学農学部学術報告*, 90(1), 21-25.
- OuYang, Y. N., Gao, J. S., Li, R. L., Zhu, M. R., Hu, X. H., Min, Z., ... & Fang, Y. L. (2015). Changes in volatile profiles and activity of hydroperoxide lyase and alcohol dehydrogenase during the development of Cabernet Sauvignon grapes (*Vitis vinifera* L.). *South African Journal of Enology and Viticulture*, 36(3), 328-342. <https://doi.org/10.21548/36-3-966>
- Panighel, A., Vedova, A. D., De Rosso, M., Gardiman, M., & Flamini, R. (2014). A solid-phase microextraction gas chromatography/ion trap tandem mass spectrometry method for simultaneous determination of 'foxy smelling compounds' and 3-alkyl-2-methoxypyrazines in grape juice. *Rapid Communications in Mass Spectrometry*, 24(14), 2023-2029. <https://doi.org/10.1002/rcm.4611>
- Park, S. K., Morrison, J. C., Adams, D. O., & Noble, A. C. (1991). Distribution of free and glycosidically bound monoterpenes in the skin and mesocarp of Muscat of Alexandria grapes during development. *Journal of agricultural and food chemistry*, 39(3), 514-518. <https://doi.org/10.1021/jf00003a017>
- Pashaei, M., & Hassanpour, H. (2024). Phenolic, amino acids, and fatty acids profiles and the nutritional properties in the fresh and dried fruits of black rosehip (*Rosa pimpinellifolia* L.). *Scientific Reports*, 14(1), 19665. <https://doi.org/10.1038/s41598-024-70574-5>
- Peinado, R. A., Moreno, J., Bueno, J. E., Moreno, J. A., & Mauricio, J. C. (2004). Comparative study of aromatic compounds in two young white wines subjected to pre-fermentative cryomaceration. *Food Chemistry*, 84(4), 585-590. [https://doi.org/10.1016/S0308-8146\(03\)00282-6](https://doi.org/10.1016/S0308-8146(03)00282-6)

- Pico, J., Nozadi, K., Gerbrandt, E. M., Dossett, M., & Castellarin, S. D. (2023). Determination of bound volatiles in blueberries, raspberries, and grapes with an optimized protocol and a validated SPME-GC/MS method. *Food Chemistry*, 403, 134304. <https://doi.org/10.1016/j.foodchem.2022.134304>
- Pollner, G., & Schieberle, P. (2016). Characterization of the key odorants in commercial cold-pressed oils from unpeeled and peeled rapeseeds by the sensomics approach. *Journal of agricultural and food chemistry*, 64(3), 627-636. <https://doi.org/10.1021/acs.jafc.5b05321>
- Qi, Y., Wang, M., Wan, N., Yin, D., Wei, M., Sun, X., ... & Ma, T. (2024). Sensory characteristics of “Shine Muscat” grapes based on consumer reviews and human and intelligent sensory evaluation. *LWT*, 115810. <https://doi.org/10.1016/j.lwt.2024.115810>
- Qin, D., Wang, Q., Jiang, X., Ni, E., Fang, K., Li, H., ... & Wu, H. (2023). Identification of key volatile and odor-active compounds in 10 main fragrance types of Fenghuang Dancong tea using HS-SPME/GC-MS combined with multivariate analysis. *Food Research International*, 173, 113356. <https://doi.org/10.1016/j.foodres.2023.113356>
- Reineccius, G. (2005). *Flavor Chemistry and Technology* (2nd ed.). CRC Press. <https://doi.org/10.1201/9780203485347>
- Reisch, B.I., Owens, C.L., Cousins, P.S. (2012). Grape. In: Badenes, M., Byrne, D. (eds) *Fruit Breeding. Handbook of Plant Breeding*, vol 8. Springer, Boston, MA. https://doi.org/10.1007/978-1-4419-0763-9_7
- ROBINSON, S. P., & DAVIES, C. (2000). Molecular biology of grape berry ripening. *Australian journal of grape and wine research*, 6(2), 175-188. <https://doi.org/10.1111/j.1755-0238.2000.tb00177.x>
- Rolle, L., Torchio, F., Zeppa, G., & Gerbi, V. (2009). Relationship between skin break force and anthocyanin extractability at different ripening stages. *American Journal of Enology and Viticulture*, 60(1), 93-97. DOI : 10.5344/ajev.2009.60.1.93
- Sasaki, K., Takase, H., Matsuyama, S., Kobayashi, H., Matsuo, H., Ikoma, G., & Takata, R. (2016). Effect of light exposure on linalool biosynthesis and accumulation in grape berries. *Bioscience, Biotechnology, and Biochemistry*, 80(12), 2376-2382. <https://doi.org/10.1080/09168451.2016.1217148>
- Sasaki, T., Koshi, E., Take, H., Michihata, T., Maruya, M., & Enomoto, T. (2017). Characterisation of odorants in roasted stem tea using gas chromatography–mass spectrometry and gas chromatography–olfactometry analysis. *Food chemistry*, 220, 177-183. <https://doi.org/10.1016/j.foodchem.2016.09.208>

- Sasaki, T., Ando, S., Miyazawa, T., Yamauchi, D., Take, H., Yamazaki, Y., & Enomoto, T. (2020). Characterisation of ‘Ruby Roman’ table grapes (*Vitis Labruscana* Bailey) by sensory evaluation and analysis of aroma and taste compounds. *Food Science and Technology Research*, 26(3), 423-434. <https://doi.org/10.3136/fstr.26.423>
- Sato, M., Suzuki, Y., Okuda, T., & Yokotsuka, K. (1997). Contents of resveratrol, piceid, and their isomers in commercially available wines made from grapes cultivated in Japan. *Bioscience, biotechnology, and biochemistry*, 61(11), 1800-1805. <https://doi.org/10.1271/bbb.61.1800>
- Schwab, W., & Schreier, P. (2002). Enzymic formation of flavor volatiles from lipids. In *Lipid biotechnology* (pp. 342-372). CRC Press.
- Schwab, W., Davidovich-Rikanati, R., & Lewinsohn, E. (2008). Biosynthesis of plant-derived flavor compounds. *The plant journal*, 54(4), 712-732. <https://doi.org/10.1111/j.1365-313X.2008.03446.x>
- SHIRAISHI, S. I., & WATANABE, Y. (1988). Anthocyanin Pigments in the Blue-Black Tetraploid Grape Cultivars ‘Black Olympia’, ‘Pione’ and ‘Izunishiki’ (*Vitis vinifera* L. × *V. labrusca* L.). *Journal of the Japanese Society for Horticultural Science*, 57(1), 17-21. <https://doi.org/10.2503/jjshs.57.17>
- Shiraishi, M. (1993). Three descriptors for sugars to evaluate grape germplasm. *Euphytica*, 71, 99-106. <https://doi.org/10.1007/BF00023472>
- Shiraishi, M. (2000). Comparison in changes in sugars, organic acids and amino acids during berry ripening of sucrose- and hexose-accumulating grape cultivars. *Journal of the Japanese Society for Horticultural Science*, 69(2), 141-148. <https://doi.org/10.2503/jjshs.69.141>
- Shiraishi, M., Shinomiya, R., & Chijiwa, H. (2012). Preliminary genetic analysis of sucrose accumulation in berries of table grapes. *Scientia horticulturae*, 137, 107-113. <https://doi.org/10.1016/j.scienta.2012.01.024>
- Shiraishi, M., Shinomiya, R., & Chijiwa, H. (2018). Varietal differences in polyphenol contents, antioxidant activities and their correlations in table grape cultivars bred in Japan. *Scientia horticulturae*, 227, 272-277. <https://doi.org/10.1016/j.scienta.2017.09.032>
- Slegers, A., Angers, P., Ouellet, É., Truchon, T., & Pedneault, K. (2015). Volatile compounds from grape skin, juice and wine from five interspecific hybrid grape cultivars grown in Québec (Canada) for wine production. *Molecules*, 20(6), 10980-11016. <https://doi.org/10.3390/molecules200610980>
- Stevens, K. L., Bomben, J., Lee, A., & McFadden, W. H. (1966). Volatiles from grapes. Muscat of Alexandria. *Journal of Agricultural and Food Chemistry*, 14(3), 249-252. <https://doi.org/10.1021/jf60145a015>

- Strauss, C. R., Wilson, B., Gooley, P. R., & Williams, P. J. (1986). Role of monoterpenes in grape and wine flavor. DOI: 10.1021/bk-1986-0317.ch018
- Suehiro, Y., Mochida, K., Tsuma, M., Yasuda, Y., Itamura, H., & Esumi, T. (2019). Effects of abscisic acid/ethephon treatments on berry development and maturation in the yellow-green skinned “Shine Muscat” grape. *The Horticulture Journal*, 88(2), 189-201. <https://doi.org/10.2503/hortj.UTD-045>
- Sugiura, T., Shiraishi, M., Konno, S., & Sato, A. (2018). Prediction of skin coloration of grape berries from air temperature. *The Horticulture Journal*, 87(1), 18-25. <https://doi.org/10.2503/hortj.OKD-061>
- Sun, Q., Gates, M. J., Lavin, E. H., Acree, T. E., & Sacks, G. L. (2011). Comparison of odor-active compounds in grapes and wines from *Vitis vinifera* and non-foxy American grape species. *Journal of agricultural and food chemistry*, 59(19), 10657-10664. <https://doi.org/10.1021/jf2026204>
- Swiegers, J. H., Bartowsky, E. J., Henschke, P. A., & Pretorius, I. (2005). Yeast and bacterial modulation of wine aroma and flavour. *Australian Journal of grape and wine research*, 11(2), 139-173. <https://doi.org/10.1111/j.1755-0238.2005.tb00285.x>
- Tanaka, F., Miyazawa, T., Okazaki, K., Tatsuki, M., & Ito, T. (2015). Sensory and metabolic profiles of “Fuji” apples (*Malus domestica* Borkh.) grown without synthetic agrochemicals: the role of ethylene production. *Bioscience, biotechnology, and biochemistry*, 79(12), 2034-2043. <https://doi.org/10.1080/09168451.2015.1062713>
- Verries, C., Pradal, M., Chatelet, P., Torregrosa, L., & Tesniere, C. (2004). Isolation and analysis of the promoter of VvAdh2, a grapevine (*Vitis vinifera* L.) ripening-related gene. *Plant Science*, 167(5), 1067-1074. <https://doi.org/10.1016/j.plantsci.2004.06.003>
- Wang, J., & Luca, V. D. (2005). The biosynthesis and regulation of biosynthesis of Concord grape fruit esters, including ‘foxy’ methylanthranilate. *The Plant Journal*, 44(4), 606-619. <https://doi.org/10.1111/j.1365-313X.2005.02552.x>
- Wang, Q., Shao, Y., Cao, T. & Shen, J. (2018). Identification of Key Odor Compounds from Three Kinds of Wood Species. In Y. Wang (Ed.), *ESSE 2017: Proceedings of the International Conference on Environmental Science and Sustainable Energy* Ed. by ZhaoYang Dong (pp. 101-116). Berlin, Boston: De Gruyter. <https://doi.org/10.1515/9783110540048-011>
- Wang, J., Han, Y., Chen, C., Sam, F. E., Guan, R., Wang, K., ... & Jiang, Y. (2023). Influence of Benzothiadiazole on the Accumulation and Metabolism of C6

- Compounds in Cabernet Gernischt Grapes (*Vitis vinifera* L.). *Foods*, 12(19), 3710. <https://doi.org/10.3390/foods12193710>
- Williams, P. J., Strauss, C. R., & Wilson, B. (1980). Hydroxylated linalool derivatives as precursors of volatile monoterpenes of Muscat grapes. *Journal of Agricultural and Food Chemistry*, 28(4), 766-771. <https://doi.org/10.1021/jf60230a037>
- Willner, B., Granvogl, M., & Schieberle, P. (2013). Characterization of the key aroma compounds in Bartlett pear brandies by means of the sensomics concept. *Journal of agricultural and food chemistry*, 61(40), 9583-9593.
- Wilson, B., Strauss, C. R., & Williams, P. J. (1984). Changes in free and glycosidically bound monoterpenes in developing Muscat grapes. *Journal of Agricultural and Food Chemistry*, 32(4), 919-924. <https://doi.org/10.1021/jf00124a054>
- Winkler, A. J. (1974). *General viticulture*. Univ of California Press.
- Wu, Y., Duan, S., Zhao, L., Gao, Z., Luo, M., Song, S., ... & Wang, S. (2016). Aroma characterization based on aromatic series analysis in table grapes. *Scientific reports*, 6(1), 31116. <https://doi.org/10.1038/srep31116>
- Wu, Y., Zhang, W., Duan, S., Song, S., Xu, W., Zhang, C., ... & Wang, S. (2018). In-depth aroma and sensory profiling of unfamiliar table-grape cultivars. *Molecules*, 23(7), 1703. <https://doi.org/10.3390/molecules23071703>
- Wu, Y., Zhang, W., Yu, W., Zhao, L., Song, S., Xu, W., ... & Wang, S. (2019). Study on the volatile composition of table grapes of three aroma types. *Lwt*, 115, 108450. <https://doi.org/10.1016/j.lwt.2019.108450>
- Wu, Y., Zhang, W., Song, S., Xu, W., Zhang, C., Ma, C., ... & Wang, S. (2020). Evolution of volatile compounds during the development of Muscat grape “Shine Muscat”(Vitis labrusca× V. vinifera). *Food chemistry*, 309, 125778. <https://doi.org/10.1016/j.foodchem.2019.125778>
- Xue, H., Sekozawa, Y., & Sugaya, S. (2023). Investigating the aromatic compound changes in table grape varieties during growth and development, using HS-SPME-GC/MS. *Horticulturae*, 9(1), 85. <https://doi.org/10.3390/horticulturae9010085>
- Yaginuma, Y. (2024). Varietal differences in aroma compounds in 12 grape varieties. Faculty of Agriculture, Yamagata University, Japan.
- Yamada M., Yamane H., Sato A., Hirakawa N., Iwanami H., Yoshinaga K., Ozawa T., Mitani N., Shiraishi M., Yoshioka M. et al. (2008). New grape cultivar “Shine Muscat”. *Bull. Natl. Inst. Fruit Tree Sci.* 7, 21-38.
- Yamada, M., & Sato, A. (2016). Advances in table grape breeding in Japan. *Breeding science*, 66(1), 34-45. <https://doi.org/10.1270/jsbbs.66.34>

- Yamada, M., Yamane, H., & Sato, A. (2017). Breeding and dissemination of a new grape variety 'Shine Muscat'. *Journal of Horticultural Research*, 16(3), 229-237. 山田昌彦, 山根弘康, & 佐藤明彦. (2017). ブドウ新品種 'シャインマスカット' の育成と普及. *園芸学研究*, 16(3), 229-237. <https://doi.org/10.2503/hrj.16.229>
- Yamane, H., Kurihara, A., Yamada, M., Nagata, K., Yoshinaga, K., Matsumoto, R., Kishi, M., Ozawa, S., Kado, T., Hirabayashi, T., Kadotani, M., Sato, A., and Hirakawa, N. (1992). A new grape variety, 'Honey Black'. *Fruit Tree Experimental Report*. 23: 25–36. 山根弘康・栗原昭夫・山田昌彦・永田賢嗣・吉永勝一・松本亮司・小沢俊治・角利昭・平林利郎・角谷真奈美. 1991. ブドウ新品種 'ノースレッド'. *果樹試報*. 20: 41–48.
- Yang, C., Wang, Y., Liang, Z., Fan, P., Wu, B., Yang, L., ... & Li, S. (2009). Volatiles of grape berries evaluated at the germplasm level by headspace-SPME with GC-MS. *Food Chemistry*, 114(3), 1106-1114. <https://doi.org/10.1016/j.foodchem.2008.10.061>
- Yang, C., Wang, Y., Wu, B., Fang, J., & Li, S. (2011). Volatile compounds evolution of three table grapes with different flavour during and after maturation. *Food Chemistry*, 128(4), 823-830. <https://doi.org/10.1016/j.foodchem.2010.11.029>
- Yang, J., & Xiao, Y. Y. (2013). Grape phytochemicals and associated health benefits. *Critical reviews in food science and nutrition*, 53(11), 1202-1225. <https://doi.org/10.1080/10408398.2012.692408>
- Yang, Y., Cuenca, J., Wang, N., Liang, Z., Sun, H., Gutierrez, B., ... & Zhong, G. Y. (2020). A key 'foxy' aroma gene is regulated by homology-induced promoter indels in the iconic juice grape 'Concord'. *Horticulture research*, 7. <https://doi.org/10.1038/s41438-020-0304-6>
- Yu, H., Xie, T., Xie, J., Ai, L., & Tian, H. (2019). Characterization of key aroma compounds in Chinese rice wine using gas chromatography-mass spectrometry and gas chromatography-olfactometry. *Food Chemistry*, 293, 8-14. <https://doi.org/10.1016/j.foodchem.2019.03.071>
- Zhang, P., Fuentes, S., Siebert, T., Krstic, M., Herderich, M., Barlow, E. W. R., & Howell, K. (2016). Terpene evolution during the development of *Vitis vinifera* L. cv. Shiraz grapes. *Food Chemistry*, 204, 463-474. <https://doi.org/10.1016/j.foodchem.2016.02.125>
- Zhou, D. D., Li, J., Xiong, R. G., Saimaiti, A., Huang, S. Y., Wu, S. X., ... & Li, H. B. (2022). Bioactive compounds, health benefits and food applications of grape. *Foods*, 11(18), 2755. <https://doi.org/10.3390/foods11182755>
- Ziółkowska, A., Wąsowicz, E., & Jeleń, H. H. (2016). Differentiation of wines according to grape variety and geographical origin based on volatiles

profiling using SPME-MS and SPME-GC/MS methods. *Food Chemistry*, 213, 714-720.

<https://doi.org/10.1016/j.foodchem.2016.06.120>