

DAFTAR PUSTAKA

- Abdelmohsen Sanad E (2023) Effect of Intra-Dialytic Physical Exercise on Depression in Hemodialysis Patients. Updates on Renal Replacement Therapy. IntechOpen. Available at: <http://dx.doi.org/10.5772/intechopen.113360>.
- Agus, I., Wwpsr, B., Ratep, N., & Westa, W. (2014). Gambaran faktor-faktor yang memengaruhi tingkat depresi pada usia lanjut usia di wilayah kerja di puskesmas Kubu II Januari-Februari 2014. E-Jurnal Medika Udayana, [S.l.], apr. 2015. ISSN 2303-1395. Available at: <https://jurnal.harianregional.com/eum/id-12599>.
- Alicja Kubanek, Renke, M., Godlewska, B.R., Paul, P., Mateusz Przybylak, Kowalska, A.S., Piotr Wąż, Michał Błaszczyk, Aleksandra Macul-Sanewska, Rutkowski, P., Kamila Czarnacka, Bednarski, K. and Grabowski, J. (2024). Screening for depression in chronic haemodialysis patients as a part of care in dialysis setting: a cross-sectional study. *Frontiers in Psychiatry*, [online] 15. doi:<https://doi.org/10.3389/fpsy.2024.1410252>.
- Almutary H (2023) Psychosocial Aspects in Hemodialysis. Updates on Hemodialysis. IntechOpen. Available at: <http://dx.doi.org/10.5772/intechopen.109592>.
- Alradaydeh, M. F., & Khalil, A. A. (2019). The Effectiveness of Physical Exercise on Psychological Status, and Sleep Quality among Jordanian Patients Undergoing Hemodialysis: Literature Review. *Open Jzrnal of Nursing*, 09(12), 1267–1280. <https://doi.org/10.4236/ojn.2019.912092>
- Andrade, F. P., Rezende, P. de S., Ferreira, T. de S., Borba, G. C., Müller, A. M., & Rovedder, P. M. E. (2019). Effects of intradialytic exercise on cardiopulmonary capacity in chronic kidney disease: systematic review and meta-analysis of randomized clinical trials. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-54953-x>
- Ariyanti, I., Maria, R., & Masfuri, M. (2021). Penerapan Latihan Intradialitik terhadap Adekuasi Hemodialisis: Literature Review. *Jurnal Penelitian Kesehatan “SUARA FORIKES”* (Journal of Health Research “Forikes Voice”), 12(3), 237. <https://doi.org/10.33846/sf12303>
- Ayman Abdulfattah Elhadad, Zein, A. and Salma (2020). Psychiatric comorbidity and quality of life in patients undergoing hemodialysis. *Middle East Current Psychiatry*, [online] 27(1). doi:<https://doi.org/10.1186/s43045-020-0018-3>.

- Beck, A. T., & Alford, B. A. (2009). *Depression: Causes and treatment* (2nd ed.). University of Pennsylvania Press. Available at: <https://psycnet.apa.org/record/2009-05809-000>
- Bennett, N., Borg, W.R. and Gall, M.D. (1984) "Educational Research: An Introduction", *British Journal of Educational Studies*, 32(3), p. 274. Available at: <https://doi.org/10.2307/3121583>.
- Bertram, J. F., Douglas-Denton, R. N., Diouf, B., Hughson, M. D., & Hoy, W. E. (2011). Human nephron number: implications for health and disease. *Pediatric Nephrology*, 26(9), 1529–1533. <https://doi.org/10.1007/s00467-011-1843-8>
- Center for disease control and prevention. (2023). Chronic Kidney Disease in the United States, 2023. <https://www.cdc.gov/kidneydisease/publications-resources/CKD-national-facts.html>
- Chand (2023) *Depression* Available at: <https://pubmed.ncbi.nlm.nih.gov/33760492/>.
- Cleveland (2022) Clinical Depression (Major Depressive Disorder): Symptoms. [online] Cleveland Clinic. Available at: <https://my.clevelandclinic.org/health/diseases/24481-clinical-depression-major-depressive-disorder>
- Cukor, D., Peterson, R. A., Cohen, S. D., & Kimmel, P. L. (2006). Depression in end-stage renal disease hemodialysis patients. In *Nature Clinical Practice Nephrology* (Vol. 2, Issue 12, pp. 678–687). <https://doi.org/10.1038/ncpneph0359>
- Fan, L., Sarnak, M. J., Tighiouart, H., Drew, D. A., Kantor, A. L., Lou, K. v., Shaffi, K., Scott, T. M., & Weiner, D. E. (2014). Depression and all-cause mortality in hemodialysis patients. *American Journal of Nephrology*, 40(1), 12–18. <https://doi.org/10.1159/000363539>
- Farinde, A. (2013). "The Beck Depression Inventory", *The Pharma Innovation Journal*, 2(1), pp. 56–62. Available at: <https://www.thepharmajournal.com/archives/?year=2013&vol=2&issue=1&ArticleId=118>.
- Feng, X., Hou, N., Chen, Z., Liu, J., Li, X., Sun, X., & Liu, Y. (2023). Secular trends of epidemiologic patterns of chronic kidney disease over three decades: an

- updated analysis of the Global Burden of Disease Study 2019. *BMJ Open*, 13(3).
<https://doi.org/10.1136/bmjopen-2022-064540>
- Hall, J. E. (2017). *Guyton and Hall Textbook of Medical Physiology*.
- Hashmi, M. F., Benjamin, O., & Lappin, S. L. (2023). End-Stage Renal Disease. Available at: <https://pubmed.ncbi.nlm.nih.gov/29763036/>
- Hayden, C.M.T., Begue, G., Gamboa, J.L., Baar, K. and Roshanravan, B. (2024). Review of Exercise Interventions to Improve Clinical Outcomes in Nondialysis CKD. *Kidney International Reports*, [online] 9(11), pp.3097–3115. doi:<https://doi.org/10.1016/j.ekir.2024.07.032>.
- Henndy Ginting, Gérard Näring, William, Wilis Srisayekti and Becker, E.S. (2013). Validating the Beck Depression Inventory-II in Indonesia's general population and coronary heart disease patients. *International Journal of Clinical and Health Psychology*, [online] 13(3), pp.235–242. doi:[https://doi.org/10.1016/s1697-2600\(13\)70028-0](https://doi.org/10.1016/s1697-2600(13)70028-0).
- Hirschfeld, R. M. A., & Weissman, M. M. (2002). Risk factors for major depression and bipolar disorder. https://acnp.org/wp-content/uploads/2017/11/CH70_1017-1026.pdf
- Jocemir R. Lugon, Elias A. Warrak, Adriano S. Lugon, Bruno A. Salvador and Antonio C. L. Nobrega (2003). Revisiting Autonomic Dysfunction in End-Stage Renal Disease Patients. *Hemodialysis International*, [online] 7(3), pp.198–203. doi:<https://doi.org/10.1046/j.1492-7535.2003.00038.x>.
- John W. Brick Mental Health Foundation (2022) Move Your Mental Health: A Review of the Scientific Evidence on the Role Exercise and Physical Activity in Mental Health Available at: <https://www.johnwbrickfoundation.org/move-your-mental-health-report/>
- Joseph F. Hair Jr, William C. Black, Barry J. Babin, & Rolph E. Anderson. (2010). *Multivariate data analysis* (7th Edition) (7th ed.). Prentice Hall.
- Kementrian kesehatan RI. (2018). Riset Kesehatan Dasar (RISKESDAS) 2018. https://kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Hasil-riskesdas-2018_1274.pdf

- King-Wing Ma, T., & Kam-Tao Li, P. (2016). Depression in dialysis patients. In *Nephrology* (Vol. 21, Issue 8, pp. 639–646). Blackwell Publishing. <https://doi.org/10.1111/nep.12742>
- Kondo, K., Antick, J. R., Ayers, C. K., Kansagara, D., & Chopra, P. (2020). Depression screening tools for patients with kidney failure a systematic review. *Clinical Journal of the American Society of Nephrology*, 15(12), 1785–1795. <https://doi.org/10.2215/CJN.05540420>
- Kovesdy, C. P. (2022). Epidemiology of chronic kidney disease: an update 2022. In *Kidney International Supplements* (Vol. 12, Issue 1, pp. 7–11). Elsevier B.V. <https://doi.org/10.1016/j.kisu.2021.11.003>
- Krogh, J., Hjorthøj, C., Speyer, H., Glud, C., & Nordentoft, M. (2017). Exercise for patients with major depression: A systematic review with meta-analysis and trial sequential analysis. *BMJ Open*, 7(9). <https://doi.org/10.1136/bmjopen-2016-014820>
- Levey, A. S., Eckardt, K. U., Tsukamoto, Y., Levin, A., Coresh, J., Rossert, J., de Zeeuw, D., Hostetter, T. H., Lameire, N., Eknoyan, G., & Willis, K. (2005). Definition and classification of chronic kidney disease: A position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney International*, 67(6), 2089–2100. <https://doi.org/10.1111/j.1523-1755.2005.00365.x>
- Li, W., Liu, Y., Deng, J. and Wang, T. (2024). Influence of aerobic exercise on depression in young people: a meta-analysis. *BMC Psychiatry*, [online] 24(1). doi:<https://doi.org/10.1186/s12888-024-06013-6>.
- Lin, C. H., Hsu, Y. J., Hsu, P. H., Lee, Y. L., Lin, C. H., Lee, M. S., & Chiang, S. L. (2021). Effects of intradialytic exercise on dialytic parameters, health-related quality of life, and depression status in hemodialysis patients: A randomized controlled trial. *International Journal of Environmental Research and Public Health*, 18(17). <https://doi.org/10.3390/ijerph18179205>
- Liyanage, T., Toyama, T., Hockham, C., Ninomiya, T., Perkovic, V., Woodward, M., Fukagawa, M., Matsushita, K., Praditpornsilpa, K., Hooi, L. S., Iseki, K., Lin, M. Y., Stirnadel-Farrant, H. A., Jha, V., & Jun, M. (2022). Prevalence of chronic kidney disease in Asia: A systematic review and analysis. *BMJ Global Health*, 7(1). <https://doi.org/10.1136/bmjgh-2021-007525>

- Minanton, M., Muzaenah, T., & Sriyati, S. (2021) Latihan Intradialitik Meningkatkan Kinerja Dialisis dan Hasil Kesehatan Pasien: mini review. *Jurnal Penelitian Keperawatan Kontemporer*, 1(1), 24–33.
<https://doi.org/10.59894/jpkk.v1i1.190>
- Murdeswar, H.N. and Anjum, F. (2023) *Hemodialysis*, Nih.gov. StatPearls Publishing. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK563296/>
- Murray, I. v., & Paolini, M. A. (2023). *Histology, Kidney and Glomerulus*. Available at: <https://pubmed.ncbi.nlm.nih.gov/32119431/>
- Mutaz Foad Alradaydeh and Khalil, A.A. (2017) “The association of spiritual well-being and depression among patients receiving hemodialysis”, *Perspectives In Psychiatric Care*, 54(3), pp. 341–347. Available at: <https://doi.org/10.1111/ppc.12249>.
- Hemodialysis* (2024) *National Kidney Foundation*. Available at: <https://www.kidney.org/kidney-topics/hemodialysis>
- Neto, M.G., Filipe, Lopes, A.A., Martinez, B.P. and Micheli Bernardone Saquetto (2018). Intradialytic exercise training modalities on physical functioning and health-related quality of life in patients undergoing maintenance hemodialysis: systematic review and meta-analysis. *Clinical Rehabilitation*, [online] 32(9), pp.1189–1202. doi:<https://doi.org/10.1177/0269215518760380>.
- Noetel, M., Sanders, T., Gallardo-Gómez, D., Taylor, P., Pozo, del, Daniel, Smith, J.J., Mahoney, J., Jemima Spathis, Moresi, M., Pagano, R., Pagano, L., Vasconcellos, R., Arnott, H., Varley, B., Parker, P., Biddle, S. and Lonsdale, C. (2024). Effect of exercise for depression: systematic review and network meta-analysis of randomised controlled trials. *BMJ*, [online] pp.e075847–e075847. doi:<https://doi.org/10.1136/bmj-2023-075847>.
- Nur Aedi. (2010). pengolahan dan analisis data hasil penelitian. Fakultas Ilmu Pendidikan Universitas Pendidikan Indonesia.
- Ouzouni, Kouidi S., , E., Sioulis, A., Grekas, D., & Deligiannis, A. (2009). Effects of intradialytic exercise training on health-related quality of life indices in haemodialysis patients. *Clinical Rehabilitation*, 23(1), 53–63.
<https://doi.org/10.1177/0269215508096760>

- Park, K., Eunju Jaekal, Yoon, S., Lee, S.-H. and Choi, K.-H. (2020). Diagnostic Utility and Psychometric Properties of the Beck Depression Inventory-II Among Korean Adults. *Frontiers in Psychology*, [online] 10. doi:<https://doi.org/10.3389/fpsyg.2019.02934>.
- Putri, D. D. H., & Nugroho, I. (2023). Depression and Anxiety in Chronic Kidney Disease Patients Undergoing Hemodialysis Therapy at Hemodialysis Installations General Hospital Dr. Moewardi Surakarta. *Contagion : Scientific Periodical of Public Health and Coastal Health*, 5(4), 1478–1488. doi: <http://dx.doi.org/10.30829/contagion.v5i4.15531>
- Report Of Indonesian Renal Registry 2018. Available at: <https://www.indonesianrenalregistry.org/>.
- Rhee, S. Y., Song, J. K., Hong, S. C., Choi, J. W., Jeon, H. J., Shin, D. H., Ji, E. H., Choi, E. H., Lee, J., Kim, A., Choi, S. W., & Oh, J. (2019). Intradialytic exercise improves physical function and reduces intradialytic hypotension and depression in hemodialysis patients. *Korean Journal of Internal Medicine*, 34(3), 588–598. <https://doi.org/10.3904/kjim.2017.020>
- Saskia Heijnen, Hommel, B., Armin Kibele and Colzato, L.S. (2016) “Neuromodulation of Aerobic Exercise—A Review”, *Frontiers in Psychology*, 6. Available at: <https://doi.org/10.3389/fpsyg.2015.01890>.
- Schnaper, H. W. (2014). Remnant nephron physiology and the progression of chronic kidney disease. In *Pediatric Nephrology* (Vol. 29, Issue 2, pp. 193–202). <https://doi.org/10.1007/s00467-013-2494-8>
- Sheng, K., Zhang, P., Chen, L., Cheng, J., Wu, C., & Chen, J. (2014). Intradialytic exercise in hemodialysis patients: A systematic review and meta-analysis. *American Journal of Nephrology*, 40(5), 478–490. <https://doi.org/10.1159/000368722>
- Singh, B., Olds, T., Curtis, R., Dumuid, D., Virgara, R., Watson, A., Szeto, K., O'Connor, E., Ferguson, T., Eglitis, E., Miatke, A., Simpson, C. E. M., & Maher, C. (2023). Effectiveness of physical activity interventions for improving depression, anxiety and distress: An overview of systematic reviews. In *British Journal of Sports Medicine*. BMJ Publishing Group. <https://doi.org/10.1136/bjsports-2022-106195>

Siti Nur Rohmah, Metalia Puspitasari, Heru Prasanto, Yulia Wardhani, Iri Kuswadi and Andika Dhamarjati (2024). Effect of intradialytic aerobic exercise intervention on dialysis adequacy and quality of life in patients with end-stage kidney disease undergoing hemodialysis at Dr. Sardjito General Hospital, Indonesia. *International Urology and Nephrology*. doi:<https://doi.org/10.1007/s11255-024-04100-x>

Suhaib Muflih, Alzoubi, K.H., Sayer Al-Azzam and Belal Al-Husein (2021). Depression symptoms and quality of life in patients receiving renal replacement therapy in Jordan: A cross-sectional study. *Annals of Medicine and Surgery*, [online] 66. doi:<https://doi.org/10.1016/j.amsu.2021.102384>.

Weisbord, S. D., Fried, L. F., Arnold, R. M., Fine, M. J., Levenson, D. J., Peterson, R. A., & Switzer, G. E. (2005). Prevalence, severity, and importance of physical and emotional symptoms in chronic hemodialysis patients. *Journal of the American Society of Nephrology*, 16(8), 2487–2494. <https://doi.org/10.1681/ASN.2005020157>

Young, S.N. (2007). How to increase serotonin in the human brain without drugs. *Journal of psychiatry & neuroscience : JPN*, [online] 32(6), pp.394–9. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2077351>

Zhang, L., Wang, F., Wang, L., Wang, W., Liu, B., Liu, J., Chen, M., He, Q., Liao, Y., Yu, X., Chen, N., Zhang, J. E., Hu, Z., Liu, F., Hong, D., Ma, L., Liu, H., Zhou, X., Chen, J., ... Wang, H. (2012). Prevalence of chronic kidney disease in China: A cross-sectional survey. *The Lancet*, 379(9818), 815–822. [https://doi.org/10.1016/S0140-6736\(12\)60033-6](https://doi.org/10.1016/S0140-6736(12)60033-6)

Zimmer, P., Stritt, C., Bloch, W., Schmidt, F.-P., Sven Thorsten Hübner, Binnebößel, S., Schenk, A. and Oberste, M. (2016). The effects of different aerobic exercise intensities on serum serotonin concentrations and their association with Stroop task performance: a randomized controlled trial. *European Journal of Applied Physiology*, [online] 116(10), pp.2025–2034. doi:<https://doi.org/10.1007/s00421-016-3456-1>.