

DAFTAR PUSTAKA

- Ahn, J. H., Cho, Y. S., Cho, G. C. (2020). Elevated procalcitonin levels in patients with acetaminophen intoxication: Two case reports: A CARE-compliant article. *Medicine (United States)*.
<https://doi.org/10.1097/MD.00000000000018882>
- Ali, S. M., Basit, A., Sheikh, T., Mumtaz, S., Hydrie, M. Z. I. (2001). Diabetic Foot Ulcer - A Prospective Study. *Journal of the Pakistan Medical Association*, 51(2).
- Apelqvist, J. (2012). Diagnostics and treatment of the diabetic foot. In *Endocrine* (Vol. 41, Issue 3). <https://doi.org/10.1007/s12020-012-9619-x>
- Arya, A., Garg, S., Kumar, S., Meena, L., Tripathi, K. (2013). Estimation of lymphocyte apoptosis in patients with chronic, non healing diabetic foot ulcer. *International Journal of Medical Science and Public Health*.
<https://doi.org/10.5455/ijmsph.2013.100620132>
- Becker, K. L., Snider, R., Nylén, E. S. (2010). Procalcitonin in sepsis and systemic inflammation: A harmful biomarker and a therapeutic target. In *British Journal of Pharmacology*. <https://doi.org/10.1111/j.1476-5381.2009.00433.x>
- Benwan, K. Al, Mulla, A. Al, Rotimi, V. O. (2012). A study of the microbiology of diabetic foot infections in a teaching hospital in Kuwait. *Journal of Infection and Public Health*. <https://doi.org/10.1016/j.jiph.2011.07.004>
- Brodská, H., Drabek, T., Malicková, K., Kazda, A., Vitek, A., Zima, T., *et al.* (2009). Marked increase of procalcitonin after the administration of anti-thymocyte globulin in patients before hematopoietic stem cell transplantation does not indicate sepsis: A prospective study. *Critical Care*.
<https://doi.org/10.1186/cc7749>
- Charan, J., Biswas, T. (2013). How to calculate sample size for different study designs in medical research? In *Indian Journal of Psychological Medicine*.
<https://doi.org/10.4103/0253-7176.116232>
- Charles, P. E., Ladoire, S., Aho, S., Quenot, J. P., Doise, J. M., Prin, S., *et al.* (2008). Serum procalcitonin elevation in critically ill patients at the onset of bacteremia caused by either gram negative or gram positive bacteria. *BMC Infectious Diseases*. <https://doi.org/10.1186/1471-2334-8-3>

- Chen, W., Chen, K., Xu, Z., Hu, Y., Liu, Y., Liu, W., *et al.* (2021). Neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio predict mortality in patients with diabetic foot ulcers undergoing amputations. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. <https://doi.org/10.2147/DMSO.S284583>
- Christ-Crain, M., Müller, B. (2005). Procalcitonin in bacterial infections - Hype, hope, more or less? In *Swiss Medical Weekly*. <https://doi.org/2005/31/smw-11169>
- Clayton, W., Elasy, T. A. (2009). A review of the pathophysiology, classification, and treatment of foot ulcers in diabetic patients. In *Clinical Diabetes*. <https://doi.org/10.2337/diaclin.27.2.52>
- Dahaba, A. A., Rehak, P. H., List, W. F. (2003). Procalcitonin and C-reactive protein plasma concentrations in nonseptic uremic patients undergoing hemodialysis. *Intensive Care Medicine*. <https://doi.org/10.1007/s00134-003-1664-8>
- Frykberg, R. G. (2002). Diabetic foot ulcers: Pathogenesis and management. *American Family Physician*.
- Grace, E., Turner, R. M. (2014). Use of procalcitonin in patients with various degrees of chronic kidney disease including renal replacement therapy. *Clinical Infectious Diseases*. <https://doi.org/10.1093/cid/ciu732>
- Guthrie, H. C., Clasper, J. C. (2011). Historical origins and current concepts of wound debridement. In *Journal of the Royal Army Medical Corps*. <https://doi.org/10.1136/jramc-157-02-01>
- Hausfater, P., Garric, S., Ayed, S. Ben, Rosenheim, M., Bernard, M., Riou, B. (2002). Usefulness of procalcitonin as a marker of systemic infection in emergency department patients: A prospective study. *Clinical Infectious Diseases*. <https://doi.org/10.1086/339198>
- Hobizal, K. B., Wukich, D. K. (2012). Diabetic foot infections: Current concept review. In *Diabetic Foot and Ankle*. <https://doi.org/10.3402/dfa.v3i0.18409>
- Jafari, N. J., Firouzabadi, M. S., Izadi, M., Firouzabadi, M. S. S., Saburi, A. (2014). Can procalcitonin be an accurate diagnostic marker for the classification of diabetic foot ulcers? *International Journal of Endocrinology and Metabolism*. <https://doi.org/10.5812/ijem.13376>
- Jalilian, M., Sarbarzeh, P. A., Oubari, S. (2020). Factors related to severity of diabetic foot ulcer: A systematic review. *Diabetes, Metabolic Syndrome and*

Obesity: Targets and Therapy. <https://doi.org/10.2147/DMSO.S256243>

- Jameson, J. L., Kasper, D. L., Longo, D. L., Fauci, A. S., Hauser, S. L., Loscalzo, J. (2018). Harrison Principles of Internal Medicine 20th edition. In *McGraw-Hill Education* (Vol. 1).
- Jeandrot, A., Richard, J. L., Combescure, C., Jourdan, N., Finge, S., Rodier, M., *et al.* (2008). Serum procalcitonin and C-reactive protein concentrations to distinguish mildly infected from non-infected diabetic foot ulcers: A pilot study. *Diabetologia.* <https://doi.org/10.1007/s00125-007-0840-8>
- Jeffcoate, W., Bakker, K. (2005). World Diabetes Day: Footing the bill. In *Lancet* (Vol. 365, Issue 9470). [https://doi.org/10.1016/S0140-6736\(05\)66437-9](https://doi.org/10.1016/S0140-6736(05)66437-9)
- Jeong, D., Lee, J. H., Lee, G. Bin, Shin, K. H., Hwang, *et al.* (2023). Application of extracorporeal shockwave therapy to improve microcirculation in diabetic foot ulcers: A prospective study. *Medicine (United States).* <https://doi.org/10.1097/MD.00000000000033310>
- Jia, L., Parker, C. N., Parker, T. J., Kinnear, E. M., Derhy, P. H., Alvarado, A. M., *et al.* (2017). Incidence and risk factors for developing infection in patients presenting with uninfected diabetic foot ulcers. *PLoS ONE.* <https://doi.org/10.1371/journal.pone.0177916>
- Jiang, Y., Ran, X., Jia, L., Yang, C., Wang, P., Ma, J., *et al.* (2015). Epidemiology of type 2 diabetic foot problems and predictive factors for amputation in china. *International Journal of Lower Extremity Wounds.* <https://doi.org/10.1177/1534734614564867>
- Lauri, C., Tamminga, M., Glaudemans, A. W. J. M., Juárez Orozco, L. E., Erba, P. A., Jutte, P. C., *et al.* (2017). Detection of osteomyelitis in the diabetic foot by imaging techniques: A systematic review and meta-analysis comparing mri, white blood cell scintigraphy, and FDG-PET. *Diabetes Care.* <https://doi.org/10.2337/dc17-0532>
- Lavery, L. A., Armstrong, D. G., Wunderlich, R. P., Mohler, M. J., Wendel, C. S., Lipsky, B. A. (2006). Risk factors for foot infections in individuals with diabetes. *Diabetes Care.* <https://doi.org/10.2337/dc05-2425>
- Level, C., Chauveau, P., Delmas, Y., Lasseur, C., Pellé, G., Peuchant, E., Montaudon, D., *et al.* (2001). Procalcitonin: A new marker of inflammation in haemodialysis patients? *Nephrology Dialysis Transplantation.* <https://doi.org/10.1093/ndt/16.5.980>
- Madmoli, M., Madmoli, Y., Taqvaeinasab, H., Khodadadi, M., Darabiyan, P., Rafi,

- A. (2019). Some influential factors on severity of diabetic foot ulcers and Predisposing of limb amputation: A 7-year study on diabetic patients. *International Journal of Ayurvedic Medicine*. <https://doi.org/10.47552/ijam.v10i1.1222>
- Mallet, M., Haq, M., Tripon, S., Bernard, M., Benosman, H., Thabut, D., *et al.* (2017). Elevated procalcitonin is associated with bacterial infection during acute liver failure only when unrelated to acetaminophen intoxication. *European Journal of Gastroenterology and Hepatology*. <https://doi.org/10.1097/MEG.0000000000000862>
- Massara, M., De Caridi, G., Serra, R., Barillà, D., Cutrupi, A., Volpe, A., *et al.* (2017). The role of procalcitonin as a marker of diabetic foot ulcer infection. *International Wound Journal*. <https://doi.org/10.1111/iwj.12536>
- Meisner, M., Schmidt, J., Hüttner, H., & Tschaikowsky, K. (2000). The natural elimination rate of procalcitonin in patients with normal and impaired renal function. *Intensive Care Medicine, Supplement*. <https://doi.org/10.1007/bf02900740>
- Monteiro-Soares, M., Russell, D., Boyko, E. J., Jeffcoate, W., Mills, J. L., Morbach, S., *et al.* (2020). Guidelines on the classification of diabetic foot ulcers (IWGDF 2019). *Diabetes/Metabolism Research and Reviews*. <https://doi.org/10.1002/dmrr.3273>
- Müller, B., Harbarth, S., Stolz, D., Bingisser, R., Mueller, C., Leuppi, J., *et al.* (2007). Diagnostic and prognostic accuracy of clinical and laboratory parameters in community-acquired pneumonia. *BMC Infectious Diseases*. <https://doi.org/10.1186/1471-2334-7-10>
- Müller, B., White, J. C., Nylén, E. S., Snider, R. H., Becker, K. L., Habener, J. F. (2001). Ubiquitous Expression of the Calcitonin-I Gene in Multiple Tissues in Response to Sepsis 1 . *The Journal of Clinical Endocrinology & Metabolism*. <https://doi.org/10.1210/jcem.86.1.7089>
- Mutluoğlu, M., Uzun, G., Ipcioğlu, O. M., Sildiroglu, O., Özcan, Ö., Turhan, V., *et al.* (2011). Can procalcitonin predict bone infection in people with diabetes with infected foot ulcers? A pilot study. *Diabetes Research and Clinical Practice*. <https://doi.org/10.1016/j.diabres.2011.05.023>
- Nargis, W., Ahamed, B., Ibrahim, M. (2014). Procalcitonin versus C-reactive protein: Usefulness as biomarker of sepsis in ICU patient. *International Journal of Critical Illness and Injury Science*. <https://doi.org/10.4103/2229-5151.141356>
- Ndosi, M., Wright-Hughes, A., Brown, S., Backhouse, M., Lipsky, B. A., Bhogal,

- M., *et al.* (2018). Prognosis of the infected diabetic foot ulcer: a 12-month prospective observational study. *Diabetic Medicine*. <https://doi.org/10.1111/dme.13537>
- Paraskevas, K. I., Baker, D. M., Pompella, A., Mikhailidis, D. P. (2008). Does Diabetes Mellitus Play a Role in Restenosis and Patency Rates Following Lower Extremity Peripheral Arterial Revascularization? A Critical Overview. In *Annals of Vascular Surgery*. <https://doi.org/10.1016/j.avsg.2007.12.012>
- Park, J. H., Suh, D. H., Kim, H. J., Lee, Y. I., Kwak, I. H., Choi, G. W. (2017). Role of procalcitonin in infected diabetic foot ulcer. *Diabetes Research and Clinical Practice*. <https://doi.org/10.1016/j.diabres.2017.04.008>
- Patel, S., Srivastava, S., Singh, M. R., Singh, D. (2019). Mechanistic insight into diabetic wounds: Pathogenesis, molecular targets and treatment strategies to pace wound healing. In *Biomedicine and Pharmacotherapy*. <https://doi.org/10.1016/j.biopha.2019.108615>
- Pitocco, D., Spanu, T., Di Leo, M., Vitiello, R., Rizzi, A., Tartaglione, L., *et al.* (2019). Diabetic foot infections: A comprehensive overview. In *European Review for Medical and Pharmacological Sciences*. https://doi.org/10.26355/EURREV_201904_17471
- Richard, J.-L. (2011). New insights in diabetic foot infection. *World Journal of Diabetes*. <https://doi.org/10.4239/wjd.v2.i2.24>
- Samsudin, I., Vasikaran, S. D. (2017). Clinical utility and measurement of procalcitonin. In *Clinical Biochemist Reviews*.
- Sanniyasi, S., Balu, J., Narayanan, C. D. (2015). Fungal Infection: A Hidden Enemy in Diabetic Foot Ulcers. *Journal of Foot and Ankle Surgery (Asia Pacific)*. <https://doi.org/10.5005/jp-journals-10040-1033>
- Sastroasmoro, S. I. S. (2014). Dasar-dasar Metodologi Penelitian Klinis Edisi 5, 2014. In *Dasar-dasar Metodologi Penelitian*.
- Serban, D., Papanas, N., Dascalu, A. M., Kempler, P., Raz, I., Rizvi, A. A., *et al.* (2021). Significance of Neutrophil to Lymphocyte Ratio (NLR) and Platelet Lymphocyte Ratio (PLR) in Diabetic Foot Ulcer and Potential New Therapeutic Targets. In *International Journal of Lower Extremity Wounds*. <https://doi.org/10.1177/15347346211057742>
- Shen, C. J., Wu, M. S., Lin, K. H., Lin, W. L., Chen, H. C., Wu, J. Y., *et al.* (2013). The use of procalcitonin in the diagnosis of bone and joint infection: A systemic review and meta-analysis. In *European Journal of Clinical Microbiology and Infectious Diseases* (Vol. 32, Issue 6).

<https://doi.org/10.1007/s10096-012-1812-6>

Sitompul, Y., Budiman, B., Soebardi, S., Abdullah, M. (2014). Profil Pasien Kaki Diabetes yang Menjalani Reamputasi di Rumah Sakit Cipto Mangunkusumo Tahun 2008 -2012. *Jurnal Penyakit Dalam Indonesia*. <https://doi.org/10.7454/jpdi.v2i1.75>

Smith-Strøm, H., Iversen, M. M., Igland, J., Østbye, T., Graue, M., Skeie, S., *et al.* (2017). Severity and duration of diabetic foot ulcer (DFU) before seeking care as predictors of healing time: A retrospective cohort study. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0177176>

Somayaji, R., Elliott, J. A., Persaud, R., Lim, M., Goodman, L., Sibbald, R. G. (2017). The impact of team based interprofessional comprehensive assessments on the diagnosis and management of diabetic foot ulcers: A retrospective cohort study. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0185251>

Soreng, K., Levy, H. R. (2011). Procalcitonin: An Emerging Biomarker of Bacterial Sepsis. *Clinical Microbiology Newsletter*. <https://doi.org/10.1016/j.clinmicnews.2011.10.004>

Su, H. Y., Yang, C. Y., Ou, H. T., Chen, S. G., Chen, J. C., Ho, H. J., *et al.* (2023). Cost-effectiveness of Novel Macrophage-Regulating Treatment for Wound Healing in Patients with Diabetic Foot Ulcers from the Taiwan Health Care Sector Perspective. *JAMA Network Open*. <https://doi.org/10.1001/jamanetworkopen.2022.50639>

Tennvall, G. R., Apelqvist, J. (2004). Health-economic consequences of diabetic foot lesions. *Clinical Infectious Diseases*. <https://doi.org/10.1086/383275>

Todorova, A. S., Dimova, R. B., Chakarova, N. Y., Serdarova, M. S., Grozeva, G. G., Georgiev, G. K. (2023). Comparative Evaluation of the Diagnostic Value of Procalcitonin and hsCRP for the Presence of Mild-to-Moderate Diabetic Foot Infections. *International Journal of Lower Extremity Wounds*. <https://doi.org/10.1177/15347346211011849>

Uzun, G., Solmazgul, E., Curuksulu, H., Turhan, V., Ardic, N., Top, C. (2007). Procalcitonin as a diagnostic aid in diabetic foot infections. *Tohoku Journal of Experimental Medicine*. <https://doi.org/10.1620/tjem.213.305>

Van Asten, S. A. V., Nichols, A., La Fontaine, J., Bhavan, K., Peters, E. J. G., Lavery, L. A. (2017). The value of inflammatory markers to diagnose and monitor diabetic foot osteomyelitis. *International Wound Journal*. <https://doi.org/10.1111/iwj.12545>

- Vijayan, A. L., Ravindran, S., Saikant, R., Lakshmi, S., Kartik, R., Manoj, G. (2017). Procalcitonin: A promising diagnostic marker for sepsis and antibiotic therapy. In *Journal of Intensive Care*. <https://doi.org/10.1186/s40560-017-0246-8>
- Wang, A., Lv, G., Cheng, X., Ma, X., Wang, W., Gui, J., *et al.* (2020). Guidelines on multidisciplinary approaches for the prevention and management of diabetic foot disease (2020 edition). In *Burns and Trauma*. <https://doi.org/10.1093/BURNST/TKAA017>
- Wang, W., Zhang, X., Ge, N., Liu, J., Yuan, H., Zhang, P., *et al.* (2014). Procalcitonin testing for diagnosis and short-term prognosis in bacterial infection complicated by congestive heart failure: A multicenter analysis of 4,698 cases. *Critical Care*. <https://doi.org/10.1186/cc13181>
- Williams, D. T., Hilton, J. R., Harding, K. G. (2004). Diagnosing foot infection in diabetes. *Clinical Infectious Diseases*. <https://doi.org/10.1086/383267>
- Xu, Y., Geng, R., Meng, X., Feng, Z., Wang, X., Zhang, G., *et al.* (2024). The impact of inflammatory biomarkers on amputation rates in patients with diabetic foot ulcers. *International Wound Journal*. <https://doi.org/10.1111/iwj.14827>
- Yunus, I., Fasih, A., Wang, Y. (2018). The use of procalcitonin in the determination of severity of sepsis, patient outcomes and infection characteristics. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0206527>
- Zhang, P., Lu, J., Jing, Y., Tang, S., Zhu, D., Bi, Y. (2017). Global epidemiology of diabetic foot ulceration: a systematic review and meta-analysis†. In *Annals of Medicine*. <https://doi.org/10.1080/07853890.2016.1231932>