

DAFTAR PUSTAKA

- Adly S, Hewedi I, Mokhtar N. 2010. Clinicopathologic Significance of Molecular Classification of Breast Cancer: Relation to Nottingham Prognosis Index. *Journal of the Egyptian Nat. Center Inst*, Vol 22, No 4, December: 209-215
- Chen et all. 2010. Molecular subtype approximated by quantitative estrogen receptor, progesterone receptor and Her2 can predict the prognosis of breast cancer. *Tumori*, 96: 103-110
- DeVita, Vincent T.; Hellman, Samuel; Rosenberg, Steven A. 2005 : *Cancer: Principles & Practice of Oncology*, 7th Edition Lippincott Williams & Wilkins
- Holm K, Hegardt C, Staaf J, Christersson J.V, Jonsson G, Olsson H. 2010. Molecular subtypes of breast cancer are associated with characteristic DNA methylation patterns. *Breast Cancer Research*.
- Ontilo A.A, Engel J.M, Greenlee R.T, Mukesh B.N. Breast Cancer Subtype Based on ER/PR and Her2 Expressions : Comparison of Clinicopathologic Feature and survival, *Clinical Medicine & Research Volume 7*, number 1 / 2 : 4-13 Marshfield Clinic. Clinmedres.org
- Onitilo A.A et all. Breast Cancer Subtypes Based on ER/PR and Her2 Expression: Comparison of Clinicopathologic features and Survival. 2009. *Volume 7*, Number ½: 4-13
- Perry S, Kowalski T.L, Chang C.H. 2007. Quality of Life in Women with Breast Cancer : Benefits, Acceptability, and Utilization, *Biomed Central*
- Steiner E, Klubert D. 2008. Assessing Breast Cancer Risk in Women. *American Family Physician*. Volume 78, Number 12
- Zaha D.C, Lazar E, Lazureanu C. 2010. Clinicopathologic features and five years survival analysis in molecular subtypes of breast cancer. *Romanian Journal of Morphology and Embryology*, 51(1):85-89