



REFERENCES

- Adatara P, Afaya A, Salia SM, Afaya RA, Konlan KD, Agyabeng-Fandoh E, Agbinku E, Ayandayo EA, Boahene IG. Risk Factors Associated with Neonatal Sepsis: A Case Study at a Specialist Hospital in Ghana. *ScientificWorldJournal*. 2019 Jan 1;2019:9369051. doi: 10.1155/2019/9369051. PMID: 30692878; PMCID: PMC6332869.
- Afolabi AF, Adeyemi AS. Grand-multiparity: is it still an obstetric risk? *BMC Preg Childbirth*. (2013). doi: 10.4236/ojog.2013.34075
- Ahmed IAM. Maternal and Fetal Outcome of Grandmultipara in Comparison to Multiparous Woman in Two hospital in Khartoum State (2013).
- Alsammani MA, Ahmed SR. Grand multiparity: risk factors and outcome in a tertiary hospital: a comparative study. *Mater Sociomed*. (2015) 27: 244. doi: 10.5455/msm.2015.27.244-247.
- Amiel-Tison, C. and Pettigrew, A.G., (1991). Adaptive changes in the developing brain during intrauterine stress. *Brain Dev*, 13, pp.67-76.
- Amur S, Parekh A, Mummaneni P. Sex differences and genomics in autoimmune diseases. *J Autoimmun*. 2012;38:J254–65. doi: 10.1016/j.jaut.2011.12.001.
- Anker M. Addressing sex and gender in epidemic-prone infectious diseases. Geneva, Switzerland: World Health Organization; 2007.
- Aydiko A, Gultie T, Fetene Abebe G, Ginbeto T, Gendisha Ukke G. Determinants of late-onset neonatal sepsis among neonates admitted to the neonatal intensive care unit of Arba-Minch general hospital, southern Ethiopia. *PLoS One*. 2022 Dec 30;17(12):e0279622. doi: 10.1371/journal.pone.0279622. PMID: 36584129; PMCID: PMC9803164.
- Bæk, O., Ren, S., Brunse, A., Sangild, P.T., & Nguyen, D.N. (2020) 'Impaired Neonatal Immunity and Infection Resistance Following Fetal Growth Restriction in Preterm Pigs', *Front Immunol*, 11, 1808. doi: 10.3389/fimmu.2020.01808. PMID: 32903565; PMCID: PMC7438575.
- Bager P., Simonsen J., Ethelberg S., Frisch M. Cesarean delivery and risk of intestinal bacterial infection. *The Journal of Infectious Diseases*. 2010;201(6):898–902. doi: 10.1086/650998.
- Baker, C.J. and Kasper, D.L., (1976). Correlation of maternal antibody deficiency with susceptibility to neonatal group B streptococcal infection. *N Engl J Med*, 294, pp.753-756.
- Bang AT, Reddy HM, Baitule SB, Deshmukh MD, Bang RA. The incidence of morbidities in a cohort of neonates in rural Gadchiroli, India: seasonal and temporal variation and a hypothesis about prevention. *J Perinatol*. 2005;25(suppl 1):S18-S28.
- Bartels, D.B., Schwab, F., Geffers, C., Poets, C.F., and Gastmeier, P., (2007). Nosocomial infection in small for gestational age newborns with birth weight <1500 g: a multicentre analysis. *Arch Dis Child Fetal Neonatal Ed*, 92(6), pp.F449-53.
- Battaglia, F.C. and Lubchenco, L.O., (1967). A practical classification of newborn infants by weight and gestational age. *J Pediatr*, 71(2), pp.159-63.



- Bech CM, Stensgaard CN, Lund S, Holm-Hansen C, Brok JS, Nygaard U, Poulsen A. Risk factors for neonatal sepsis in Sub-Saharan Africa: a systematic review with meta-analysis. *BMJ Open*. 2022 Sep 1;12(9):e054491. doi: 10.1136/bmjopen-2021-054491. PMID: 36253895; PMCID: PMC9438195.
- Bekele, T., Merga, H., Tesfaye, T., et al., (2022). Predictors of mortality among neonates hospitalized with neonatal sepsis: a case control study from southern Ethiopia. *BMC Pediatr*, 22(1), p.1.
- Belinda M, Linda R, Jay K, et al. Lois sur l'âge minimum du mariage et prévalence du mariage précoce et de la maternité à l'adolescence: données d'Afrique subsaharienne. *Perspectives sexuelles sur la santé sexuelle et génésique*. 2016; 29-39.
- Bone, R.C. (1996). A continuing evolution in our understanding of the systemic inflammatory response syndrome (SIRS) and the multiple organ dysfunction syndromes (MODS). *Ann Intern Med*, 125(8), pp.680-7.
- Brundtland, G.H. (2002). Plenary Address delivered at United Nations General Assembly Special Session on Children. New York: WHO.
- Buckley, R.H. (2004). The Immunologic System and Disorders: T Lymphocytes, B Lymphocytes, and Natural Killer Cells. In: Richard E, Behrman MD, Robert M, Kliegman MD, Hal B, Jenson MD, editors. *Nelson Text Book of Pediatrics*, 17th Ed. SAUNDERS An Imprint of Elsevier Science.
- Camacho-Gonzalez, A., et al., (2013). Neonatal infections Disease: Evaluation of Neonatal Sepsis. *Pediatric Clinical North America*, 60(2), pp.367-389.
- Capasso, L., Borrelli, A., Cerullo, J., Pisanti, R., Figliuolo, C., Izzo, F., Paccone, M., Ferrara, T., Lama, S., Raimondi, F. (2014). Role of immunoglobulins in neonatal sepsis. *Transl Med UniSa*, 11, 28-33. PMID: 25674546; PMCID: PMC4309652.
- Cavazos-Rehg PA, Krauss MJ, Spitznagel EL, Bommarito K, Madden T, Olsen MA, Subramaniam H, Peipert JF, Bierut LJ. Maternal age and risk of labor and delivery complications. *Matern Child Health J*. 2015 Jun;19(6):1202-11. doi: 10.1007/s10995-014-1624-7. PMID: 25366100; PMCID: PMC4418963.
- Chang CJ, Chi H, Jim WT, Chiu NC, Chang L. Risk of infection in neonates born in accidental out-of-hospital deliveries. *PLoS One*. 2022 Feb 10;17(2):e0263825. doi: 10.1371/journal.pone.0263825. PMID: 35143570; PMCID: PMC8830727.
- Cleland JG and van Ginneken JK. Maternal education and child survival in developing countries: the search for pathways of influence. *Soc. Sci. Med*. 27, 1357, 1988.
- Cohen-Wolkowicz M, Moran C, Benjamin DK, Cotten CM, Clark RH, Benjamin DK Jr, Smith PB. Early and late onset sepsis in late preterm infants. *Pediatr Infect Dis J*. 2009 Dec;28(12):1052-6. doi: 10.1097/inf.0b013e3181acf6bd. PMID: 19953725; PMCID: PMC2798577.
- Colella, M., Frérot, A., Novais, A.R.B., et al. (2018) 'Neonatal and long-term consequences of fetal growth restriction', *Curr Pediatr Rev*, 14, 212–218. doi: 10.2174/1573396314666180712114531.
- Cortese, F., et al. (2016). Early and Late Infections in Newborns: Where Do We Stand? A Review. *Pediatr Neonatol*, 57(4), pp.265-73.



- Cunnington AJ. What's so bad about teenage pregnancy? *J Fam Plann Reprod Health Care*. 2001;27(1):36–41. doi: 10.1783/147118901101194877.
- Cutland, C.L., Lackritz, E., Alonso, A.B., Bodjick, T., Chandrasekaran, R., Lahariya, C. (2016). Low birth weight: case definition & guidelines for data collection, analysis, and presentation of maternal immunisation safety data. *Vaccine*.
- De Onis, M., Blössner, M., and Villar, J. (1998). Levels and patterns of intrauterine growth retardation in developing countries. *Eur J Clin Nutr*, 52 Suppl 1, pp.S5.
- De Rose DU, Ronchetti MP, Martini L, Rechichi J, Iannetta M, Dotta A, Auriti C. Diagnosis and Management of Neonatal Bacterial Sepsis: Current Challenges and Future Perspectives. *Tropical Medicine and Infectious Disease*. 2024; 9(9):199. <https://doi.org/10.3390/tropicalmed9090199>
- Dien Bard J, McElvania TeKippe E. Diagnosis of bloodstream infections in children. *J Clin Microbiol* 2016;54:1418e24.
- Downey LC, Smith PB, Benjamin DK Jr. Risk factors and prevention of late-onset sepsis in premature infants. *Early Hum Dev*. 2010 Jul;86 Suppl 1(Suppl 1):7-12. doi: 10.1016/j.earlhumdev.2010.01.012. Epub 2010 Jan 29. PMID: 20116186; PMCID: PMC2889164.
- Edwards, W.H., et al. (2004). The effect of prophylactic ointment therapy on nosocomial sepsis rates and skin integrity in infants with birth weights of 501 to 1000 g. *Pediatrics*, 113, pp.1195-1203.
- English, B.K., Hammond, W.P., Lewis, D.B., Brown, C.B., Wilson, C.B. (1992). Decreased granulocyte-macrophage colony-stimulating factor production by human neonatal blood mononuclear cells and T cells. *Pediatr Res*, 31, 211–26.
- Ershad, M., et al. (2019). Neonatal Sepsis. *Curr Emerg Hosp Med Rep*, 7(3), pp.83-90. doi: 10.1007/s40138-019-00188-z.
- Escalante MJ, Ceriani-Cernadas JM, D'Apremont I, Bancalari A, Webb V, Genes L, Villarroel L, Munoz E, Tapia JL; NEOCOSUR Neonatal Network. Late Onset Sepsis in Very Low Birth Weight Infants in the South American NEOCOSUR Network. *Pediatr Infect Dis J*. 2018 Oct;37(10):1022-1027. doi: 10.1097/INF.0000000000001958. PMID: 29561518.
- Fish EN. The X-files in immunity: sex-based differences predispose immune responses. *Nat Rev Immunol*. 2008;8:737–44. doi: 10.1038/nri2394.
- Fraser AM, Brockert JE, Ward RH. Association of young maternal age with adverse reproductive outcomes. *N Engl J Med*. 1995;332(17):1113–1118. doi: 10.1056/NEJM199504273321701.
- GBD 2017 Causes of Death Collaborators. (2018) 'Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the Global Burden of Disease Study 2017.'
- Gidi, N.W., et al. (2020). Comparison of neonatal outcomes of small for gestational age and appropriate for gestational age preterm infants born at 28-36 weeks of gestation: a multicentre study in Ethiopia. *BMJ Paediatr Open*, 4(1), e000740.



- Goldstein B, Giroir B, Randolph A. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med* 2005; 6: 2-8.
- Gro Harlem, B., 2002. Plenary Address delivered at United Nations General Assembly Special Session on Children. New York: WHO.
- Haque, K.N. (2005). Definitions of bloodstream infection in the newborn. *Pediatr Crit Care Med*, 6(3 Suppl), pp.S45-S49. doi:10.1097/01.PCC.0000161946.73305.0A.
- Haksari, E.L., et al., 2016. bech
ence curves of birth weight, length, and head circumference for gestational ages in Yogyakarta, Indonesia. *BMC Pediatrics*, 16(1), p.188.
- Heffner LJ. Advanced maternal age — how old is too old? *N Engl J Med*. 2004;351(19):1927–1929. doi: 10.1056/NEJMp048087.
- Herbst, A. and Kallen, K., 2007. Time between membrane rupture and delivery and septicemia in term neonates. *Obstet. Gynecol*, 110, pp.612–618.
- Hoffman MC, Jeffers S, Carter J, Duthely L, Cotter A, González-Quintero VH. Pregnancy at or beyond age 40 years is associated with an increased risk of fetal death and other adverse outcomes. *Am J Obstet Gynecol*. 2007;196(5):e11–e13. doi: 10.1016/j.ajog.2006.10.862.
- Hornik, C.P., et al., 2012. Early and late onset sepsis in very-low-birth-weight infants from a large group of neonatal intensive care units. *Early Hum Dev*, 88 Suppl 2(Suppl 2), pp.S69-74.
- Howie SR. Blood sample volumes in child health research: review of safe limits. *Bull World Health Organ* 2011;89:46e53.
- Huber, S.; Hetzer, B.; Crazzolaro, R.; Orth-Höller, D. The Correct Blood Volume for Paediatric Blood Cultures: A Conundrum? *Clin. Microbiol. Infect.* 2020, 26, 168–173.
- Iroh Tam, P.Y. and Bendel, C.M., 2017. Diagnostics for neonatal sepsis: current approaches and future directions. *Pediatr Res*, 82(4), pp.574-583. doi: 10.1038/pr.2017.134.
- Javaudin F, Hamel V, Legrand A, Goddet S, Templier F, Potiron C, et al. Unplanned out-of-hospital birth and risk factors of adverse perinatal outcome: findings from a prospective cohort. *Scand J Trauma Resusc Emerg Med*. 2019;27(1):26. doi: 10.1186/s13049-019-0600-z ; PubMed Central PMCID: PMC6397745.
- Johansson-Gudjónsdóttir, M., et al., 2019. Changes in incidence and etiology of early-onset neonatal infections 199
Here is the entry in Harvard style for the mentioned source:
- Kapur, R., Yoder, M.C., & Polin, R.A. (2006). The Immune System: Developmental Immunology. In: Martin RJ, Fanaroff AA, Walsh MC, editors. Neonatal Perinatal Medicine: Diseases of the Fetus and Infant. Elsevier Mosby.
- Karim, F., Li, S.S., Pao, D., Ying, R., & Stover, J. (2009). HIV in South Asia. *BMJ*, 338, b2141. doi: 10.1136/bmj.b2141.
- Katz, J., Lee, A.C.C., Kozuki, N., Lawn, J.E., Cousens, C., Blencowe, H., et al. (2013) 'Mortality risk in preterm and small for gestational age infants in low-



- income and middle-income countries: a pooled country analysis', *Lancet*, 82(9890), 417–25.
- Katz, J., Wu, L.A., Mullany, L.C., Coles, C.L., Lee, A.C.C., Kozuki, N., et al. (2014) 'Prevalence of small for gestational age and its mortality risk varies by choice of birth weight for gestational reference population', *PLoS ONE*, 9(3), e92074.
- Kayange, N., Kamugisha, E., Mwizamholya, D.L., Jeremiah, S., & Mshana, S.E. (2010). Predictors of positive blood culture and deaths among neonates with suspected neonatal sepsis in a tertiary hospital, Mwanza-Tanzania. *BMC Pediatrics*, 10, 39. doi: 10.1186/1471-2431-10-39.
- Kim, S.J., Kim, G.E., Park, J.H., Lee, S.L., & Kim, C.S. (2019) 'Clinical features and prognostic factors of early-onset sepsis: a 7.5-year experience in one neonatal intensive care unit', *Korean Journal of Pediatrics*, 62(1), pp. 36-41. doi: 10.3345/kjp.2018.06807. PMID: 30304900; PMCID: PMC6351802.
- King, A., Juszczak, E., Kingdom, U., Zealand, N., Haque, K., Salt, A., et al. (2011) 'Treatment of neonatal sepsis with intravenous immune globulin', *New England Journal of Medicine*, 365(13), pp. 1201–11.
- Klein JD. Adolescent pregnancy: current trends and issues. *Pediatrics*. 2005;116:281–286. doi: 10.1542/peds.2005-0999.
- Klein, J.O. (2001). Bacterial Sepsis and Meningitis. In: Remington JS, Klein JO, editors. *Infectious Diseases of the Fetus and Newborn Infant*. Philadelphia: W.B. Saunders Company. pp. 943–1010.
- Klinger G, Bromiker R, Zaslavsky-Paltiel I, Klinger S, Sokolover N, Lerner-Geva L, Reichman B, Israel Neonatal Network. Late-Onset Sepsis in Very Low Birth Weight Infants. *Pediatrics*. 2023;152(5):e2023062223. doi:10.1542/peds.2023-062223.
- Kosyreva, A. M., The Sex Differences of Morphology and Immunology of SIRS of Newborn Wistar Rats, *International Scholarly Research Notices*, 2014, 190749, 7 pages, 2014. doi:10.1155/2014/190749.
- Kwatra, G., Cunningham, M.C., Merrall, E., Adrian, P.V., Ip, M., Klugman, K.P., & Heyderman, R.S. (2014). Prevalence of maternal colonisation with Group B Streptococcus: a systematic review and meta-analysis. *Lancet Infect Dis*, 16(9), 1076-1084. doi: 10.1016/S1473-3099(16)30055-X.
- Lancaster DP, Friedman DF, Chiotos K, Sullivan KV. Blood volume required for detection of low levels and ultralow levels of organisms responsible for neonatal bacteremia by use of Bactec Peds Plus/F, Plus Aerobic/F Medium, and the BD Bactec FX System: an in vitro study. *J Clin Microbiol* 2015;53:3609e13.
- Lao TT, Ho LF. The obstetric implications of teenage pregnancy. *Hum Reprod*. 1997;12:2303–2305. doi: 10.1093/humrep/12.10.2303.
- Lazic Z, Takac I. Outcomes and risk factors for unplanned delivery at home and before arrival to the hospital. *Wien Klin Wochenschr*. 2011;123(1–2):11–4. doi: 10.1007/s00508-010-1505-z
- Lee, A.C., Katz, J., Blencowe, H., Cousens, S., Kozuki, N., Vogel, J.P., et al. (2013) 'National and regional estimates of term and preterm babies born small for gestational age in 138 low-income and middle-income countries in 2010', *Lancet Glob Health*, 1, e26–e36.



- Longo, S., Bollani, L., Decembrino, L., et al. (2013) 'Short-term and long-term sequelae in intrauterine growth retardation (IUGR)', *Journal of Maternal-Fetal & Neonatal Medicine*, 26, pp. 222–225.
- Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2095–128. doi: 10.1016/S0140-6736(12)61728-0.
- Malhotra A, Allison BJ, Castillo-Melendez M, et al. Neonatal morbidities of fetal growth restriction: pathophysiology and impact. *Front Endocrinol* 2019;10:55.
- Mariani, M., Parodi, A., Minghetti, D., Ramenghi, L.A., Palmero, C., Ugolotti, E., ... & Castagnola, E. (2022). Early and Late Onset Neonatal Sepsis: Epidemiology and Effectiveness of Empirical Antibacterial Therapy in a III Level Neonatal Intensive Care Unit. *Antibiotics*, 11(2), 284. doi: 10.3390/antibiotics11020284.
- McLelland G, McKenna L, Morgans A, Smith K. Epidemiology of unplanned out-of-hospital births attended by paramedics. *BMC Pregnancy Childbirth*. 2018;18(1):15. doi: 10.1186/s12884-017-1638-4 ; PubMed Central PMCID: PMC5759287.
- Mogollón CA, Bautista EE, Hernández-Arriaga G, et al. Factors associated with early-onset neonatal sepsis in children of Peruvian military personnel. *Electron J Gen Med* 2019; 16: em156.
- Mugadza G., Zvinavashe M., Gumbo F. Z., Pedersen B. S. Early breastfeeding initiation and incidence of neonatal sepsis in Chipinge District Zimbabwe. *International Journal of Contemporary Pediatrics*. 2018;5:1–5
- Mukhopadhyay P, Chaudhuri RN, Paul B. Hospital-based perinatal outcomes and complications in teenage pregnancy in India. *J Health Popul Nutr*. 2010;28:494–500. doi: 10.3329/jhpn.v28i5.6158.
- Munan R, Kakudji Y, Nsambi J, Mukuku O, Maleya A, et al. Accouchement chez la primipare à Lubumbashi : pronostic maternel et périnatal. *Pan African Med J*. 2017; 28: 77.
- Müller, W., Resch, B., & Hofer, N. (2015). *Definitions of SIRS and sepsis in correlation with early and late onset neonatal sepsis*. *Journal of Pediatric Intensive Care*, 01(01), 017–023. doi:10.3233/pic-2012-004.
- Nyenga AM, Mukuku O, Mutombo AK, Mpoy CW, Luboya ON, et al. Predictors of mortality in neonatal sepsis in a resource-limited setting. *J Adv Pediatr Child Health*. 2021; 4: 057-061.
- Odabasi, I.O., & Bulbul, A. (2020). Neonatal Sepsis. *Med Bull Sisli Etfal Hosp*, 54, 142-158. DOI: 10.14744/SEMB.2020.00236.
- Olivier, F., Bertelle, V., Shah, P.S., Drolet, C., & Piedboeuf, B. (2016). Association between birth route and late-onset sepsis in very preterm neonates. *Journal of Perinatology*, 36(12), 1083–1087.
- Onyedibe K. I., Utoh-Nedosa A. U., Okolo M. O., et al. Impact of socioeconomic factors on neonatal sepsis in Jos, Nigeria. *Jos Journal of Medicine*. 2011;6(2)
- Orimadegun AE, Akinbami FO, Tongo OO, Okereke JO. Comparison of neonates born outside and inside hospitals in a children emergency unit, southwest of



- Nigeria. *Pediatr Emerg Care*. 2008 Jun;24(6):354-8. doi: 10.1097/PEC.0b013e318177a73c. PMID: 18562876.
- Ovaskainen K, Ojala R, Gissler M, Luukkaala T, Tammela O. Out-of-hospital deliveries have risen involving greater neonatal morbidity: Risk factors in out-of-hospital deliveries in one University Hospital region in Finland. *Acta Paediatr*. 2015;104(12):1248–52. doi: 10.1111/apa.13117
- Oza, S., Lawn, J.E., Hogan, D.R., Mathers, C., Cousens, S.N., & the Lancet Child Mortality Study Group. (2015). Neonatal cause-of-death estimates for the early and late neonatal periods for 194 countries: 2000–2013. *Bulletin of the World Health Organization*, 93(1), 19–28. doi: 10.2471/BLT.14.139790.
- Polin, R.A., & the Committee on Fetus and Newborn. (2012). Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics*, 129, 1006–1015. doi: 10.1542/peds.2012-0541.
- Qiu, X., Lodha, A., Shah, P.S., et al. (2012). Neonatal outcomes of small for gestational age preterm infants in Canada. *American Journal of Perinatology*, 29, 87–94. doi: 10.1055/s-0031-1295647.
- Ota, E., Ganchimeg, T., Morisaki, N., et al. (2014). Risk factors and adverse perinatal outcomes among term and preterm infants born small-for-gestational-age: secondary analyses of the WHO Multi-Country Survey on Maternal and Newborn Health. *PLoS ONE*, 9(1), e105155. doi: 10.1371/journal.pone.0105155.
- Rodie VA, Thomson AJ, Norman JE. Accidental out-of-hospital deliveries: an obstetric and neonatal case control study. *Acta obstetrica et gynecologica Scandinavica*. 2002;81(1):50–4. doi: 10.1046/j.0001-6349.2001.00420.x
- Rosa-Mangeret, F., Benski, A.C., Golaz, A., Zala, P.Z., Kyokan, M., Wagner, N., Muhe, L.M., & Pfister, R.E. 2022, '2.5 Million Annual Deaths-Are Neonates in Low- and Middle-Income Countries Too Small to Be Seen? A Bottom-Up Overview on Neonatal Morbi-Mortality', *Tropical Medicine and Infectious Disease*, vol. 7, no. 5, p. 64. doi: 10.3390/tropicalmed7050064. PMID: 35622691; PMCID: PMC9148074.
- Rowe-Murray H. J., Fisher J. R. W. Baby Friendly Hospital practices: Cesarean section is a persistent barrier to early initiation of breastfeeding. *Women and Birth*. 2002;29(2):124–131. doi: 10.1046/j.1523-536X.2002.00172.x.
- Saqeeb KN, Hasan SMT, Khan MA, Ahmed T, Chisti MJ. Determinants and Outcome of Community-Acquired Late-Onset Neonatal Sepsis in Rural Bangladesh. *Glob Pediatr Health*. 2019 Mar 15;6:2333794X19833730. doi: 10.1177/2333794X19833730. PMID: 30906816; PMCID: PMC6421610.
- Schrag SJ, Cutland CL, Zell ER, et al. Risk factors for neonatal sepsis and perinatal death among infants enrolled in the prevention of perinatal sepsis trial, Soweto, South Africa. *Pediatr Infect Dis J*. 2012;31:821-826.
- Singh, M., Alsaleem, M., & Gray, C.P. Neonatal Sepsis. [Updated 2022 Sep 29]. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK531478/>.
- Singh, M. (1999). Perinatal infections. In: *Care of the Newborn*, M. Singh, Ed., pp. 198–221, New Delhi, India, 5th edition.



- Singer, M., Deutschman, C.S., Seymour, C.W., et al. (2016). The third International consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA*, 315, 801–10.
- Suwarna, N.O., Yuniati, T., Cahyadi, A.I., Achmad, T.H., & Agustian, D. (2022). Faktor Risiko Kejadian Sepsis Neonatorum Awitan Dini di Rumah Sakit Umum Pusat Dr. Hasan Sadikin Bandung. *Sari Pediatri*, 24(2), 100. DOI: 10.14238/sp24.2.2022.99-105.
- Stoll, B.J., Hansen, N., Fanaroff, A.A., Wright, L.L., Carlo, W.A., Ehrenkranz, R.A., ... & the Neonatal Research Network. (2002). Changes in pathogens causing early-onset sepsis in very-low-birth-weight infants. *New England Journal of Medicine*, 347, 240–7.
- Suhag, A., & Berghella, V. (2013). Intrauterine growth restriction (IUGR): etiology and diagnosis. *Current Obstetrics and Gynecology Reports*, 2, 102–11. doi: 10.1007/s13669-013-0041-z.
- Singer, M., Deutschman, C.S., Seymour, C.W., et al. (2016). The third International consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA*, 315, 801–10.
- Simchen, M.J., Beiner, M.E., Strauss-Liviathan, N., Dulitzky, M., Kuint, J., Mashiach, S., et al. (2000). Neonatal outcome in growth-restricted versus appropriately grown preterm infants. *American Journal of Perinatology*, 17, 187–92.
- Simonsen, K.A., Anderson-Berry, A.L., Delair, S.F., & Davies, H.D. (2014). Early-onset neonatal sepsis. *Clinical Microbiology Reviews*, 27(1), 21–47. doi: 10.1128/CMR.00031-13.
- Tanjaroen, S., Rungamornrat, S., & Thampanichawat, W. . (2024). Predictive Factors of Late-onset Sepsis in Preterm Infants and Nurses' Practices for the Prevention of Sepsis in Pediatric Intensive Care Units. *Nursing Science Journal of Thailand*, 42(1), 13–27. Retrieved from <https://he02.tcithaijo.org/index.php/ns/article/view/263863>
- Tidy C, Payne J. Gravidity and Parity Definitions. Implications in Risk Assessment. *Professional Articles* (2019).
- Tomar, P.; Garg, A.; Gupta, R.; Singh, A.; Gupta, N.K.; Upadhyay, A. Simultaneous Two-Site Blood Culture for Diagnosis of Neonatal Sepsis. *Indian Pediatr.* 2017, 54, 199–203.
- Tsai L.Y., Chen Y.L., Tsou K.I., Mu S.C. (2015) 'The Impact of Small-for-gestational-age on Neonatal Outcome Among Very-low-birth-weight Infants.' *Pediatrics & Neonatology*, 56(2), 101-107. doi: 10.1016/j.pedneo.2014.07.007.
- Utting D, Bewley S. Family planning and age-related reproductive risk. *Obstet Gynecol.* 2011;13(1):35–41
- Utomo, M.T., Sumitro, K.R., Etika, R., & Widodo, A.D.W. (2021). Current-proven neonatal sepsis in Indonesian tertiary neonatal intensive care unit: a hematological and microbiological profile. *Iranian Journal of Microbiology*, 13(3), 266-273. doi: 10.18502/ijm.v13i3.6386.



- Utomo M. T. Risk Factors of Neonatal Sepsis: A Preliminary Study in Dr. Soetomo Hospital. *Indonesian Journal of Tropical and Infectious Disease*. 2016;1(1):23–26. doi: 10.20473/ijtid.v1i1.3718.
- Utomo, M.T., Harum, N.A., Nurrosyida, K., Sampurna, M.T.A., & Aden, T.Y. (2022). The Association between Birth Route and Early/Late-onset Neonatal Sepsis in Term Infants: A Case-control Study in the NICU of a Tertiary Hospital in East Java, Indonesia. *Iranian Journal of Neonatology*, 13(4), 11–16. ISSN 2251-7510-2322-2158.
- Victoria, C. G., Huttly, S. R. A., Barros, F. C., Lombardi, C., & Vaughan, J. P. (1992). Maternal education in relation to early and late child health outcomes: Findings from a Brazilian cohort study. *Social Science & Medicine*, 34(8), 899–905. doi:10.1016/0277-9536(92)90258-r
- Wang, H. & Chow, S.-C. (2007) 'Sample Size Calculation for Comparing Proportions', *Wiley Encyclopedia of Clinical Trials*.
- Ward, N., & Levy, M. (2017). Sepsis: Definitions, Pathophysiology and the Challenge of Bedside Management. doi: 10.1007/978-3-319-48470-9.
- Ware H. Effects of maternal education, women's roles, and child care on child mortality. In *Child Survival: Strategies for Research* (Edited by Mosley WH and Chen LC). *Popul. Devl. Rev.* 10, (suppl.), 191, 1984.
- Woodford, E.C.; Dhudasia, M.B.; Puopolo, K.M.; Skerritt, L.A.; Bhavsar, M.; DeLuca, J.; Mukhopadhyay, S. Neonatal Blood Culture Inoculant Volume: Feasibility and Challenges. *Pediatr. Res.* 2021, 90, 1086–1092.
- World Health Organization. (2010). 'Health Statistics and Information Systems on Maternal Mortality'.
- World Health Organization. (2022) 'Newborn Mortality.' [Online] Available at: <https://www.who.int/news-room/fact-sheets/detail/levels-and-trends-in-child-mortality-report-2021#:~:text=Globally%202.4%20million%20children%20died,up%20from%2040%25%20in%201990> (Accessed: September 7, 2023).
- World Health Organisation, “Indicators for Assessing Infant and Young Child Feeding Practices”. Part 3: Country profiles. Geneva. 2010.
- Wynn, J.L., & Levy, O. (2010). Role of Innate Host Defenses in Susceptibility to Early Onset Neonatal Sepsis. *Clinics in Perinatology*, 37, 307–337. doi: 10.1016/j.clp.2010.03.001.
- Yachie, A., Takano, N., Ohta, K., Uehara, T., Fujita, S., Miyawaki, T., et al. (1992). Defective production of interleukin-6 in very small premature infants in response to bacterial pathogens. *Infect Immun*, 60, 749–753.
- Yogev Y, Melamed N, Bardin R, Tenenbaum-Gavish K, Ben-Shitrit G, Ben-Haroush A. Pregnancy outcome at extremely advanced maternal age. *Am J Obstet Gynecol.* 2010;203(6):e551–e558. doi: 10.1016/j.ajog.2010.07.039.
- Zaidi AK, Thaver D, Ali SA, Khan TA. Pathogens associated with sepsis in newborns and young infants in developing countries. *Pediatr Infect Dis J.* 2009;28(1 suppl):S10-S18.
- Zea-Vera, A., & Ochoa, T.J. (2015). Challenges in the diagnosis and management of neonatal sepsis. *Journal of Tropical Pediatrics*, 61(1), 1-13. doi: 10.1093/tropej/fmu079.