UNIVERSITAS
GADJAH MADA
**Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit
Ginjal Tahap Akhir yang Menjalani Hemodialisis**
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Abe, T., Kato, S., Tsuruta, Y., Sugiura, S., Katsuno, T., Kosugi, T., Tsuboi, N., Matsuo, S., & Maruyama, S. (2015). Neutrophil/lymphocyte ratio as a predictor of cardiovascular events in incident dialysis patients: a Japanese prospective cohort study. Clinical and experimental nephrology, 19(4), 718–724. https://doi.org/10.1007/s10157-014-1046-2

Adelin, P., Putra, A. E., Puspita, T. P., Triyana, R., & Puspita, D. (2024). Hubungan Kadar Low Density Lipoprotein dengan Derajat Stenosis Arteri Pasien Penyakit Jantung Koroner RSUP M Djamil Padang Tahun 2021 – 2022. *HEME: health and medical journal*. Vol. 7 (3). https://doi.org/10.33854/heme.v6i3.1625

Ahbap, E., Sakaci, T., Kara, E., Sahutoglu, T., Koc, Y., Basturk, T., Sevinc, M., Akgol, C., Kayalar, A. O., Ucar, Z. A., Bayraktar, F., & Unsal, A. (2016). Neutrophil-to-lymphocyte ratio and platelet-tolymphocyte ratio in evaluation of inflammation in end-stage renal disease. *Clinical nephrology*, 85(4), 199–208. https://doi.org/10.5414/CN108584

Ahuja, N., Bhinder, J., Nguyen, J., Langan, T., Jr, O'Brien-Irr, M., Montross, B., Khan, S., Sharma, A. M., & Harris, L. M. (2021). Venous thromboembolism in patients with COVID-19 infection: risk factors, prevention, and management. Seminars in vascular surgery, 34(3), 101–116. https://doi.org/10.1053/j.semvascsurg.2021.06.002

Al-Hafeez, H. A. A., Al-Alashkar, A. M., Mohamed, N. M. A., Zoheir, M. I. K. (2019). Associatio between Neutrophil to Lymphocyte Ratio and Inflammatory Markers in Hemodialysis Patients. *The Egyptian Journal of Hospital Medicine*. Vol. 77 (5), 5681-5689. DOI: 10.21608/ejhm.2019.62671

Anees, M., Jawed, K., Ali, Z., Khan, A. M., & Siddiqui, N. A. (2024). Association of neutrophil-to-lymphocyte ratio and hemodialysis access failure in patients with end stage renal disease: A systematic review and meta-analysis. *The journal of vascular access*, 11297298241276560. Advance online publication. https://doi.org/10.1177/11297298241276560

Ao, G., Wang, Y., Qi, X., Wang, F., & Wen, H. (2021). Association of neutrophil-to-lymphocyte ratio and risk of cardiovascular or all-cause mortality in chronic kidney disease: a meta-analysis. *Clinical and experimental nephrology*, 25(2), 157–165. https://doi.org/10.1007/s10157-020-01975-9

UNIVERSITAS GADJAH MADA
Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit Ginjal Tahap Akhir yang Menjalani Hemodialisis
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

Biswas, M., Suvarna, R., Krishnan S, V., Devasia, T., Shenoy Belle, V., & Prabhu, K. (2022). The mechanistic role of neutrophil lymphocyte ratio perturbations in the leading non communicable lifestyle diseases. F1000Research, 11, 960. https://doi.org/10.12688/f1000research.123245.1

Blokhin, I. O., & Lentz, S. R. (2013). Mechanisms of thrombosis in obesity. *Current opinion in hematology*, 20(5), 437–444. https://doi.org/10.1097/MOH.0b013e3283634443

Broeders, M. A., Tangelder, G. J., Slaaf, D. W., Reneman, R. S., & oude Egbrink, M. G. (2002). Hypercholesterolemia enhances thromboembolism in arterioles but not venules: complete reversal by L-arginine. Arteriosclerosis, thrombosis, and vascular biology, 22(4), 680–685. https://doi.org/10.1161/01.atv.0000013287.08141.74

Brahmbhatt, A., Remuzzi, A., Franzoni, M., & Misra, S. (2016). The molecular mechanisms of hemodialysis vascular access failure. *Kidney international*, 89(2), 303–316. https://doi.org/10.1016/j.kint.2015.12.019

Buonacera, A., Stancanelli, B., Colaci, M., & Malatino, L. (2022). Neutrophil to Lymphocyte Ratio: An Emerging Marker of the Relationships between the Immune System and Diseases. International journal of molecular sciences, 23(7), 3636. https://doi.org/10.3390/ijms23073636

Carobbio, A., Vannucchi, A.M., De Stefano, V. et al. (2022). Neutrophil-to-lymphocyte ratio is a novel predictor of venous thrombosis in polycythemia vera. Blood Cancer J. 12(28). https://doi.org/10.1038/s41408-022-00625-5

Carvalho, H. A., & Villar, R. C. (2018). Radiotherapy and immune response: the systemic effects of a local treatment. Clinics (Sao Paulo, Brazil), 73(suppl 1), e557s. https://doi.org/10.6061/clinics/2018/e557s

Cobo, G., Lindholm, B., & Stenvinkel, P. (2018). Chronic inflammation in end-stage renal disease and dialysis. Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association, 33(suppl_3), iii35–iii40. https://doi.org/10.1093/ndt/gfy175

Dielis A.W.J.H., Smid, M. Spronk, H.M.H., Hamulyak, K., Kroon A.A., ten Cate, H., de Leeuw, P.W. (2005). The Prothrombotic Paradox of Hypertension: Role of the Renin-Angiotensin and Kallikrein-Kinin Systems. *Hypertension*. 2005;46:1236-1242. https://doi.org/10.116110.1161/01.HYP.0000193538.20705.23

UNIVERSITAS
GADJAH MADA
**Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit
Ginjal Tahap Akhir yang Menjalani Hemodialisis**
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

Departemen Kesehatan. Direktorat Jenderal Bina Pelayanan Medik. (2008). Pedoman pelayanan hemodialisis di sarana pelayanan kesehatan. Jakarta : Departemen Kesehatan RI.

Donal N. R., Preston S. K., Lynda A. S., Joseph A. C., Susan O'Shea, William F. O. Jr., Edmund G. L. (2003). White blood cells as a novel mortality predictor in haemodialysis patients. *Nephrology Dialysis Transplantation*. Volume 18, Issue 6, June 2003, Pages 1167–1173, https://doi.org/10.1093/ndt/gfg066

Elmahdi, A. M., Gawish, R. I., Shalaan, W. R., Eldin, M. G., Gamal, N. A., & Mabrouk, M. H. (2022). Vascular access thrombosis among end-stage renal disease patients with acute COVID19 infection (a retrospective cohort study). J*ournal of The Egyptian Society of Nephrology and Transplantation* 22(3):p 168-178. https://doi.org/10.4103/jesnt.jesnt_39_21

Fauziyati, A. (2017). Global Challenge of Early Detection and Management of Chronic Kidney Disease. *J. Kedokt. dan Kesehat. Indones*. 8, 1–2.

Fernandes, C. J., Morinaga, L. T. K., Alves, J. L., Jr, Castro, M. A., Calderaro, D., Jardim, C. V. P., & Souza, R. (2019). Cancer-associated thrombosis: the when, how and why. European respiratory review : an official journal of the European Respiratory Society, 28(151), 180119. https://doi.org/10.1183/16000617.0119-2018

Foucar, K. (2006) 'Neutrophilia', in Practical diagnosis of hematologic disorders. 4th edn. Chicago, Illinois: ASCP Press, pp. 191–229.

Francis, J. L., Palmer, G. J., 3rd, Moroose, R., & Drexler, A. (2003). Comparison of bovine and porcine heparin in heparin antibody formation after cardiac surgery. The Annals of thoracic surgery, 75(1), 17–22. https://doi.org/10.1016/s0003-4975(02)04349-7

Gubensek, J., Lolic, M., Ponikvar, R., & Buturovic-Ponikvar, J. (2016). D-dimer levels in maintenance hemodialysis patients: High prevalence of positive values also in the group without predisposing diseases. Hemodialysis international. International Symposium on Home Hemodialysis, 20(2), 198–203. https://doi.org/10.1111/hdi.12371

Hall, Y. N., & Chertow, G. M. (2007). End stage renal disease. *BMJ clinical evidence*, 2007, 2002.

Hamadi, R., Sakr, F., Aridi, H., et al. (2023) Heparin-induced thrombocytopenia in chronic hemodialysis patients. *Clinical and Applied Thrombosis/Hemostasis*, 29. doi:10.1177/10760296231177993

UNIVERSITAS GADJAH MADA
Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit Ginjal Tahap Akhir yang Menjalani Hemodialisis
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

Han, B. K., Wysham, K. D., Cain, K. C., Tyden, H., Bengtsson, A. A., & Lood, C. (2020). Neutrophil and lymphocyte counts are associated with different immunopathological mechanisms in systemic lupus erythematosus. *Lupus Science & Medicine* 2020;7:e000382. doi:10.1136/lupus-2020-000382

Hashmi, M. F., Benjamin, O., Lappin, S. L., End-Stage Renal Disease. [Updated 2023 Aug 28]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK499861/

Indrastuti, B.A. (2021). Hubungan neutrophil to lymphocyte ratio (NLR) dan platelet to lymphocyte ratio (PLR) terhadap skor geriatric nutritional risk index (GNRI) dan luaran klinis pasien penyakit ginjal kronis (PGK) yang menjalani hemodialisis reguler. Skripsi thesis. Universitas Hasanuddin. http://repository.unhas.ac.id:443/id/eprint/11728

International Society of Nephrology. (2012). KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney International Supplements*. Vol. 3 (1).

Kaller, R., Arbănași, E. M., Mureșan, A. V., Voidăzan, S., Arbănași, E. M., Horváth, E., Suciu, B. A., Hosu, I., Halmaciu, I., Brinzaniuc, K., & Russu, E. (2022). The Predictive Value of Systemic Inflammatory Markers, the Prognostic Nutritional Index, and Measured Vessels' Diameters in Arteriovenous Fistula Maturation Failure. Life (Basel, Switzerland), 12(9), 1447. https://doi.org/10.3390/life12091447

Kim, J. H., Lee, H. S., Kwon, M. J., Song, Y. R., Kim, S. G., & Kim, J. K. (2023). Role of neutrophil extracellular traps in vascular access thrombosis in hemodialysis patients. *Thrombosis research*, 231, 121–127. https://doi.org/10.1016/j.thromres.2023.10.008

Li, H., Lu, X., Xiong, R., & Wang, S. (2017). High Neutrophil-to-Lymphocyte Ratio Predicts Cardiovascular Mortality in Chronic Hemodialysis Patients. *Mediators of inflammation*, 2017, 9327136. https://doi.org/10.1155/2017/9327136

Liao, J., Wei, D., Sun, C., Yang, Y., Wei, Y., & Liu, X. (2022). Prognostic value of the combination of neutrophil-to-lymphocyte ratio, monocyte-to-lymphocyte ratio and platelet-to-lymphocyte ratio on mortality in patients on maintenance hemodialysis. BMC nephrology, 23(1), 393. https://doi.org/10.1186/s12882-022-03020-1

Limijadi, E. K. S., Setyadi, A., Utami, S. B., Purohito, B., & Sofia, S. N. (2020). The correlation between neutrophil lymphocyte ratio, C-reactive protein, and serum amyloid a with the degree of stenosis in acute

UNIVERSITAS GADJAH MADA
Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit Ginjal Tahap Akhir yang Menjalani Hemodialisis
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

coronary syndrome. Open Access Maced J Med Sci. 8(B): 1234-1239. https://doi.org/10.3889/oamjms.2020..5232

Litjens, N. H., van Druningen, C. J., & Betjes, M. G. (2006). Progressive loss of renal function is associated with activation and depletion of naive T lymphocytes. Clinical immunology (Orlando, Fla.), 118(1), 83–91. https://doi.org/10.1016/j.clim.2005.09.007

López-Novoa, J. M., Martínez-Salgado, C., Rodríguez-Peña, A. B., & López-Hernández, F. J. (2010). Common pathophysiological mechanisms of chronic kidney disease: therapeutic perspectives. *Pharmacology & therapeutics*, 128(1), 61–81. https://doi.org/10.1016/j.pharmthera.2010.05.006

MacRae, J. M., Dipchand, C., Oliver, M., Moist, L., Lok, C., Clark, E., Hiremath, S., Kappel, J., Kiaii, M., Luscombe, R., Miller, L. M., & Canadian Society of Nephrology Vascular Access Work Group (2016). Arteriovenous Access Failure, Stenosis, and Thrombosis. *Canadian journal of kidney health and disease*, 3, 2054358116669126. https://doi.org/10.1177/2054358116669126

Magwenzi, S., Woodward, C., Wraith, K. S., Aburima, A., Raslan, Z., Jones, H., McNeil, C., Wheatcroft, S., Yuldasheva, N., Febbriao, M., Kearney, M., & Naseem, K. M. (2015). Oxidized LDL activates blood platelets through CD36/NOX2-mediated inhibition of the cGMP/protein kinase G signaling cascade. Blood, 125(17), 2693–2703. https://doi.org/10.1182/blood-2014-05-574491

Marra, A., Bondesan, A., Caroli, D., Grugni, G., & Sartorio, A. (2023). The neutrophil to lymphocyte ratio (NLR) positively correlates with the presence and severity of metabolic syndrome in obese adults, but not in obese children/adolescents. BMC endocrine disorders, 23(1), 121. https://doi.org/10.1186/s12902-023-01369-4

Martinod, K., & Wagner, D. D. (2014). Thrombosis: tangled up in NETs. Blood, 123(18), 2768–2776. https://doi.org/10.1182/blood-2013-10-463646

Matsumoto, Y., Shinzato, T., Amano, I., Takai, I., Kimura, Y., Morita, H., Miwa, M., Nakane, K., Yoshikai, Y., & Maeda, K. (1995). Relationship between susceptibility to apoptosis and Fas expression in peripheral blood T cells from uremic patients: a possible mechanism for lymphopenia in chronic renal failure. *Biochemical and biophysical research communications*, *215*(1), 98–105. https://doi.org/10.1006/bbrc.1995.2438

Menteri Kesehatan Republik Indonesia. (2017). Keputusan Menteri Kesehatan Republik Indonesia Nomor HK.01.07/MENKES/642/2017 tentang

UNIVERSITAS GADJAH MADA
Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit Ginjal Tahap Akhir yang Menjalani Hemodialisis
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

Pedoman Nasional Pelayanan Kedokteran Tata Laksana Penyakit Ginjal Tahap Akhir.

Meola, M., Marciello, A., Di Salle, G., & Petrucci, I. (2021). Ultrasound evaluation of access complications: Thrombosis, aneurysms, pseudoaneurysms and infections. *The journal of vascular access*, 22(1_suppl), 71–83. https://doi.org/10.1177/11297298211018062

Mims M. P. (2018). Lymphocytosis, Lymphocytopenia, Hypergammaglobulinemia, and Hypogammaglobulinemia. *Hematology*, 682–690. https://doi.org/10.1016/B978-0-323-35762-3.00049-4

Montagnana, M., Meschi, T., Borghi, L., & Lippi, G. (2011). Thrombosis and occlusion of vascular access in hemodialyzed patients. *Seminars in thrombosis and hemostasis*, *37*(8), 946–954. https://doi.org/10.1055/s-0031-1297373

Murdeshwar, H. N., Anjum, F., Hemodialysis. [Updated 2023 Apr 27]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK563296/

Mureşan, A. V., Russu, E., Arbănaşi, E. M., Kaller, R., Hosu, I., Arbănaşi, E. M., & Voidăzan, S. T. (2022). The Predictive Value of NLR, MLR, and PLR in the Outcome of End-Stage Kidney Disease Patients. *Biomedicines*, 10(6), 1272. https://doi.org/10.3390/biomedicines10061272

National Kidney Foundation. (2006). KDOQI Clinical Practice Guidelines and Clinical Practice Recommendations for 2006 Updates: Hemodialysis Adequacy, Peritoneal Dialysis Adequacy and Vascular *Access.Am J Kidney Dis*.2006;48:S1-S322

Nicolas D, Nicolas S, Hodgens A, et al. Heparin-Induced Thrombocytopenia. [Updated 2023 May 16]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK482330/

Oe, Y., & Takahashi, N. (2022). Tissue Factor, Thrombosis, and Chronic Kidney Disease. *Biomedicines*, 10(11), 2737. https://doi.org/10.3390/biomedicines10112737

Oh, D. J., Lee, J. H., Kwon, Y. E., & Choi, H. M. (2021). Relationship Between Arteriovenous Fistula Stenosis and Circulating Levels of Neutrophil Granule Proteins in Chronic Hemodialysis Patients. Annals of vascular surgery, 77, 226–235. https://doi.org/10.1016/j.avsg.2021.05.056

UNIVERSITAS
GADJAH MADA
**Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit
Ginjal Tahap Akhir yang Menjalani Hemodialisis**
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

Ouyang, J., Bajracharya, S., John, S., Wagner, J., Xu, J., Luo, Y., Thaxton, M., Salifu, M., Yap, E., & Mallappallil, M. (2022). Clotting of Hemodialysis Access in Patients with COVID-19 in an Inner-City Hospital. Nephron, 146(2), 179–184. https://doi.org/10.1159/000520174

Pasqui, E., de Donato, G., Lazzeri, E., Molino, C., Galzerano, G., Giubbolini, M., & Palasciano, G. (2022). High Neutrophil-to-Lymphocyte and Platelet-to-Lymphocyte Ratios Are Associated with a Higher Risk of Hemodialysis Vascular Access Failure. *Biomedicines*, 10(9), 2218. https://doi.org/10.3390/biomedicines10092218

Pechlivani, N., & Ajjan, R. A. (2018). Thrombosis and Vascular Inflammation in Diabetes: Mechanisms and Potential Therapeutic Targets. *Frontiers in cardiovascular medicine*, 5, 1. https://doi.org/10.3389/fcvm.2018.00001

Perhimpunan Nefrologi Indonesia. (2003). Konsensus Dialisis PERNEFRI. 1st ed.

Pernerdri. (2017). *Report of Indonesian Renal Registry*. 10th ed.

Pernefri. (2018). *Report of Indonesian Renal Registry*. 11th ed.

Pernefri. (2020). *Report of Indonesian Renal Registry*. 13th ed.

Peterson, L. (2006) 'Infectious mononucleosis and other reactive disorders of lymphocytes' & 'Lymphocytopenia', in Practical diagnosis of hematologic disorders. 4th edn. Chicago, Illinois: ASCP Press, pp. 259–273.

Pham, P. T., Miller, J. M., Demetrion, G., & Lew, S. Q. (1995). Clotting by heparin of hemoaccess for hemodialysis in an end-stage renal disease patient. American journal of kidney diseases : the official journal of the National Kidney Foundation, 25(4), 642–647. https://doi.org/10.1016/0272-6386(95)90138-8

Quencer, K. B., & Oklu, R. (2017). Hemodialysis access thrombosis. *Cardiovascular diagnosis and therapy*, 7(Suppl 3), S299–S308. https://doi.org/10.21037/cdt.2017.09.08

Rosales C. (2018). Neutrophil: A Cell with Many Roles in Inflammation or Several Cell Types?. *Frontiers in physiology*, 9, 113. https://doi.org/10.3389/fphys.2018.00113

Santoro, D., Benedetto, F., Mondello, P., Pipitò, N., Barillà, D., Spinelli, F., Ricciardi, C. A., Cernaro, V., & Buemi, M. (2014). Vascular access for hemodialysis: current perspectives. *International journal of nephrology and renovascular disease*, 7, 281–294. https://doi.org/10.2147/IJNRD.S46643

UNIVERSITAS
GADJAH MADA
**Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit
Ginjal Tahap Akhir yang Menjalani Hemodialisis**
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

Sawako Kato, et al. (2016). Neutrophil/lymphocyte ratio: A promising prognostic marker in patients with chronic kidney disease. *Inflamm Cell Signal*; 3: e683. doi: 10.14800/ics.683.

Solak, Y., Yilmaz, M. I., Sonmez, A., Saglam, M., Cakir, E., Unal, H. U., Gok, M., Caglar, K., Oguz, Y., Yenicesu, M., Karaman, M., Ay, S. A., Gaipov, A., Turk, S., Vural, A., & Carrero, J. J. (2013). Neutrophil to lymphocyte ratio independently predicts cardiovascular events in patients with chronic kidney disease. Clinical and experimental nephrology, 17(4), 532–540. https://doi.org/10.1007/s10157-012-0728-x

Stein, P. D., Hull, R. D., Kayali, F., Ghali, W. A., Alshab, A. K., & Olson, R. E. (2004). Venous thromboembolism according to age: the impact of an aging population. arch intern med, 164(20), 2260–2265. https://doi.org/10.1001/archinte.164.20.2260

Thakur, M., Junho, C. V. C., Bernhard, S. M., Schindewolf, M., Noels, H., & Döring, Y. (2023). NETs-Induced Thrombosis Impacts on Cardiovascular and Chronic Kidney Disease. Circulation research, 132(8), 933–949. https://doi.org/10.1161/CIRCRESAHA.123.321750

Usman, R., Jamil, M., & Naveed, M. (2017). High Preoperative Neutrophil-Lymphocyte Ratio (NLR) and Red Blood Cell Distribution Width (RDW) as Independent Predictors of Native Arteriovenous Fistula Failure. Annals of vascular diseases, 10(3), 205–210. https://doi.org/10.3400/avd.oa.17-00016

Vaidya, A. R., Wolska, N., Vara, D., Mailer, R. K., Schröder, K., & Pula, G. (2021). Diabetes and Thrombosis: A Central Role for Vascular Oxidative Stress. Antioxidants, 10(5), 706. https://doi.org/10.3390/antiox10050706

Wang, L., Wang, C., Jia, X., Yang, M., & Yu, J. (2020). Relationship between Neutrophil-to-Lymphocyte Ratio and Systemic Lupus Erythematosus: A Meta-analysis. Clinics (Sao Paulo, Brazil), 75, e1450. https://doi.org/10.6061/clinics/2020/e1450

Warnock, L. B., & Huang, D. (2023). Heparin. In StatPearls. StatPearls Publishing. Available from: https://pubmed.ncbi.nlm.nih.gov/30855835/

Wilkerson, W. R., & Sane, D. C. (2002). Aging and thrombosis. *Seminars in thrombosis and hemostasis*, 28(6), 555–568. https://doi.org/10.1055/s-2002-36700

Wongmahisorn Y. (2019). Role of neutrophil-to-lymphocyte ratio as a prognostic indicator for hemodialysis arteriovenous fistula failure. The journal of

UNIVERSITAS GADJAH MADA
Hubungan Antara Rasio Neutrofil-Limfosit (RNL) dengan Trombosis Akses Vaskular pada Pasien Penyakit Ginjal Tahap Akhir yang Menjalani Hemodialisis
Faiza Husna Alifia, Dr. dr. Usi Sukorini, M.Kes., Sp.PK(K); Dr. dr. Umi Solekhah Intansari, Sp.PK(K), M.Kes
Universitas Gadjah Mada, 2024 | Diunduh dari http://etd.repository.ugm.ac.id/

vascular access, 20(6), 608–614. https://doi.org/10.1177/1129729819826037

Yang, G., Shan, H., Wu, D., Li, S., Lai, Z., Zheng, F., Xiong, Z., Xiong, Z., Diao, Y., Shan, Y., Chen, Y., Wang, A., Liang, W., & Yin, Y. (2024). COVID-19 increases extracorporeal coagulation during hemodialysis associated with upregulation of vWF/FBLN5 signaling in patients with severe/critical symptoms. BMC infectious diseases, 24(1), 427. https://doi.org/10.1186/s12879-024-09245-9

Yao, Q., Axelsson, J., Stenvinkel, P., & Lindholm, B. (2004). Chronic systemic inflammation in dialysis patients: an update on causes and consequences. *ASAIO journal* (American Society for Artificial Internal Organs : 1992), 50(6), lii–lvii. https://doi.org/10.1097/01.mat.0000147958.87989.eb

Yilmaz, H., Bozkurt, A., Cakmak, M., Celik, H. T., Bilgic, M. A., Bavbek, N., & Akcay, A. (2014). Relationship between late arteriovenous fistula (AVF) stenosis and neutrophil–lymphocyte ratio (NLR) in chronic hemodialysis patients. Renal Failure, 36(9), 1390–1394. https://doi.org/10.3109/0886022X.2014.945183

Yoshitomi, R., Nakayama, M., Sakoh, T., Fukui, A., Katafuchi, E., Seki, M., Tsuda, S., Nakano, T., Tsuruya, K., & Kitazono, T. (2019). High neutrophil/lymphocyte ratio is associated with poor renal outcomes in Japanese patients with chronic kidney disease. *Renal failure*, 41(1), 238–243. https://doi.org/10.1080/0886022X.2019.1595645

Yuan, Q., Wang, J., Peng, Z. et al. (2019). Neutrophil-to-lymphocyte ratio and incident end-stage renal disease in Chinese patients with chronic kidney disease: results from the Chinese Cohort Study of Chronic Kidney Disease (C-STRIDE). *J Transl Med* 17, 86 (2019). https://doi.org/10.1186/s12967-019-1808-4

Zhang, J., Lu, X., Wang, S., & Li, H. (2021). High Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio Are Associated with Poor Survival in Patients with Hemodialysis. *BioMed research international*, 2021, 9958081. https://doi.org/10.1155/2021/9958081