

REFERENCES

- Abdel-Hamid, M., Goda, H. A., De Gobba, C., Jenssen, H., & Osman, A. (2016). Antibacterial activity of papain hydrolysed camel whey and its fractions. *International Dairy Journal*, *61*, 91-98.
- Abdelhedi, O., & Nasri, M. (2019). Basic and recent advances in marine antihypertensive peptides: Production, structure-activity relationship and bioavailability. *Trends in Food Science & Technology*, *88*, 543-557.
- Abeyrathne, E. D. N. S., Nam, K., & Ahn, D. U. (2021). Analytical methods for lipid oxidation and antioxidant capacity in food systems. *Antioxidants* *2021*, *10*, 1587.
- Ahmad, A. S., Sae-Leaw, T., Zhang, B., & Benjakul, S. (2024). Antioxidant and antimicrobial activities of ethanolic jik (*Barringtonia acutangula*) leaf extract and its application for shelf-life extension of Pacific white shrimp meat during refrigerated storage. *Food Control*, *155*, 110037.
- Alavi, F., & Ciftci, O. N. (2023). Purification and Fractionation of Bioactive Peptides Through Membrane Filtration: A Critical and Application Review. *Trends in Food Science & Technology*, *131*, 118-128.
- Azman, A. T., Isa, N. S. M., Zin, Z. M., Abdullah, M. A. A., Aidat, O., & Zainol, M. K. (2023). Protein hydrolysate from underutilized legumes: unleashing the potential for future functional foods. *Preventive nutrition and food science*, *28*(3), 209.
- Baker, R. W. (2012). *Membrane Transport Theory. Membrane Technology and Applications* (3rd ed.). John Wiley & Sons.
- Benjakul, S., & Morrissey, M. T. (1997). Protein hydrolysates from Pacific whiting solid wastes. *Journal of agricultural and food chemistry*, *45*(9), 3423-3430.
- Benjakul, S., Yarnpakdee, S., Senphan, T., Halldorsdottir, S. M., & Kristinsson, H. G. (2014). Fish protein hydrolysates: production, bioactivities, and applications. *Antioxidants and functional components in aquatic foods*, 237-281.
- Benjakul, S., Yarnpakdee, S., Senphan, T., Halldorsdottir, S. M., & Kristinsson, H. G. (2014). Fish protein hydrolysates: production, bioactivities, and applications. *Antioxidants and functional components in aquatic foods*, 237-281.
- Bigelow, C. (1967). On the average hydrophobicity of proteins and the relation between it and protein structure. *Journal of Theoretical Biology*, *16*(2), 187-211. doi:10.1016/0022-5193(67)90004-5.



- Buamard, N., Singh, A., Zhang, B., Hong, H., Singh, P., & Benjakul, S. (2023). Ethanolic extract of *Duea Ching* fruit: Extraction, characterization and its effect on the properties and storage stability of sardine surimi gel. *Foods*, *12*(8), 1635.
- Chaieb, N., González, J. L., López-Mesas, M., Bouslama, M., & Valiente, M. (2011). Polyphenols content and antioxidant capacity of thirteen faba bean (*Vicia faba L.*) genotypes cultivated in Tunisia. *Food Research International*, *44*(4), 970-977.
- Clemente, A. (2000). Enzymatic protein hydrolysates in human nutrition. *Trends in Food Science & Technology*, *11*(7), 254-262.
- Chalamaiah, M., Hemalatha, R., & Jyothirmayi, T. (2012). Fish protein hydrolysates: Proximate composition, amino acid composition, antioxidant activities and applications: A review. *Food chemistry*, *135*(4), 3020-3038.
- Clemente, A. (2000). Enzymatic protein hydrolysates in human nutrition. *Trends in Food Science & Technology*, *11*(7), 254-262.
- Dávalos, A., Miguel, M., Bartolome, B., & Lopez-Fandino, R. (2004). Antioxidant activity of peptides derived from egg white proteins by enzymatic hydrolysis. *Journal of food protection*, *67*(9), 1939-1944.
- Dhull, S. B., Kidwai, M. K., Noor, R., Chawla, P., & Rose, P. K. (2022). A review of nutritional profile and processing of faba bean (*Vicia faba L.*). *Legume Science*, *4*(3), e129.
- do Evangelho, J. A., Vanier, N. L., Pinto, V. Z., De Berrios, J. J., Dias, A. R. G., & da Rosa Zavareze, E. (2017). Black bean (*Phaseolus vulgaris L*) protein hydrolysates: Physicochemical and functional properties. *Food Chemistry*, *214*, 460-467.
- Dumoulin, L., Jacquet, N., Malumba, P., Richel, A., & Blecker, C. (2021). Dry and wet fractionation of plant proteins: How a hybrid process increases yield and impacts nutritional value of faba beans proteins. *Innovative Food Science & Emerging Technologies*, *72*, 102747.
- Felix, M., Cermeño, M., & FitzGerald, R. J. (2021). Structure and in vitro bioactive properties of O/W emulsions generated with faba bean protein hydrolysates. *Food Research International*, *150*, 110780.
- Food and Agriculture Organization of the United Nations. (2024). The State Of World Fisheries And Aquaculture 2024. Retrieved from <https://openknowledge.fao.org/items/ef79a6ba-d8df-41b9-9e87-2b6edd811511>. Accessed on 26 September 2024.
- Gani, A., Benjakul, S., & Nuthong, P. (2018). Effect of virgin coconut oil on properties of surimi gel. *Journal of Food Science and Technology*, *55*, 496-505.



- Gao, R., Yu, Q., Shen, Y., Chu, Q., Chen, G., Fen, S., Yang, M., Yuan, L., McClements, D.J., & Sun, Q. (2021). Production, bioactive properties, and potential applications of fish protein hydrolysates: Developments and challenges. *Trends in Food Science & Technology*, 110, 687-699.
- Gautam, A. R., Benjakul, S., Kadam, D., Tiwari, B., & Singh, A. (2024). Enhanced Antioxidant and Digestive Enzyme Inhibitory Activities of Pacific White Shrimp Shell Protein Hydrolysates via Conjugation with Polyphenol: Characterization and Application in Surimi Gel. *Foods*, 13(24), 4022.
- Gautam, A. R., Benjakul, S., Patil, U., Surasani, V. K. R., Nuthong, P., & Singh, A. (2025). Synergistic Effects of Partially Purified Transglutaminase from Asian Seabass Liver and Plant Protein Isolates on Gelation, Cross-Linking, Molecular Interactions and Rheological Properties of Sardine Surimi Proteins. *Food Bioscience*, 105971.
- Idowu, A. T., Benjakul, S., Sinthusamran, S., Sookchoo, P., & Kishimura, H. (2019). Protein hydrolysate from salmon frames: Production, characteristics and antioxidative activity. *Journal of Food Biochemistry*, 43(2), e12734.
- Idowu, A. T., & Benjakul, S. (2019). Bitterness of fish protein hydrolysate and its debittering prospects. *Journal of Food Biochemistry*, 43(9), e12978.
- Idowu, A. T., Benjakul, S., Sinthusamran, S., Pongsetkul, J., Sae-Leaw, T., & Sookchoo, P. (2019). Whole wheat cracker fortified with biocalcium and protein hydrolysate powders from salmon frame: characteristics and nutritional value. *Food Quality and Safety*, 3(3), 191-199.
- Idowu, A. T., Igiehon, O. O., Idowu, S., Olatunde, O. O., & Benjakul, S. (2021). Bioactivity potentials and general applications of fish protein hydrolysates. *International Journal of Peptide Research and Therapeutics*, 27, 109-118.
- Kamani, M. H., Liu, J., Fitzsimons, S. M., Fenelon, M. A., & Murphy, E. G. (2024). Determining the influence of fava bean pre-processing on extractability and functional quality of protein isolates. *Food Chemistry: X*, 21, 101200.
- Kamani, M. H., Singh, A., Nuthong, P., Fenelon, M. A., Murphy, E. G., & Benjakul, S. (2024). Insight into gelation quality of low-grade surimi as affected by fava bean protein isolate. *International Journal of Food Science and Technology*, 59(11), 8743-8754.
- Klomklao, S., & Benjakul, S. (2017). Utilization of tuna processing byproducts: Protein hydrolysate from skipjack tuna (*Katsuwonus pelamis*) viscera. *Journal of Food Processing and Preservation*, 41(3), e12970.



- Klompong, V., & Benjakul, S. (2015). Antioxidative and antimicrobial activities of the extracts from the seed coat of Bambara groundnut (*Voandzeia subterranea*). *The Royal Society of Chemistry Advances*, 5(13), 9973-9985.
- Klompong, V., Benjakul, S., Kantachote, D., & Shahidi, F. (2007). Antioxidative activity and functional properties of protein hydrolysate of yellow stripe trevally (*Selaroides leptolepis*) as influenced by the degree of hydrolysis and enzyme type. *Food chemistry*, 102(4), 1317-1327.
- Ma, Y. S., Sun, X. H., & Wang, L. T. (2015). Study on optimal conditions of alcalase enzymatic hydrolysis of soybean protein isolate. *Advance Journal of Food Science and Technology*, 9(2), 154-158.
- Martineau-Côté, D., Achouri, A., Karboune, S., & L'Hocine, L. (2022). Faba bean: an untapped source of quality plant proteins and bioactives. *Nutrients*, 14(8), 1541.
- Mishra, R., & Bisht, S. S. (2011). Antioxidants and their characterization. *J. Pharm. Res*, 4(8), 2744-2746.
- Mittal, A., Singh, A., Aluko, R. E., & Benjakul, S. (2020). Pacific white shrimp (*Litopenaeus vannamei*) shell chitosan and the conjugate with epigallocatechin gallate: Antioxidative and antimicrobial activities. *Journal of Food Biochemistry*.
- Mittal, A., Singh, A., Zhang, B., Visessanguan, W., & Benjakul, S. (2022). Chitooligosaccharide conjugates prepared using several phenolic compounds via ascorbic acid/H₂O₂ free radical grafting: Characteristics, antioxidant, antidiabetic, and antimicrobial activities. *Foods*, 11(7), 920.
- Najafian, L., & Babji, A. S. (2012). A review of fish-derived antioxidant and antimicrobial peptides: Their production, assessment, and applications. *Peptides*, 33(1), 178-185.
- Ozogul, F., Cagalj, M., Šimat, V., Ozogul, Y., Tkaczewska, J., Hassoun, A., Kaddour, A. A., Kuley, E., Rathod, N. B., & Phadke, G. G. (2021). Recent developments in valorisation of bioactive ingredients in discard/seafood processing by-products. *Trends in Food Science & Technology*, 116, 559-582.
- Park, J. W. (2005). *Surimi and surimi seafood*. CRC press.
- Petrova, I., Tolstorebrov, I., & Eikevik, T. M. (2018). Production of fish protein hydrolysates step by step: technological aspects, equipment used, major energy costs and methods of their minimizing. *International Aquatic Research*, 10, 223-241.



- Pezeshek, S., Ojagh, S. M., Rezaei, M., & Shabanpour, B. (2019). Fractionation of protein hydrolysates of fish waste using membrane ultrafiltration: investigation of antibacterial and antioxidant activities. *Probiotics and antimicrobial proteins*, 11, 1015-1022.
- Pownall, T. L., Udenigwe, C. C., & Aluko, R. E. (2010). Amino acid composition and antioxidant properties of pea seed (*Pisum sativum L.*) enzymatic protein hydrolysate fractions. *Journal of agricultural and food chemistry*, 58(8), 4712-4718.
- Raju, T. S. (2019). *Co-and post-translational modifications of therapeutic antibodies and proteins*. John Wiley & Sons. 183-202.
- Ross, K. A., T. Beta, and S. D. Arntfield. (2009). A comparative study on the phenolic acids identified and quantified in dry beans using HPLC as affected by different extraction and hydrolysis methods. *Food Chemistry*, 113.1, 336-344.
- Sae-Leaw, T., O'Callaghan, Y. C., Benjakul, S., & O'Brien, N. M. (2016). Antioxidant activities and selected characteristics of gelatin hydrolysates from seabass (*Lates calcarifer*) skin as affected by production processes. *Journal of food science and technology*, 53, 197-208.
- Sarmadi, B. H., & Ismail, A. (2010). Antioxidative peptides from food proteins: A review. *Peptides*, 31(10), 1949-1956.
- Sbroggio, M. F., Montilha, M. S., Figueiredo, V. R. G. D., Georgetti, S. R., & Kurozawa, L. E. (2016). Influence of the degree of hydrolysis and type of enzyme on antioxidant activity of okara protein hydrolysates. *Food Science and Technology*, 36, 375-381.
- Singh, A., Hong, H., & Benjakul, S. (2022). Threadfin bream surimi gel containing squid fin protein hydrolysate: Textural properties, acceptability, and volatile profile. *Journal of food science*, 87(6), 2337-2349.
- Singh, A., Kadam, D., Gautam, A. R., Rengasamy, K. R., Aluko, R. E., & Benjakul, S. (2024). Angiotensin-I-converting enzyme and renin inhibitions by antioxidant shrimp shell protein hydrolysate and ultrafiltration peptide fractions. *Food Bioscience*, 60, 104524.
- Singh, A., Putri, G. A. U., Mittal, A., Hong, H., Yesilsu, A. F., & Benjakul, S. (2022). Protein hydrolysate from splendid squid (*Loligo formosana*) fins: antioxidant, functional properties, and flavoring profile. *Turkish Journal of Fisheries and Aquatic Sciences*, 22(6).
- Sridhar, K., Inbaraj, B. S., & Chen, B. H. (2021). Recent developments on production, purification and biological activity of marine peptides. *Food Research International*, 147, 110468.



- Supmee, V., Songrak, A., Suppapan, J., & Sangthong, P. (2021). Population genetic structure of ornate threadfin bream (*Nemipterus hexodon*) in Thailand. *Tropical Life Sciences Research*, 32(1), 63.
- Sridhar, K., Inbaraj, B. S., & Chen, B. H. (2021). Recent developments on production, purification and biological activity of marine peptides. *Food Research International*, 147, 110468.
- Song, R., Wei, R., Zhang, B., & Wang, D. (2012). Optimization of the antibacterial activity of half-fin anchovy (*Setipinna taty*) hydrolysates. *Food and Bioprocess Technology*, 5, 1979-1989.
- Sahasakul, Y., Aursalung, A., Thangsiri, S., Wongchang, P., Sangkasa-Ad, P., Wongpia, A., Polpanit, A., Inthachat, W., Temviriyaaanukul, P., & Suttisansanee, U. (2022). Nutritional compositions, phenolic contents, and antioxidant potentials of ten original lineage beans in Thailand. *Foods*, 11(14), 2062.
- Tacias-Pascacio, V. G., Morellon-Sterling, R., Siar, E. H., Tavano, O., Berenguer-Murcia, A., & Fernandez-Lafuente, R. (2020). Use of Alcalase in the production of bioactive peptides: A review. *International journal of biological macromolecules*, 165, 2143-2196.
- Thongkam, P., Chaijan, M., Cheong, L. Z., & Panpipat, W. (2023). Impact of washing with antioxidant-infused soda-saline solution on gel functionality of mackerel (*Auxis thazard*) surimi. *Foods*, 12(17), 3178.
- Varelis, P., Melton, L., & Shahidi, F. (2018). *Encyclopedia of food chemistry*. Elsevier Inc.
- Walayat, N., Liu, J., Nawaz, A., Aadil, R. M., López-Pedrouso, M., & Lorenzo, J. M. (2022). Role of food hydrocolloids as antioxidants along with modern processing techniques on the surimi protein gel textural properties, developments, limitation and future perspectives. *Antioxidants*, 11(3), 486.
- Wang, C., Rao, J., Li, X., He, D., Zhang, T., Xu, J., Chen, X., Wang, L., Yuan, Y., & Zhu, X. (2023). Chickpea protein hydrolysate as a novel plant-based cryoprotectant in frozen surimi: Insights into protein structure integrity and gelling behaviors. *Food Research International*, 169, 112871.
- Wang, X., Yu, H., Xing, R., & Li, P. (2017). Characterization, preparation, and purification of marine bioactive peptides. *BioMed research international*, 2017(1), 9746720.
- Xia, E., Zhai, L., Huang, Z., Liang, H., Yang, H., Song, G., Li, W., & Tang, H. (2019). Optimization and identification of antioxidant peptide from underutilized *Dunaliella*



- salina* protein: Extraction, in vitro gastrointestinal digestion, and fractionation. *BioMed research international*, 2019(1), 6424651.
- You, L., Zhao, M., Cui, C., Zhao, H., & Yang, B. (2009). Effect of degree of hydrolysis on the antioxidant activity of loach (*Misgurnus anguillicaudatus*) protein hydrolysates. *Innovative food science & emerging technologies*, 10(2), 235-240.
- Zhang, L., Li, Q., Hong, H., & Luo, Y. (2020). Prevention of protein oxidation and enhancement of gel properties of silver carp (*Hypophthalmichthys molitrix*) surimi by addition of protein hydrolysates derived from surimi processing by-products. *Food Chemistry*, 316, 126343.
- Zhang, X., Zhang, Y., Liu, J., & Dai, Z. (2024). Effects of surimi by-product hydrolysates on processing quality and difference analysis of volatile compounds of silver carp surimi gel. *LWT*, 211, 116906.
- Zhao, X., Mei, T., & Cui, B. (2024). Effect and mechanism of different exogenous biomolecules on the thermal-induced gel properties of surimi: A review. *Journal of Food Science*, 89(12), 10266-10282.
- Zhu, B., Yang, J., Dou, J., Ning, Y., Qi, B., & Li, Y. (2024). Comparison of the physical stability, microstructure and protein-lipid co-oxidation of O/W emulsions stabilized by l-arginine/l-lysine-modified soy protein hydrolysate. *Food Chemistry*, 447, 138901.