

DAFTAR PUSTAKA

- Adindaputri, Z., Karunia, D. dan Sutantyo, D., (2023) Accuracy of mesiodistal teeth and dental arch width measurement with conventional plaster and digital model study. *JDMFS*. 8(1). doi:<https://doi.org/10.15562/jdmfs.v8i1.1450>.
- Ahmed, Z.S.H. dan Diab, B.S., (2015) The effect of nutritional status on arch width and length of primary teeth among fifteen year old students. *IJSR*. 6(4): 1139-1144. doi:<https://doi.org/10.12816/0015315>.
- Al-Dulaimy, D.A., Al-Khannaq, M. R. A., Nahidh, M., Marrapodi, M.M., Cervino, G., Ciccì, M., Minervini, G., (2023) Assessment of dental arch forms in a sample of children. *J Clin Pediatr Dent*. 47(5): 51-56.
DOI:10.22514/jocpd.2023.045
- Aljayousi, M., Al-Khateeb, S., Badran, S. dan Abu Alhajja, E.S., (2021) Correction to: Maxillary and mandibular dental arch forms in a Jordanian population with normal occlusion. *BMC Oral Health*. 21(1): 105.
doi:<https://doi.org/10.1186/s12903-021-01523-1>.
- Alpiah, D.R.A., Anindita, P.S., Juliatri., (2015) Ukuran dan Bentuk Lengkung Gigi Mandibula pada Suku Minahasa. *e-Gigi*. 3(2): 373-378.
doi:<https://doi.org/10.35790/eg.3.2.2015.9629>.
- Alvesalo, L., (2009) Human sex chromosomes in oral and craniofacial growth. *Archives of Oral Biology*. 54: S18-S24.
doi:<https://doi.org/10.1016/j.archoralbio.2008.06.004>.
- Aznar, T., Gala'na, A.F., Marin, I., Dominguez, A., (2006) Dental arch diameters and relationships to oral habits. *Angle Orthod*. 76 (3): 441-445.
doi:[https://doi.org/10.1043/0003-3219\(2006\)07](https://doi.org/10.1043/0003-3219(2006)07).
- Barnett, E.M. dan Mehta, J.D., (1970) Oral growth stage-the key to guiding occlusal development. *JADA*. 81(6): 1360-1368. doi:[https://doi.org/10.1043/0003-3219\(2006\)07](https://doi.org/10.1043/0003-3219(2006)07).
- Burdi, A.R. dan Moyers, R.E., (1988) Development of the dentition and the occlusion. Dalam: Moyers, R.E. Handbook of orthodontics 4th ed. Chicago, London: Year Book Medical Publisher. pp. 99-136.
- Carter, G.A. dan McNamara, J.A., (1998) Longitudinal Dental Arch Changes in Adults. *Am J Orthod Dentofacial Orthop*. 144(1): 88-99.
doi:[https://doi.org/10.1016/s0889-5406\(98\)70243-4](https://doi.org/10.1016/s0889-5406(98)70243-4).
- Chusida, A., Kurniawan, A., Rizky, B.N., Pribadi, S.N.P., Diva, A.T., Anandhiyah, H.D. dan Alias, A., (2024) Gonial angle and mandibular ramus height in

Surabaya population: comparison and correlation analysis in panoramic radiograph. *Majalah Kedokteran Gigi Indonesia*. 10(2):116.
doi:<https://doi.org/10.22146/majkedgiind.98837>.

Dasgupta, M., Roy, B.K., Bora, G.R., Bharali, T., (2021) Relationship between dental arch width and vertical facial morphology in multiethnic assamese adults. *Indian J Oral Health Res*. (7): 26-35.
doi:https://doi.org/10.4103/ijohr.ijohr_27_20.

Dayataka, R.P., Herawati, H., Darwis, R.S., (2019) Hubungan tingkat keparahan maloklusi dengan status karies pada remaja. *PJDRS*. 3(1): 43.
doi:<https://doi.org/10.24198/pjdrs.v2i2.22224>.

Dean, J.A., (2016) Managing the developing occlusion. Dalam: Dean, J.A., Jones, J.E., Vinson, L.A.W. (editor). McDonald and Avery's dentistry for the child and adolescent 10th ed. St. Louis: Elsevier. pp. 417.

Demirjian, A. dan Levesque, G.Y., (1980). *Sexual Differences in Dental Development and Prediction of Emergence*. *J Dent Res*. 59(7): 1110–1122. doi:10.1177/00220345800590070301.

Fleming, P., (2017) Timing orthodontic treatment: early or late?. *Aust Dent J*. 62(1): 11-19. doi:<https://doi.org/10.1177/00220345800590070301>.

Germec-Cakan, D.G., Taner, T.U., Akan, S., (2010) Arch-width and perimeter changes in patients with borderline class I malocclusion treated with extractions or without extractions with air-rotor stripping. *Am J Orthod Dentofacial Orthop*. 137(6): 734.e1-734.e7.
doi:<https://doi.org/10.1016/j.ajodo.2009.12.023>.

Gutiérrez, D.A.R., Garzón, J.S., Franco, J.Q., Botero-Mariaca, P., (2021) Anterior open bite and its relationship with dental arch dimensions and tongue position during swallowing and phonation in individuals aged 8–16 years: A retrospective case-control study. *Int Orthod*. 19(1): 107-116.
doi:<https://doi.org/10.1016/j.ortho.2020.12.005>.

Halim, H. dan Sylvia, M., (2015) Posisi gigi dipengaruhi oleh faktor ras (studi pustaka). *J Dent Indones*. 10(1): 183-187.
doi:<https://doi.org/10.14693/jdi.v10i2.419>.

Himammi, A.N. dan Hartomo, B.T., (2021) Kegunaan radiografi panoramik pada masa mixed dentition. *Jurnal Radiologi Dentomaksilofasial Indonesia (JRDI)*. 5(1): 39. doi:<https://doi.org/10.32793/jrdi.v5i1.663>.

Jain, M. dan Dhakar, N., (2013). Arch forms: An overview. *Univers. Res. J. Dent*. 3(1): 16–16. doi:<https://doi.org/10.4103/2249-9725.119045>.

- Lin, T., Hughes, T., Meade, M.J., (2023) The genetic and environmental contributions to variation in the permanent dental arch form: a twin study. *Eur J Orthod.* 45(6): 868-974.
- Maharjan, S., Rajbhandari, A., Pradhan, R., Bajracharya, M., Manandhar, P. dan Pant, B.D., (2023) Arch Form and Mathematical Ratio in Orthodontics Patients: A Descriptive Cross-sectional Study. *Journal of Institute of Medicine Nepal*, 45(2): 49–53.
doi:<https://doi.org/10.59779/jiomnepal.1267>.
- Martins-Júnior, P.A., Ramos-Jorge, M.L., de Paiva, S.M., Pereira, L.J., Marques, L.S., (2016) Premature deciduous tooth loss and orthodontic treatment need: a 6-year prospective study. *J Public Health.* 25(2). pp.173–179.
doi:<https://doi.org/10.1007/s10389-016-0775-y>.
- Mclaughlin, R.P., Bennett, J.C., Trevisi, H.J., (2001) *Systemized orthodontic treatment mechanics*. Edinburgh: Mosby. Pp. 75-76.
- Melalatoa, M.J., (1995) *Ensiklopedi suku bangsa di indonesia*. Jakarta: CV. Eka Putra. pp.96-97.
- Mazumder, P., Bahety, H., Das, A., Mahanta, P., Saikia, D., Konwar, R., (2023) Sexual dimorphism in teeth dimension and arch perimeter of individuals of four ethnic groups of northeastern India. *Cureus.* 15(4). e37905.
doi:<https://doi.org/10.7759/cureus.37905>.
- Moyers, R.E. (1988) Analysis of the dentitions and occlusion. Dalam: Moyers, R.E. *Handbook of Orthodontics* 4th ed. Chicago, London: Year Book Medical Publisher. pp. 211-246.
- Najm, A.A., Abbas, B. (2020) Evaluation of gonial angle, ramus height and bigonial width in relation to age and gender using digital panoramic radiograph. *Diyala Journal of Medicine.* 18(2):55-61. doi: 10.26505/DJM.18024950922
- Omar, H., Alhajrasi, M., Felemban, N., Hassan, A., (2018) Dental arch dimensions, form and tooth size ratio among a Saudi sample. *Saudi Med J.* 39(1): 86-91.
doi:<https://doi.org/10.15537/smj.2018.1.21035>.
- Othman, S.A., Xinwei, E.S., Lim, S.Y., Jamaludin, M., Mohamed, N.H., Yusof, Z.Y.M., Shoaib, L.A., Hussein, N.N.N., (2012) Comparison of arch form between ethnic Malays and Malaysian aborigines in peninsular Malaysia. *Korean J Orthod.* 42(1): 47-54.
doi:<https://doi.org/10.4041/kjod.2012.42.1.47>.
- Paulino, V., Paredes, V., Cibrian, R., Gandia, J.L., (2011) Dental arch changes from adolescence to adulthood in a Spanish population: A cross-sectional study.

Med Oral Patol Oral Cir Bucal. 16(4): e607-e613.
doi:<https://doi.org/10.4317/medoral.16.e607>.

Poosti, M. dan Jalali, T., (2007) Tooth size and arch dimension in uncrowded versus crowded class I malocclusion. *J Contemp Dent Pract.* 8(3): 45-52.
doi:<https://doi.org/10.5005/jcdp-8-3-45>.

Prihatiningrum, B. (2014) *Ukuran dan bentuk lengkung gigi fase gigi bercampur pada anak Suku Jawa (kajian di Kecamatan Pakem, Kabupaten Sleman, Yogyakarta)*. Yogyakarta: Tesis Fakultas Kedokteran Gigi UGM. pp 62.

Proffit, W.R. (2019) Early stage of development. Dalam: Proffit, W.R., Field, W.F., Larson, B.E., Sarver, D.M. *Contemporary orthodontics* 6th ed. Philadelphia: Elsevier. pp. 60-89.

Proffit, W.R. (2019) Concepts of Growth and Development. Dalam: Proffit, W.R., *Contemporary orthodontics* 6th ed. Philadelphia: Elsevier. pp. 34-37.

Rahardjo, P., (2012) *Ortodonti dasar edisi 2*. Surabaya: Airlangga University Press. pp. 12-13.

Rahmawati, A.D., Sudarso, I.S.R., Pramono, D., Arguni, E., (2020) Correlation between age and dental arch dimension of Javanese children. *Dent J.* 53(2): 93-98. doi:<https://doi.org/10.20473/j.djmk.v53.i2.p93-98>.

Rawlani, S., Rawlani, S., Bhowate, R., Chandak, R., Khubchandani, M., (2017) Racial characteristics of human teeth. *Int. J. Forensic Odontol.* 2(1): 38
doi:https://doi.org/10.4103/ijfo.ijfo_21_16.

Sabóia, T.M., Tannure, P.N., Luiz, R.R., Costa, M.C., Granjeiro, J.M., Küchler, E.C., Antunes, L.S., (2013) Sexual dimorphism involved on the mesiodistal and buccolingual dimensions of permanent teeth. *Dentistry 3000.* 1 (1): 2-6. doi:<https://doi.org/10.5195/d3000.2013.10>.

Samitha, S., Nagar, P., Abinaya, R., Janani, J., (2020) Comparing the arch forms between Mongoloid race and Dravidian race in 11–14-year-old children. *Int J Clin Pediatr Dent.* 13(S-1): S26-S28. doi:<https://doi.org/10.5005/jp-journals-10005-1836>.

Saghiri, M.A., Eid, J., Tang, C.K. and Freag, P., (2021) Factors influencing different types of malocclusion and arch form-A review. *J Stomatol Oral Maxillofac Surg.*122(2). pp.185–191.
doi:<https://doi.org/10.1016/j.jormas.2020.07.002>.

Sangwan, S., Chawla, H.S., Goyal, A., Gauba, K., Mohanty, U., (2011) Progressive changes in arch width from primary to early mixed dentition period: A longitudinal study. *J Indian Soc Pedop.* 29(1): 14-19.
doi:<https://doi.org/10.4103/0970-4388.79915>.

- Sharaf, R.F., Radwan, E., Salem, G.A., El-yazeed, M.A., (2022) Dental arch form and arch dimensions among a group of Egyptian children and adolescents. *Bull Natl Res Cent.* 46(1): 201. doi:<https://doi.org/10.1186/s42269-022-00887-w>.
- Shetty, S., Ratnaparkhi, I., Pereira, T., Acharya, S., Swati Gotmare dan Kamath, P., (2019). Odontometric analysis of canines to establish sexual dimorphism in an urban population. *Indian J Dent Res.* 30(6): 855–855. doi:https://doi.org/10.4103/ijdr.ijdr_75_18.
- Singh, G., (2007) Etiologi of malocclusion-local factor. Dalam: Singh, G. Textbook of orthodontics 2nd Edition. New Delhi, India: Jaypee Brothers Medical Publishers. pp.191-200.
- Singh, S., Saraf, B.G., Indushekhar, K.R., Sheoran, N., (2021) Estimation of the Intercanine Width, Intermolar Width, Arch Length, and Arch Perimeter and Its Comparison in 12–17-year-old Children of Faridabad. *Int J Clin Pediatr Dent.* 14(3): 369–375. doi:<https://doi.org/10.5005/jp-journals-10005-1957>.
- Soetjningsih, Ranuh, G., (1995) *Tumbuh kembang anak*. Jakarta: EGC. pp 1-2.
- Thilander, B., (2009) Dentoalveolar development in subjects with normal occlusion. A longitudinal study between the ages of 5 and 31 years. *Eur J Orthod.* 31(2): 109–120. doi:<https://doi.org/10.1093/ejo/cjn124>.
- Tiwari, A., Garg, A., Virang, B., Sahu, S., Shah, N., Verma, N., (2018) Archform in Orthodontics: A Review. *JOADMS.* 4(1):2018.
- Yaacob, H., Nambiar, P., Naidu (1996) Racial characteristics of human teeth with special emphasis on the Mongoloid dentition. *Malaysian J Pathol.* 18(1): 1–7.
- Yang D., Liang S., Zhang K., Gao W., Bai Y., (2019) Evaluation of growth and development of late mixed dentition upper dental arch with normal occlusion using 3-dimensional digital models. *J Healthc Eng,* 1-8. doi:<https://doi.org/10.1155/2019/4191848>.