

DAFTAR PUSTAKA

- Anindita, A.N., Setyoboedi, B., Arief, S., & Prihaningtyas, R.A. (2024). Accuracy of 2-phase abdominal ultrasound for diagnosing biliary atresia. *Bali Med. J.* 13 : 54–58.
- Arsena, H., Gunadi, Makhmudi, A., & Andi Dwihantoro (2018). AKURASI TEMUAN ULTRASONOGRAFI UNTUK MENDIAGNOSIS ATRESIA BILIER. Universitas Gadjah Mada.
- Arua, H.N., Eze, C.U., & Arua, C.L. (2021). Correlation of Sonographic Measurements of the Gallbladder and somatometric variables in Primary School Age Pupils in Nigeria. *Adv. Res. Gastroenterol. Hepatol.* 17.
- Brahee, D.D., & Lampl, B.S. (2022). Neonatal diagnosis of biliary atresia: a practical review and update. *Pediatr. Radiol.* 52 : 685–692.
- Cavallo, L., Kovar, E.M., Aqul, A., McLoughlin, L., Mittal, N.K., Rodriguez-Baez, N., *et al.* (2022). The Epidemiology of Biliary Atresia: Exploring the Role of Developmental Factors on Birth Prevalence. *J. Pediatr.* 246 : 89-94.e2.
- Choochuen, P., Kritsaneepaiboon, S., Charoonratana, V., & Sangkhathat, S. (2019a). Is “gallbladder length-to-width ratio” useful in diagnosing biliary atresia? *J. Pediatr. Surg.* 54 : 1946–1952.
- Choochuen, P., Kritsaneepaiboon, S., Charoonratana, V., & Sangkhathat, S. (2019b). Is “gallbladder length-to-width ratio” useful in diagnosing biliary atresia? *J. Pediatr. Surg.* 54 : 1946–1952.
- Dahlan, S. (2010). Besar Sampel dan Cara Pengambilan Sampel. Jakarta : Salemba Medika.
- Gautam, A.D., Yadav, R.R., Sarma, M.S., Mandelia, A., Agrawal, V., Lal, R., *et al.* (2023). Shear Wave Elastography: A Reliable Secondary Parameter for Diagnosing Biliary Atresia in Infants With Neonatal Cholestasis. *Cureus* 15.
- Ghazy, R.M., Adawy, N.M., Khedr, M.A., & Tahoun, M.M. (2018). Biliary atresia recent insight. *Egypt. Pediatr. Assoc. Gaz.* 66 : 1–8.
- Hope, W.W., Fanelli, R., Walsh, D.S., Narula, V.K., Price, R., Stefanidis, D., *et al.* (2017). SAGES clinical spotlight review: intraoperative cholangiography. *Surg. Endosc.* 31 : 2007–2016.
- Isella, V., & Dewi, M.R. (2021). Kolestasis Neonatal di Rumah Sakit Umum Daerah Wangaya, Bali. *Cermin Dunia Kedokt.* 48 : 346.
- J Richard Landis, & Koch., G.G. (1977). The Measurement of Ob_server Agreement for Categorical Data. *Biometrics* 33 : 159–174.
- Kanegawa, K., Akasaka, Y., Kitamura, E., Nishiyama, S., Muraji, T., Nishijima,

- E., *et al.* (2003). Sonographic Diagnosis of Biliary Atresia in Pediatric Patients Using the “Triangular Cord” Sign Versus Gallbladder Length and Contraction. *Am. J. Roentgenol.* 181 : 1387–1390.
- Kelly, D.A., & Davenport, M. (2007). Current management of biliary atresia. *Arch. Dis. Child.* 92 : 1132–1135.
- Kim, S.S., Kim, M.J., Lee, M.J., Yoon, C.S., Han, S.J., & Koh, H. (2014). Ultrasonographic findings of type IIIa biliary atresia. *Ultrasonography* 33 : 267–274.
- Koo, T.K., & Li, M.Y. (2016). A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J. Chiropr. Med.* 15 : 155–163.
- Lee, S.M., Cheon, J.E., Choi, Y.H., Kim, W.S., Cho, H.H., Kim, I.O., *et al.* (2015). Ultrasonographic diagnosis of biliary atresia based on a decision-making tree model. *Korean J. Radiol.* 16 : 1364–1372.
- Lefkowitz, J.H. (1998). Biliary Atresia. *Mayo Clin. Proc.* 73 : 90–95.
- Malik, A., Thanekar, U., Mourya, R., & Shivakumar, P. (2020). Recent developments in etiology and disease modeling of biliary atresia: a narrative review. *Dig. Med. Res.* 3.
- Marciani, L., Cox, E.F., Hoad, C.L., Totman, J.J., Costigan, C., Singh, G., *et al.* (2013). Effects of various food ingredients on gall bladder emptying. *Eur. J. Clin. Nutr.* 67 : 1182–1187.
- Napolitano, M., Franchi-Abella, S., Damasio, M.B., Augdal, T.A., Avni, F.E., Bruno, C., *et al.* (2021). Correction to: Practical approach to imaging diagnosis of biliary atresia, Part 1: prenatal ultrasound and magnetic resonance imaging, and postnatal ultrasound (Pediatric Radiology, (2021), 51, 2, (314–331), 10.1007/s00247-020-04840-9). *Pediatr. Radiol.* 51 : 332–333.
- Oswari, H., & Lidvihin, A.S. (2019). Pemberian Steroid untuk Meningkatkan Bilirubin Clearance pada Pasien dengan Atresia Bilier Pasca Prosedur Kasai. *Sari Pediatr.* 20 : 249.
- Ramaswamy, P.K., Jana, M., Sharma, R., Kandasamy, D., Gupta, A.K., Bhatnagar, V., *et al.* (2024). Novel Scoring Systems and Age-Based Hepatic Shear Wave Stiffness Cut-Offs for Improving Sonographic Diagnosis of Biliary Atresia. *Indian J. Pediatr.* 91 : 667–674.
- Sastroasmoro, S., & Ismael, S. (2011). Dasar-dasar metodologi penelitian klinis. Jakarta : Sagung Seto.
- Schreiber, R.A., Harpavat, S., Hulscher, J.B.F., & Wildhaber, B.E. (2022). Biliary Atresia in 2021: Epidemiology, Screening and Public Policy. *J. Clin. Med.* 11.

- Shah, R., & Savio John (2023). Cholestatic Jaundice [WWW Document]. *StatPearls*. URL <https://www.ncbi.nlm.nih.gov/books/NBK482279/> (accessed 10.8.24).
- Sherwani, P., Bolia, R., Kaushik, A., Kumar, S., Kishore, S., & Chauhan, U. (2021). An Unusual Cause of Cholestasis in an Infant: Biliary Atresia Type IIB. *Indian J. Radiol. Imaging* 31 : 689–692.
- Sugiono (2007). Statistik untuk penelitian, 11th ed. Bandung : CV Alfabeta.
- Sutowo, D.W., Gunadi, & Makhmudi, A. (2019). Faktor Prognosis Kolangitis, Hipertensi Porta, dan Varises Esofagus untuk Kesintasan Pasien Atresia Bilier pasca Operasi Kasai di RSUP Dr. Sardjito. *Repos. UGM*.
- Tan Kendrick, A.P.A., Phua, K.B., Ooi, B.C., & Looi Tan, C.E. (2003). Biliary atresia: Making the diagnosis by the gallbladder ghost triad. *Pediatr. Radiol.* 33 : 311–315.
- Vij, M., & Rela, M. (2020). Biliary atresia: Pathology, etiology and pathogenesis. *Futur. Sci. OA* 6.
- Wang, G., Zhang, N., Zhang, X., Zhou, W., Xie, X., & Zhou, L. (2021). Ultrasound characteristics combined with gamma-glutamyl transpeptidase for diagnosis of biliary atresia in infants less than 30 days. *Pediatr. Surg. Int.* 37 : 1175–1182.
- Yan, H., Liu, J., Jin, S., Du, L., Wang, Q., & Luo, Y. (2023). A novel prediction tool based on shear wave elastography, gallbladder ultrasound, and serum biomarkers for the early diagnosis of biliary atresia in infants younger than 60 days old. *Quant. Imaging Med. Surg.* 13 : 259–270.
- Zagory, J.A., Nguyen, M. V, & Wang, K.S. (2015). Recent advances in the pathogenesis and management of biliary atresia. *Curr. Opin. Pediatr.* 27 : 389–394.
- Zhang, K., Tang, Y., Zheng, Z., Tang, C., Zhu, D., Du, Q., *et al.* (2023). Value of gallbladder length-to-width ratio for diagnosis of biliary atresia by correlation with age. *Updates Surg.* 75 : 915–920.
- Zhou, L.Y., Wang, W., Shan, Q.Y., Liu, B.X., Zheng, Y.L., Xu, Z.F., *et al.* (2015). Optimizing the us diagnosis of biliary atresia with a modified triangular cord thickness and gallbladder classification1. *Radiology* 277 : 181–191.
- Zhou, W., Chen, D., Jiang, H., Shan, Q., Zhang, X., Xie, X., *et al.* (2019). Ultrasound Evaluation of Biliary Atresia Based on Gallbladder Classification: Is 4 Hours of Fasting Necessary? *J. Ultrasound Med.* 38 : 2447–2455.
- Zhou, W., & Zhou, L. (2022). Ultrasound for the diagnosis of biliary atresia: From conventional ultrasound to artificial intelligence. *Diagnostics* 12.