

DAFTAR PUSTAKA

- Adem, M., Mekonen, W., Ausman, A., Ahmed, M. & Yimer, A. 2024. Prevalence of chronic kidney disease and its associated factors among diabetes mellitus patients in Dessie Referral Hospital, South Wollo, Ethiopia. *Scientific Reports*, 14, 9229.
- Adiyanti, S. S., Yusra, Y., Immanuel, S., Aulia, D., Henrika, F. & Kumalawati, J. 2021. The Role of Neutrophil Lymphocyte Ratio, Vitamin D, and NGAL as Cardiovascular Disease Marker in Chronic Kidney Disease. *eJournal Kedokteran Indonesia*, 192-6.
- Adwaney, A., Lim, C., Blakey, S., Duncan, N. & Ashby, D. R. 2019. Central venous stenosis, access outcome and survival in patients undergoing maintenance hemodialysis. *Clinical Journal of the American Society of Nephrology*, 14, 378-384.
- Afiatin, Supriadi, R., Sarly, D., Rusanti, I., 2018. 11th Report Of Indonesian Renal Registry. Indonesian Renal Registry.
- Anees, M., Jawed, K., Ali, Z., Khan, A. M. & Siddiqui, N. A. 2024. Association of neutrophil-to-lymphocyte ratio and hemodialysis access failure in patients with end stage renal disease: A systematic review and meta-analysis. *The Journal of Vascular Access*, 11297298241276560.
- Aneez, F. A., Shariffdeen, N., Haleem, F. A., Thangarajah, B. R. & Rasaratnam, K. 2024. Correlation between neutrophil to lymphocyte ratio and platelet to lymphocyte ratio with proteinuria in different stages of chronic kidney disease. *The Egyptian Journal of Internal Medicine*, 36, 6.
- Arhuidese, I. J., Purohit, A., Elemuo, C., Parkerson, G. R., Shames, M. L. & Malas, M. B. 2020. Outcomes of autogenous fistulas and prosthetic grafts for hemodialysis access in diabetic and nondiabetic patients. *Journal of Vascular Surgery*, 72, 2088-2096.

- Arhuidese, I. J., Tran, C. K., Thayer, A., Ottinger, M. E., Calero, A. T., Malas, M. & Shames, M. 2022. Effect of active smoking on outcomes of arteriovenous fistulas and grafts for hemodialysis access. *Journal of Vascular Surgery*, 75, e48.
- Bargman, M., Skorecki, K., 2010. Chronic Kidney Disease. In: Jameson L, Loscalzo D (eds) *Harrison's Nephrology and Acid-Base Disorders*. 17th edition. McGraw Hill Companies. Pp : 113 – 129.
- Bettaieb, A., Jebali, H., Kaab, B. B., Mami, I., Mhadhbi, B., Chermiti, M., Keteb, H., Hmida, F. B., Raies, L. & Zouaghi, M. 2022. POS-995 OBESITY PARADOX IN THE ADEQUACY OF CHRONIC HEMODIALYSIS PATIENTS. *Kidney International Reports*, 7, S435-S436.
- Bhadade, R., Harde, M. & de Souza, R. 2024a. Neutrophil-to-lymphocyte Ratio as a Marker for Diagnosis and Prognostication of Sepsis. *The Journal of the Association of Physicians of India*, 72, 33-38.
- Bhadade, R., Naik, I., Harde, M. & de Souza, R. 2024b. Neutrophil-to-lymphocyte Ratio as a Marker for Diagnosis and Prognostication of Sepsis. *The Journal of the Association of Physicians of India*, 72, 33-38.
- Brahmbhatt, Akshar., Remuzzi, Andrea., Franzoni, Marco., Misra, Sanjaya. 2016. The molecular mechanisms of hemodialysis vascular access failure. *HHS Journal* ; 303-316.
- Deigendesch, N. & Stenzel, W. 2018. Acute and chronic viral infections. *Handbook of Clinical Neurology*, 145, 227-243.
- Echefu, G., Stowe, I., Lukan, A., Sharma, G., Basu-Ray, I., Guidry, L., Schellack, J. & Kumbala, D. 2023. Central vein stenosis in hemodialysis vascular access: clinical manifestations and contemporary management strategies. *Frontiers in nephrology*, 3, 1280666.
- Gao, Meiling., Wang, jing., 2022. Risk Factors of Arteriovenous Fistula Stenosis of Patients with Maintenance Hemodialysis. *Hidawi Journal* ; 4.

- García, G. G., Iyengar, A., Kaze, F., Kierans, C., Padilla-Altamira, C. & Luyckx, V. A. Sex and gender differences in chronic kidney disease and access to care around the globe. *Seminars in nephrology*, 2022. Elsevier, 101-113.
- Ge, H., Zhang, L., Zhang, W., Yuan, Q. & Xiao, X. 2024. Neutrophil-to-lymphocyte ratio predicts poor prognosis in patients with chronic kidney disease–related pulmonary hypertension: A retrospective study. *Medicine*, 103, e40161.
- Ghonemy, T. A., Farag, S. E., Soliman, S. A., Amin, E. M. & Zidan, A. A. 2016. Vascular access complications and risk factors in hemodialysis patients: A single center study. *Alexandria journal of medicine*, 52, 67-71.
- Grogan, J., Castilla, M., Lozanski, L., Griffin, A., Loth, F., Bassioun, H., 2005. Frequency of critical stenosis in primary arteriovenous fistulae before hemodialysis access: Should duplex ultrasound surveillance be the standard of care?. *J Vasc Surg* 2005;41:1000-6.
- Gupta, R., Woo, K. & Jeniann, A. Y. Epidemiology of end-stage kidney disease. *Seminars in vascular surgery*, 2021. Elsevier, 71-78.
- Han, Q., Lin, S., He, F., Zhang, R., Xie, X., Qing, F., Huang, R., Chen, C. & Yang, Q. 2022. A high neutrophil to lymphocyte ratio is associated with poor nutritional status in chronic kidney disease patients. *British Journal of Nutrition*, 128, 1990-1996.
- Higashi, Y. 2023. Smoking cessation and vascular endothelial function. *Hypertension Research*, 46, 2670-2678.
- Hsu, B.G., Tsai, J.P., 2021. Vascular calcification of chronic kidney disease: A brief review. *Tzu Chi Medical Journal*; 33 (1): 34–41
- Hustrini, N. M. 2023. Chronic kidney disease care in Indonesia: challenges and opportunities. *Acta Medica Indonesiana*, 55, 1.
- Hustrini, N. M., Susalit, E. & Rotmans, J. I. 2022. Prevalence and risk factors for chronic kidney disease in Indonesia: An analysis of the National Basic Health Survey 2018. *Journal of global health*, 12.

- Iglesias, R., Valespin, J., Ibeas, J. 2018. Handbook On Ultrasound For Vascular Access Examination. European Dialysis and Transplant Nurses Association/ European Renal Care Association (EDTNA/ERCA). Pp : 64 – 69
- Ismail, M. T., Hariawan, H., Wardhani, Y., Puspitasari, M., Artayasa, I. P. A., Ramadhan, G., Tarigan, T. & Triatmaja, R. 2021. Prevalence and Risk Factors of Arterio-Venous Fistula Obstruction on Patient with Chronic Kidney Disease. *ACI (Acta Cardiologia Indonesiana)*, 7, 5-5.
- Ismail, M.T., Hariawan, H., Wardhani, Y., Puspitasari, M., *et al.* 2021. Prevalence and Risk Factors of AV Fistula Stenosis on patient with CKD. *ACI (Acta Cardiologia Indonesiana)* (Vol.7 No.2): 23-28.
- Jacob, P. & McCafferty, K. 2023. Assessment and management of chronic kidney disease in people living with obesity. *Clinical Medicine*, 23, 353-356.
- Johansen, K. L., Chertow, G. M., Gilbertson, D. T., Ishani, A., Israni, A., Ku, E., Li, S., Li, S., Liu, J. & Obrador, G. T. 2023. US renal data system 2022 annual data report: epidemiology of kidney disease in the United States. *American journal of kidney diseases: the official journal of the National Kidney Foundation*, 81, A8.
- Kanbay, M., Solak, Y., Unal, H.U., Kurt, Y.G., Gok, M., Cetinkaya, H.,. 2014. Monocyte count/HDL cholesterol ratio and cardiovascular events in patients with chronic kidney disease. *Int Urol Nephrol*
- Kats, M., Hawxby, A., Barker, J. & Allon, M. 2007. Impact of obesity on arteriovenous fistula outcomes in dialysis patients. *Kidney international*, 71, 39-43.
- Kazemzadeh, G.H., Modaghegh, MHS, Ravari, H, Daliri M, Hoseini L, Nateghi M. 2012. Primary Patency Rate of Native AV Fistula: Long Term Follow Up. *International Journal of Clinical and Experimental Medicine* Vol.5(2); 173–178.

- KDOQI Advisory Board Members. 2006. NKF-KDOQI Clinical Practice Guidelines for Vascular Access. American Journal of Kidney Disease Supplement 1 Vol. 48; 1-322.
- Khadatkar, A., Mahakalkar, C., Pradhan, T. & Bora, A. 2016. Comparative analysis of radiocephalic versus brachiocephalic native arteriovenous fistula for hemodialysis in end stage renal disease. *Int J Res Med Sci*, 5, 171.
- Kler, A., Khambalia, H., Pondor, Z., Donne, R., Giuffrida, G., Augustine, T., Campbell, B. & van Dellen, D. 2023. Smoking is associated with a higher complication and failure rate in arteriovenous grafts for haemodialysis: A multi-centre experience. *The Journal of Vascular Access*, 24, 747-753.
- Kovesdy, C. P. 2022. Epidemiology of chronic kidney disease: an update 2022. *Kidney international supplements*, 12, 7-11.
- Ku, E., Lee, B. J., Wei, J. & Weir, M. R. 2019. Hypertension in CKD: core curriculum 2019. *American Journal of Kidney Diseases*, 74, 120-131.
- Lee, Y. H. 2023. Reassessing the role of the neutrophil-to-lymphocyte ratio in chronic kidney disease patients. *The Korean Journal of Internal Medicine*, 38, 579.
- Limono, A. B., Setyawan, Y. & Silitonga, H. T. H. 2024. Characteristics and quality of life of chronic kidney disease patients undergoing hemodialysis in Surabaya.
- Liu, B.C., Li, L., Gao, M., Wang, YJ., Yu, JR., 2008. Microinflammation is involved in the dysfunction of arteriovenous fistula in patients with maintenance hemodialysis. *Chin Med J (Engl)*;121:2157-61.
- Lok, C. E., Huber, T. S., Lee, T., Shenoy, S., Yevzlin, A. S., Abreo, K., Allon, M., Asif, A., Astor, B. C. & Glickman, M. H. 2020. KDOQI clinical practice guideline for vascular access: 2019 update. *American Journal of Kidney Diseases*, 75, S1-S164.
- Lok, C. E., Oliver, M. J., Su, J., Bhola, C., Hannigan, N. & Jassal, S. V. 2005. Arteriovenous fistula outcomes in the era of the elderly dialysis population. *Kidney international*, 67, 2462-2469.

- Lu, P., Boros, S., Chang, Q., Bindels, RJ., Hoenderop, JG., 2008. The betaglucuronidase klotho exclusively activates the epithelial Ca channels TRPV5 and TRPV6. *Nephrol Dial Transplant*. 23:3397–3402.
- Noble, R. & Taal, M. W. 2019. Epidemiology and causes of chronic kidney disease. *Medicine*, 47, 562-566.
- Nordheim, E. & Jenssen, T. G. 2021. Chronic kidney disease in patients with diabetes mellitus. *Endocrine connections*, 10, R151-R159.
- Oliva, A., Vogel, B., Sartori, S., Cao, D., Bay, B., Smith, K., Gitto, M., Dangas, G., Pocock, S. & Bhatt, D. 2024. Impact of neutrophil-to-lymphocyte ratio on clinical outcomes in patients with or without chronic kidney disease undergoing percutaneous coronary intervention. *European Heart Journal*, 45, ehae666. 1297.
- Ozaki, K., Hori, T., Ishibashi, T., Nishio, M. & Aizawa, Y. 2010. Effects of chronic cigarette smoking on endothelial function in young men. *Journal of cardiology*, 56, 307-313.
- Park, KA., Jo, MH., Han, SJ., *et al.* 2013. Features of atherosclerosis in hemodialysis patients. *Kidney Res Clin Pract* 32: 177–182.
- Pasqui, E., de Donato, G., Lazzeri, E., Molino, C., Galzerano, G., Giubbolini, M. & Palasciano, G. 2022. High neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios are associated with a higher risk of hemodialysis vascular access failure. *Biomedicines*, 10, 2218.
- Pasqui, Edoardo., Donato, Gianmarco., Lazzer, Elisa., Molino, Cecilia., Galzerano, Giuseppe., *et al.* 2022. High Neutrophil-to-Lymphocyte and Platelet-to-Lymphocyte Ratios Are Associated with a Higher Risk of Hemodialysis Vascular Access Failure. *Biomedicines* ; 2218.
- Petrovic, M., Rabrenovic, V., Plicevic, D. & Vavic, N. 2023. WCN23-1152 THE SIGNIFICANCE OF DETERMINING RATIO OF NEUTROPHIL-LYMPHOCYTES AND PLATELET-LYMPHOCYTES IN CHRONIC

KIDNEY FAILURE PATIENTS. *Kidney International Reports*, 8, S127-S128.

Plumb, T. J., Adelson, A. B., Groggel, G. C., Johanning, J. M., Lynch, T. G. & Lund, B. 2007. Obesity and hemodialysis vascular access failure. *American journal of kidney diseases*, 50, 450-454.

Quencer, K. B. & Arici, M. 2015. Arteriovenous fistulas and their characteristic sites of stenosis. *American Journal of Roentgenology*, 205, 726-734.

Quencer, K.B., Arici, M., 2015. Arteriovenous Fistulas and Their Characteristic Sites of Stenosis. *AJR*; 205:726–734

Rooijens, P., Burgmans, J., Yo, T.-I., Hop, W., De Smet, A., Van Den Dorpel, M., Fritschy, W., de Groot, H., Burger, H. & Tordoir, J. 2005. Autogenous radial-cephalic or prosthetic brachial-antecubital forearm loop AVF in patients with compromised vessels? A randomized, multicenter study of the patency of primary hemodialysis access. *Journal of vascular surgery*, 42, 481-487.

Sanjaya, A. A. K. A., Budiarta, I. B. & Sudarsa, I. W. 2024. Hubungan antara kadar serum total kolesterol, neutrophil-to-lymphocyte ratio, dan diameter pembuluh darah terhadap kejadian stenosis arteriovenous fistula tipe radiocephalic di RSUP Prof. Dr. IGNG Ngoerah Denpasar. *Intisari Sains Medis*, 15, 1312-1319.

Shankar, M., Satheesh, G., CG, S. & Reddy, G. G. 2024. Gender disparity in maintenance hemodialysis units in South India: a cross-sectional observational study. *Frontiers in Nephrology*, 4, 1322003.

Silswal, N., Touchberry, C. D., Daniel, D. R., McCarthy, D. L., Zhang, S., Andresen, J., *et al.* 2014. FGF23 directly impairs endothelium-Dependent vasorelaxation by increasing superoxide levels and reducing nitric oxide bioavailability. *Am. J. Physiol. Endocrinol. Metab.* 307, 426–436.

Sladoje-Martinovic, B., Mikolasevic, I., Bubic, I., Racki, S. & Orlic, L. 2014. Survival of chronic hemodialysis patients over 80 years of age. *Clinical interventions in aging*, 689-696.

- Suwitra, K. 2014. Penyakit Ginjal Kronik. Buku Ajar Ilmu Penyakit Dalam Edisi Keenam Jilid II ; 2159-2165.
- Wang, Z., Zhang, J., Chan, S., Cameron, A., Healy, H. G., Venuthurupalli, S. K., Tan, K.-S. & Hoy, W. E. 2019. BMI and its association with death and the initiation of renal replacement therapy (RRT) in a cohort of patients with chronic kidney disease (CKD). *BMC nephrology*, 20, 1-10.
- Weale, A. R., Bevis, P., Neary, W. D., Boyes, S., Morgan, J. D., Lear, P. A. & Mitchell, D. C. 2008. Radiocephalic and brachiocephalic arteriovenous fistula outcomes in the elderly. *Journal of vascular surgery*, 47, 144-150.
- Xia, J., Wang, L., Ma, Z., Zhong, L., Wang, Y., Gao, Y., He, L. & Su, X. 2017. Cigarette smoking and chronic kidney disease in the general population: a systematic review and meta-analysis of prospective cohort studies. *Nephrology Dialysis Transplantation*, 32, 475-487.
- Yacoub, R., Habib, H., Lahdo, A., Al Ali, R., Varjabedian, L., Atalla, G., Kassis Akl, N., Aldakheel, S., Alahdab, S. & Albitar, S. 2010. Association between smoking and chronic kidney disease: a case control study. *BMC public health*, 10, 1-6.
- Yan, Q. & Davies, M. G. 2023. Obesity drives secondary procedures to achieve access maturation in end-stage renal disease. *Journal of Vascular Surgery*, 78, 1531-1540. e4.
- Yilmaz, Hakki., Bozkurt, Alper., Cakmak, Muzaffer., Celik, Huseyin Tugrul., Bilgic, Mukadder Ayse., *et al.* 2014. Relationship between late arteriovenous fistula (AVF) stenosis and neutrophil-lymphocyte ratio (NLR) in chronic hemodialysis patients. *Informa Healthcare Journal* ; 1390-1394.
- Yin, J. & Tian, L. 2014. Joint confidence region estimation for area under ROC curve and Youden index. *Statistics in medicine*, 33, 985-1000.
- Yoshitomi, R., Nakayama, M., Sakoh, T., Fukui, A., Katafuchi, E., Seki, M., Tsuda, S., Nakano, T., Tsuruya, K. & Kitazono, T. 2019. High

neutrophil/lymphocyte ratio is associated with poor renal outcomes in Japanese patients with chronic kidney disease. *Renal failure*, 41, 238-243.

Yuan, Q., Wang, J., Peng, Z., Zhou, Q., Xiao, X., Xie, Y., Wang, W., Huang, L., Tang, W. & Sun, D. 2019. Neutrophil-to-lymphocyte ratio and incident end-stage renal disease in Chinese patients with chronic kidney disease: results from the Chinese Cohort Study of Chronic Kidney Disease (C-STRIDE). *Journal of translational medicine*, 17, 1-8.

Zainuddin AA, Faqih DM, Trisna DV, Waluyo DA, Ekayanti F, Hariyani I, Hendarto J, Paranadipa M., 2004. Panduan praktik klinis bagi dokter di fasilitas pelayanan kesehatan primer.

Zhao, W.-M., Tao, S.-M. & Liu, G.-L. 2020. Neutrophil-to-lymphocyte ratio in relation to the risk of all-cause mortality and cardiovascular events in patients with chronic kidney disease: a systematic review and meta-analysis. *Renal failure*, 42, 1059-1066.