

DAFTAR PUSTAKA

- Akhter Ansari, W. *et al.* (2019) 'Influence of Drought Stress on Morphological, Physiological and Biochemical Attributes of Plants: A Review', *Biosciences, Biotechnology Research Asia*, 16(04), pp. 697–709. doi: 10.13005/bbra/2785.
- Ammar, M.H.; Anwar, F.; El-Harty, E.H.; Migdadi, H.M.; Abdel-Khalik, S.M.; Al-Faifi, S.A.; Farooq, M.; Alghamdi, S.S. Physiological and Yield Responses of Faba Bean (*Vicia faba* L.) to Drought Stress in Managed and Open Field Environments. *J. Agron. Crop Sci.* 2015, 201, 280–287.
- Apriyani, M. *et al.* (2020) 'AGROPROSS National Conference Proceedings of Agriculture Proceedings: Peran Teaching Factory Di Perguruan Tinggi Vokasi Dalam Mendukung Ketahanan Pangan Pada Era New Normal Pengaruh Kondisi Kapasitas Lapang Yang Berbeda Terhadap Pertumbuhan Vegetatif', pp. 8–9.
- Aroca, R. (2013) Plant responses to drought stress. From morphological to molecular feature, por Ricardo Aroca, 2012. *Springer Ecosistemas*, 22(1):93. doi: 10.1007/978-3-642-32653-0.
- Bradford, W. D. *et al.* (2005) 'An inexpensive gel electrophoresis-based polymerase chain reaction method for quantifying mRNA levels', *Cell Biology Education*, 4(SUMMER), pp. 157–168. doi: 10.1187/cbe.04-09-0051.
- Cardoso AA, Brodribb TJ, Kane CN, DaMatta FM, McAdam SAM. 2020. Osmotic adjustment and hormonal regulation of stomatal responses to vapour pressure deficit in sunflower. *AoB Plants* 12:plaa025.
- Cheng, M. C. *et al.* (2013) 'The arabidopsis ETHYLENE RESPONSE FACTOR1 Regulates abiotic stress-responsive gene expression by binding to different cis-acting elements in response to different stress signals', *Plant Physiology*, 162(3), pp. 1566–1582. doi: 10.1104/pp.113.221911.
- Conesa, M.R.; de la Rosa, J.M.; Domingo, R.; Bañon, S.; Pérez-Pastor, A. Changes Induced by Water Stress on Water Relations, Stomatal Behaviour and Morphology of Table Grapes (Cv. Crimson Seedless) Grown in Pots. *Sci. Hort.* 2016, 202, 9–16.
- Costa-Júnior, J. M. *et al.* (2022) 'Paternal exercise improves the metabolic health of offspring via epigenetic modulation of the germline', *International Journal of Molecular Sciences*, 23(1), pp. 1–15. doi: 10.3390/ijms23010001.
- Dama, H., Aisyah, S. I. and Dewi, A. K. (2018) 'Respon Kerapatan Stomata dan Kandungan Klorofil Padi (*Oryza sativa* L.) Mutan terhadap Toleransi Kekeringan Response of

- Stomata Density and Chlorophyll Content Rice (*Oryza sativa* L.) Mutants to Drought Tolerance', *Jurnal Ilmiah Aplikasi Isotop dan Radiasi*, 16(1), pp. 1–6.
- Daniell, H., Streatfield, S. J. and Wycoff, K. (2001) '1-s2.0-S1360138501019227-main', *Trends in Plant Science*, 6(5), pp. 219–226.
- Dewey, R. E. and Xie, J. (2013) 'Molecular genetics of alkaloid biosynthesis in *Nicotiana tabacum*', *Phytochemistry*, 94, pp. 10–27. doi: 10.1016/j.phytochem.2013.06.002.
- Djajadi, D. (2015) 'Tobacco Diversity in Indonesia', *Berkala Penelitian Hayati*, 20(2), pp. 27–32. doi: 10.23869/bphjbr.20.2.20155.
- Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A, et al. Crop production under drought and heat stress: plant responses and management options. *Front Plant Sci*. 2017;8(June):1–16
- Fathi A, Tari DB. Effect of drought stress and its mechanisms in plants. *Int J Life Sci*. 2016;10(1):1–6.
- Häkkinen, S. T. *et al.* (2007) 'Functional characterisation of genes involved in pyridine alkaloid biosynthesis in tobacco', *Phytochemistry*, 68(22–24), pp. 2773–2785. doi: 10.1016/j.phytochem.2007.09.010.
- Hashimoto T. and Y. Yamada. 1994. Alkaloid biogenesis: Molecular aspects. *Annual Review of Plant Physiology and Plant Molecular Biology*, 45:257– 285.
- Haworth, M.; Elliott-Kingston, C.; McElwain, J.C. Co-Ordination of Physiological and Morphological Responses of Stomata to Elevated [CO₂] in Vascular Plants. *Oecologia* 2013, 171, 71–82.
- Hibi, N. *et al.* (1994) 'Gene expression in tobacco low-nicotine mutants', *Plant Cell*, 6(5), pp. 723–735. doi: 10.2307/3869875.
- Huang Q-l, Zhang M-m, Li C-s, Li B-y, Zhuo S-l, Yang Y-s, Chen Y-d, Zhong A-n, Liu H-y, Lai W-f, Huang Z-b, Cao M-h, Yuan Z-s and Zhang G-f (2025) Response mechanism of water status and photosynthetic characteristics of *Cotoneaster multiflorus* under drought stress and rehydrated conditions. *Front. Plant Sci*. 15:1457955. doi: 10.3389/fpls.2024.1457955
- Hussain, H. A. *et al.* (2018) 'Chilling and drought stresses in crop plants: Implications, cross talk, and potential management opportunities', *Frontiers in Plant Science*, 9(April), pp. 1–21. doi: 10.3389/fpls.2018.00393.
- Iguz, M.C., M. Aycan, E.Oguz, I. Poyraz, M. Yildiz. 2022. Drought stress tolerance in plants: interplay of molecular, biochemical and physiological responses in important development stages. *Physiologia*, 2: 180–197.

- Jaleel, C. A. *et al.* (2009) 'Drought stress in plants: A review on morphological characteristics and pigments composition', *International Journal of Agriculture and Biology*, 11(1), pp. 100–105.
- Jamil, A. (2016) 'Mekanisme Respon Tanaman Padi terhadap Cekaman Kekeringan dan Kultivar Toleran', *Iptek Tanaman Pangan*, 11(1).
- Kajikawa, M., Shoji, T., Kato, A., & Hashimoto, T. (2011). Vacuole-localized berberine bridge enzyme-like proteins are required for a late step of nicotine biosynthesis in tobacco. *Plant Physiology*, 155(4), 2010–2022.
- Katoh A., T. Shoji, T. Hashimoto. 2007. Molecular cloning of N-methylputrescine oxidase from tobacco. *Plant and Cell Physiology*, 48:550–554.
- Khalvandi, M. *et al.* (2021) 'Salicylic acid alleviated the effect of drought stress on photosynthetic characteristics and leaf protein pattern in winter wheat', *Heliyon*, 7(1), p. e05908. doi: 10.1016/j.heliyon.2021.e05908.
- Kishore, K. (2014). Monograph of tobacco (*Nicotiana tabacum*). *Indian Journal of Drugs*, 2(1), 5–23.
- Kumar, S. *et al.* (2018) 'Plant responses to drought stress: Physiological, biochemical and molecular basis', *Biotic and Abiotic Stress Tolerance in Plants*, (August), pp. 1–25. doi: 10.1007/978-981-10-9029-51.
- Kusumawati, D. A. *et al.* (2022) 'Pengaruh Pemberian Kompos Limbah Batang Tembakau Terhadap Pertumbuhan dan Produksi Tanaman Tembakau (*Nicotiana tabacum* L.) Kasturi', *Agropross : National Conference Proceedings of Agriculture*, pp. 65–78. doi: 10.25047/agropross.2022.275.
- Kutchan, T. M. (1995) 'Nicotine alkaloid biosynthesis', *The Plant Cell*, 7(July), pp. 1059–1070.
- Larionov, A., Krause, A. and Miller, W. R. (2005) 'A standard curve based method for relative real time PCR data processing', *BMC Bioinformatics*, 6. doi: 10.1186/1471-2105-6-62.
- Lewis, R. S. *et al.* (2015) 'Transgenic and mutation-based suppression of a *berberine bridge enzyme-like* (BBL) gene family reduces alkaloid content in field-grown tobacco', *PLoS ONE*, 10(2). doi: 10.1371/journal.pone.0117273.
- Levitt, J. (1981). *Responses of plants to environmental stresses*. Volume II: Water, radiation, salt, and other stresses (2nd ed., 607 pp.). Academic Press.
- Liu, D. *et al.* (2012) 'Validation of Reference Genes for Gene Expression Studies in Virus-Infected *Nicotiana benthamiana* Using Quantitative Real-Time PCR', *PLoS ONE*,

7(9). doi: 10.1371/journal.pone.0046451.

- Liu, M. *et al.* (2024) ‘Tobacco production under global climate change: combined effects of heat and drought stress and coping strategies’, *Frontiers in Plant Science*, 15(November), pp. 1–10. doi: 10.3389/fpls.2024.1489993.
- Livak, K. J. and Schmittgen, T. D. (2001) ‘Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method’, *Methods*, 25(4), pp. 402–408. doi: 10.1006/meth.2001.1262.
- Lorenz, T. C. (2012) ‘*Polymerase Chain Reaction : Basic Protocol Plus Troubleshooting and Optimization Strategies*’, 63(May), pp. 1–15. doi: 10.3791/3998.
- Luo, X. *et al.* (2025) ‘Physiological and transcriptomic analysis of the effect of overexpression of the NTPIP2;4 gene on drought tolerance in tobacco’, *BMC plant biology*, 25(1), p. 63. doi: 10.1186/s12870-024-05896-7.
- Ma, D. M. *et al.* (2019) ‘Untargeted Metabolomics of *Nicotiana tabacum* Grown in United States and India Characterizes the Association of Plant Metabolomes With Natural Climate and Geography’, *Frontiers in Plant Science*, 10(October). doi: 10.3389/fpls.2019.01370.
- Medrano, Marisa L. Wong, J. F. (2005) ‘Real-time PCR for mRNA quantitation’, *BioTechniques*, 39(1), pp. 1–11. doi: 10.1109/TCOM.1979.1094474.
- Mudhor, M. A. *et al.* (2022) ‘Pengaruh Cekaman Kekeringan Terhadap Pertumbuhan dan Produksi Tanaman Padi Hitam Kultivar Jeliteng’, *Agrikultura*, 33(3), p. 247. doi: 10.24198/agrikultura.v33i3.40361.
- Mukarram, M., Choudhary, S., Kurjak, D., Petek, A., and Khan, M. M. A. (2021). Drought: Sensing, signalling, effects and tolerance in higher plants. *Physiologia Plantarum* 172, 1291–1300. doi: 10.1111/ppl.13423
- Müller, M., & Munné-Bosch, S. (2015). Ethylene response factors: A key regulatory hub in hormone and stress signaling. *Plant Physiology*, 169(1), 32–41. <https://doi.org/10.1104/pp.15.00677>.
- Murphy, L. D., Herzog, C. E., Rudick, J. B., Fojo, A. T., & Bates, S. E. (1990). Use of the polymerase chain reaction in the quantitation of *mdr-1* gene expression. *Biochemistry*, 29(43), 10351–10356.
- Nakano T, Suzuki K, Fujimura T, Shinshi H (2006) Genome-wide analysis of the ERF gene family in Arabidopsis and rice. *Plant Physiol* 140: 411–432
- Nezhadahmadi, A.; Prodhan, Z.H.; Faruq, G. Drought Tolerance in Wheat. *Sci. World J.* 2013, 2013, 610721.

- Nicot, N. *et al.* (2005) 'Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress', *Journal of Experimental Botany*, 56(421), pp. 2907–2914. doi: 10.1093/jxb/eri285.
- Ortiz, N. *et al.* (2015) 'Contribution of arbuscular mycorrhizal fungi and/or bacteria to enhancing plant drought tolerance under natural soil conditions: Effectiveness of autochthonous or allochthonous strains', *Journal of Plant Physiology*, 174, pp. 87–96. doi: 10.1016/j.jplph.2014.08.019.
- Pamungkas, S.S.T.; Suwanto; Suprayogi; Farid, N. Drought Stress: Responses and Mechanism in Plants. *Rev. Agric. Sci.* 2022, 10, 168–185
- Pangisian, J., Sangi, M. S. and Kumaunang, M. (2022) 'Analisis Senyawa Metabolit Sekunder dan Uji Aktivitas Antioksidan serta Antibakteri Biji Buah Pangi (*Pangium edule* Reinw)', *Jurnal LPPM Bidang Sains dan Teknologi*, 7(1), pp. 11–19.
- Purnomo, R., Santoso, M. and Heddy, S. (2013) 'Terhadap Pertumbuhan Dan Hasil Tanaman Mentimun (*Cucumis Sativus* L .) The Effect Of Various Dosages Of Organic And Inorganic Fertilizers On Plant Growth And Yield Of Cucumber (*Cucumis sativus* L .)', *Jurnal Produksi Tanaman*, 1(3), pp. 93–100.
- Reshi, Z. A. *et al.* (2023) 'From Nature to Lab: A Review of Secondary Metabolite Biosynthetic Pathways, Environmental Influences, and In Vitro Approaches', *Metabolites*, 13(8). doi: 10.3390/metabo13080895.
- Rochman, F., Suwarso, S. And A.S. Murdiyati, A. S. M. (2020) 'Galur Harapan Tembakau Temanggung Produksi Tinggi dan Tahan Penyakit Lincat', *Jurnal Penelitian Tanaman Industri*, 13(2), p. 57. doi: 10.21082/jlitri.v13n2.2007.57-63.
- Rutledge, R. G. and Stewart, D. (2008) 'Critical evaluation of methods used to determine amplification efficiency refutes the exponential character of real-time PCR', *BMC Molecular Biology*, 9(1), pp. 1–12. doi: 10.1186/1471-2199-9-96.
- Schmidt, G. W. and Delaney, S. K. (2010) 'Stable internal reference genes for normalization of real-time RT-PCR in tobacco (*Nicotiana tabacum*) during development and abiotic stress', *Molecular Genetics and Genomics*, 283(3), pp. 233–241. doi: 10.1007/s00438-010-0511-1.
- Shewmaker, C. K. *et al.* (1990) 'Nucleotide sequence of an *EF-1 α* genomic clone from tomato', *Nucleic Acids Research*, 18(14), p. 4276. doi: 10.1093/nar/18.14.4276.
- Shi, Q., Li, C. and Zhang, F. (2006) 'Nicotine synthesis in *Nicotiana tabacum* L. induced by mechanical wounding is regulated by auxin', *Journal of Experimental Botany*, 57(11), pp. 2899–2907. doi: 10.1093/jxb/erl051.

- Shitan, N. *et al.* (2014) 'Involvement of the leaf-specific multidrug and toxic compound extrusion (mate) transporter Nt-JAT2 in vacuolar sequestration of nicotine in *Nicotiana tabacum*', *PLoS ONE*, 9(9), pp. 1–10. doi: 10.1371/journal.pone.0108789.
- Shoji, T. *et al.* (2009) 'Multidrug and toxic compound extrusion-type transporters implicated in vacuolar sequestration of nicotine in tobacco roots', *Plant Physiology*, 149(2), pp. 708–718. doi: 10.1104/pp.108.132811.
- Shoji, T., Hashimoto, T. and Saito, K. (2024) 'Genetic regulation and manipulation of nicotine biosynthesis in tobacco: strategies to eliminate addictive alkaloids', *Journal of Experimental Botany*, 75(6), pp. 1741–1753. doi: 10.1093/jxb/erad341.
- Shoji, T., Kajikawa, M. and Hashimoto, T. (2010) 'Clustered transcription factor genes regulate nicotine biosynthesis in tobacco', *Plant Cell*, 22(10), pp. 3390–3409. doi: 10.1105/tpc.110.078543.
- Shoji T. and T. Hashimoto. 2011. Tobacco MYC2 regulates jasmonate-inducible nicotine biosynthesis genes directly and by way of the *NIC2*-locus ERF genes. *Plant Cell Physiology*, 52:1117–1130
- Shoji, T. (2020). Nicotine biosynthesis, transport, and regulation in tobacco: Insights into the evolution of a metabolic pathway. In N. V. Ivanov, N. Sierra, & M. C. Peitsch (Eds.), *The tobacco plant genome* (pp. 147–156). *Springer*. https://doi.org/10.1007/978-3-030-18768-0_12
- Subantoro, R. (2014) 'Pengaruh cekaman kekeringan terhadap respon fisiologis perkecambahan benih kacang tanah', *Mediagro*, 10(2), pp. 32–44.
- Sulistiyarningsih, E. *et al.* (2024) 'Identification and validation of reference genes for gene expression study using Quantitative Real-Time PCR in shallot (*Allium cepa* L. *Aggregatum* group) based on bulb development phase', *Biodiversitas*, 25(6), pp. 2645–2651. doi: 10.13057/biodiv/d250634.
- Sulistiyono, E., Suwarno, & Lubis, I. (2012). Karakterisasi morfologi dan fisiologi untuk mendapatkan marka morfologi dan fisiologi padi sawah tahan kekeringan (-30 kPa) dan produktivitas tinggi (> 8 ton/ha). *Jurnal Ilmu Pertanian Indonesia*, 17(2).
- Sulistiyowati, S., Nurchayati, Y. and Setiari, N. (2021) 'Pertumbuhan dan Produksi Tomat (*Lycopersicon esculentum* Mill.) Kultivar Servo pada Frekuensi Penyiraman yang Berbeda', *Buletin Anatomi dan Fisiologi*, 6(1), pp. 26–34. doi: 10.14710/baf.6.1.2021.26-34.
- Sun, B. *et al.* (2013) 'Genetic variation in alkaloid accumulation in leaves of *Nicotiana*', *Journal of Zhejiang University: Science B*, 14(12), pp. 1100–1109. doi:

10.1631/jzus.B1300130.

- Suwarso. 2014. Source of Tobacco Genetic: Management Status and Application (in Source of Genetic Agriculture: 45 Food, Estate, and Horticulture Crops Eds Sumarno, Hasnam, I. Mustika, and Bahagiawati). *IAARD Press*. 165-185.
- Syahid Muttaqin, A. *et al.* (2019) 'The impact of climate variability on tobacco productivity over Temanggung Regency, Indonesia', *E3S Web of Conferences*, 76, pp. 3–7. doi: 10.1051/e3sconf/20197604003.
- Talapatra, S., Wagner, J. D. O. and Thompson, C. B. (2002) 'Elongation factor-1 alpha is a selective regulator of growth factor withdrawal and ER stress-induced apoptosis', *Cell Death and Differentiation*, 9(8), pp. 856–861. doi: 10.1038/sj.cdd.4401078.
- Twaij, B. M. and Hasan, M. N. (2022) 'Bioactive Secondary Metabolites from Plant Sources: Types, Synthesis, and Their Therapeutic Uses', *International Journal of Plant Biology*, 13(1), pp. 4–14. doi: 10.3390/ijpb13010003.
- Ursin, V. M. *et al.* (1991) 'Developmental Analysis of Elongation Factor- ICW Expression in Transgenic Tobacco', 3(June), pp. 583–591.
- Wang, J. *et al.* (2021) 'Exploitation of drought tolerance-related genes for crop improvement', *International Journal of Molecular Sciences*, 22(19). doi: 10.3390/ijms221910265.
- Wang, Q. *et al.* (2024) 'Genotypic Differences in Morphological, Physiological and Agronomic Traits in Wheat (*Triticum aestivum* L.) in Response to Drought', *Plants*, 13(2). doi: 10.3390/plants13020307.
- Willems, E., Leyns, L. and Vandesompele, J. (2008) 'Standardization of real-time PCR gene expression data from independent biological replicates', *Analytical Biochemistry*, 379(1), pp. 127–129. doi: 10.1016/j.ab.2008.04.036.
- Xu, S. *et al.* (2017) 'Wild tobacco genomes reveal the evolution of nicotine biosynthesis', *Proceedings of the National Academy of Sciences of the United States of America*, 114(23), pp. 6133–6138. doi: 10.1073/pnas.1700073114.
- Yang, X. *et al.* (2019) 'The different influences of drought stress at the flowering stage on rice physiological traits, grain yield, and quality', *Scientific Reports*, 9(1), pp. 1–12. doi: 10.1038/s41598-019-40161-0.
- Yang, X. *et al.* (2021) 'Response mechanism of plants to drought stress', *Horticulturae*, 7(3). doi: 10.3390/horticulturae7030050.
- Yuan, L. (2020) 'Clustered ERF Transcription Factors: Not All Created Equal', *Plant and Cell Physiology*, 61(6), pp. 1025–1027. doi: 10.1093/pcp/pcaa067.

- Yusra Faperta. (2013, Maret 4). *Deskripsi kultivar unggul tembakau (Nicotiana tabacum)* [Dokumen kerja]. Scribd. <https://www.scribd.com/document/129783262/Deskripsi-Kultivar-Unggul-Tembakau>
- Zaman, Q. uz *et al.* (2024) ‘Combined application of biochar and peatmoss for mitigation of drought stress in tobacco’, *BMC Plant Biology*, 24(1). doi: 10.1186/s12870-024-05576-6.
- Zenkner, F. F., Margis-Pinheiro, M. and Cagliari, A. (2019) ‘Nicotine Biosynthesis in *Nicotiana*: A Metabolic Overview’, *Tobacco Science*, 56(1), pp. 1–9. doi: 10.3381/18-063.
- Zhang, H., Zhao, Y. and Zhu, J. K. (2020) ‘Thriving under Stress: How Plants Balance Growth and the Stress Response’, *Developmental Cell*, 55(5), pp. 529–543. doi: 10.1016/j.devcel.2020.10.012.