

## DAFTAR PUSTAKA

- Arora, D., & Sharma, P. (2010). Ligninolytic fungal laccases and their biotechnological applications. *Applied Biochemistry and Biotechnology*, 160(6), 1760–1788. <https://doi.org/10.1007/s12010-009-8676-y>
- Ashenafi, M., & Busse, M. (1991). Growth of lactic acid bacteria during fermentation of tef (*Eragrostis tef*), the raw material of injera. *World Journal of Microbiology and Biotechnology*, 7, 607–611.
- Bai, Y., Wang, J., Zheng, H., Liu, J., & Chen, T. (2021). Dynamics of fungal communities during soybean fermentation for tempeh production. *Food Research International*, 140(1), 109870. <https://doi.org/10.1016/j.foodres.2020.109870>
- Barus, T., Simanjuntak, P., & Lubis, M. (2020). Komposisi mikrobiota tempe lokal berdasarkan analisis metagenomik. *Jurnal Mikrobiologi Indonesia*, 15(2), 72–81. <https://doi.org/10.5454/mi.15.2.3>
- Chen, B., Du, H., & Xu, Y. (2014). Changes in the microbial community during the spontaneous fermentation of traditional Chinese soy sauce. *International Journal of Food Microbiology*, 182–183(1), 132–138. <https://doi.org/10.1016/j.ijfoodmicro.2014.01.008>
- Feng, Y., Duan, C., & Xu, Y. (2007). Dynamics of microbial community during Chinese soy sauce fermentation. *International Journal of Food Microbiology*, 120, 87–95.
- Hachmeister, K. A., & Fung, D. Y. C. (1993). Tempeh: A mold-modified indigenous fermented food made from soybeans and/or cereal grains. *Critical Reviews in Microbiology*, 19(3), 137–188.
- Huang, C., Wu, H., Liu, Z., Liu, G., Di, J., & Lin, X. (2012). Effect of organic acids on the growth and lipid accumulation of oleaginous yeast *Trichosporon fermentans*. *Biotechnology for Biofuels*, 5(4), Article 4. <https://doi.org/10.1186/1754-6834-5-4>
- Hou, Q., Bai, X., Li, W., Gao, X., Zhang, F., Sun, Z., & Zhang, H. (2018). Design of primers for evaluation of lactic acid bacteria populations in complex biological samples. *Frontiers in Microbiology*, 9(1), 2045. <https://doi.org/10.3389/fmicb.2018.02045>
- Illegghems, K., De Vuyst, L., Papalexandratou, Z., & Weckx, S. (2015). Importance

of the microbial consortia in cocoa bean fermentation. *Applied and Environmental Microbiology*, 81(19), 5996–6006.

<https://doi.org/10.1128/AEM.00574-15>

Kohlmeier, S., Smits, T. H. M., Ford, R. M., Keel, C., Harms, H., & Wick, L. Y. (2005). Taking the fungal highway: Mobilization of pollutant-degrading bacteria by fungi. *Environmental Science & Technology*, 39(12), 4640–4646.

<https://doi.org/10.1021/es048153z>

Kusumoto, K., Yamamoto, M., & Nagai, K. (2022). Temporal dynamics of microbial diversity during fermentation of soy-based food. *Microbiology Spectrum*, 10(4), e01178-22. <https://doi.org/10.1128/spectrum.01178-22>

Leech, J., Cabrera-Rubio, R., Walsh, A. M., Finnegan, L., Ahern, G., & Cotter, P. D. (2020). Fermented foods, microbiota, and mental health: Ancient practice meets nutritional psychiatry. *Journal of Functional Foods*, 75(1), 104294.

<https://doi.org/10.1016/j.jff.2020.104294>

Li, X., Chen, Q., & Zhang, L. (2022). Geographically associated fungus–bacterium interactions contribute to flavor formation during complex spontaneous fermentation. *Microbiology Spectrum*, 10(2), e01844-22.

<https://doi.org/10.1128/spectrum.01844-22>

Magdalena, S., Hogaputri, J. E., Yulandi, A., & Yogiara, Y. (2022). The addition of lactic acid bacteria in the soybean soaking process of tempeh. *Food Research*, 6(3), 27–33. [https://doi.org/10.26656/fr.2017.6\(3\).304](https://doi.org/10.26656/fr.2017.6(3).304)

Mulyowidarso, R. K., Fleet, G. H., Buckle, K. A., & Wills, R. B. H. (1989). Microbiology of tempe fermentation. *International Journal of Food Microbiology*, 8, 213–222.

Nguyen, T. T. P., Loiseau, G., & Guyot, J. P. (2007). Antibacterial activity of fermented extracts from legumes and seeds. *International Journal of Food Microbiology*, 113(3), 304–312.

<https://doi.org/10.1016/j.ijfoodmicro.2006.07.014>

Nurdini, C., Nurhidayat, N., & Harahap, D. (2015). Studi pengaruh suhu dan aktivitas air pada pertumbuhan kapang tempe. *Jurnal Teknologi dan Industri Pangan*, 26(2), 101–108.

Parra-Flores, J., Rodríguez, A., García-Herrera, N., Bruna, F., & Aguayo-Reyes, A. (2018). *Cronobacter sakazakii* and microbiological parameters in powdered infant formula in Chile. *Frontiers in Microbiology*, 9(1), 1127.

<https://doi.org/10.3389/fmicb.2018.01127>

- Parnell, J. J., Berka, R., Young, H. A., Sturino, J. M., Kang, Y., & Barnhart, D. M. (2016). From the lab to the farm: An industrial perspective of plant beneficial microorganisms. *Frontiers in Plant Science*, 7(1), 1110. <https://doi.org/10.3389/fpls.2016.01110>
- Rizzatti, G., Lopetuso, L. R., Gibiino, G., Binda, C., & Gasbarrini, A. (2017). Proteobacteria: A common factor in human diseases. *BioMed Research International*, 2017(1), 9351507. <https://doi.org/10.1155/2017/9351507>
- Romulo, R., & Sashihara, T. (2021). Expression profile of *Rhizopus oligosporus* protease genes during tempeh fermentation. *Journal of Applied Microbiology*, 130(3), 887–898. <https://doi.org/10.1111/jam.14810>
- Smith, J., Clark, B., & Jones, M. (2020). Temperature-induced variation in microbial community and metabolome shifts in controlled fermentation. *mSystems*, 5(5), e00555-20. <https://doi.org/10.1128/mSystems.00555-20>
- Takahashi, S., Tomita, J., Nishioka, K., Hisada, T., & Nishijima, M. (2014). Development of a prokaryotic universal primer for simultaneous analysis of bacteria and archaea using next-generation sequencing. *PLOS ONE*, 9(8), e105592. <https://doi.org/10.1371/journal.pone.0105592>
- Vymazal, O., Villas-Bôas, S., & Duarte, Q. (2020). Succession of bacterial and fungal communities during fermentation of medicinal plants. *Fermentation*, 8(8), 383. <https://doi.org/10.3390/fermentation8080383>
- Wei, S., Morrison, M., & Yu, Z. (2020). Comparative microbial community analysis using metadata standards (MIxS) improves reproducibility. *Microbiome*, 8(1), 52. <https://doi.org/10.1186/s40168-020-00827-y>
- Wiesel, I., Reuter, G., & Kandler, O. (1997). Interactions between lactic acid bacteria and molds in traditional fermented foods. *Food Microbiology*, 14, 91–98.
- Wolfe, B. E., Button, J. E., Santarelli, M., & Dutton, R. J. (2014). Cheese rind communities provide tractable systems for in situ and in vitro studies of microbial diversity. *Cell*, 158(2), 422–433. <https://doi.org/10.1016/j.cell.2014.05.04>
- Xu, Y., Zhang, Z., & Wu, Q. (2021). Changes in microbial community structure during digitally managed fermentation of medium-temperature Daqu. *Food Science and Technology*, 42(1), e02021.

- Yatsu, F., Sawano, Y., & Ito, K. (2015). Morphological and physiological responses of *Rhizopus* spp. under competitive conditions in solid-state fermentation. *Journal of Food Biochemistry*, 39(5), 591–599. <https://doi.org/10.1111/jfbc.12149>
- Yuan, C., Zhu, X., & Li, J. (2020). SparCC-based correlation network analysis reveals microbial interactions during tempeh fermentation. *Frontiers in Microbiology*, 11(1), 1925. <https://doi.org/10.3389/fmicb.2020.01925>
- Zhang, T., Fang, Y., & Li, S. (2020). Microbial community and its association with physicochemical factors during compost bedding for dairy cows. *Frontiers in Microbiology*, 11(1), 254. <https://doi.org/10.3389/fmicb.2020.00254>
- Zhou, X., Riffel, A., & Brandelli, A. (2005). Production of an extracellular keratinase from *Chryseobacterium* sp. growing on raw feathers. *Applied Microbiology and Biotechnology*, 70(4), 620–626. <https://doi.org/10.1007/s00253-005-0116-7>