

DAFTAR PUSTAKA

- Aaqil, M., Peng, C., Kamal, A., Nawaz, T., Zhang, F., & Gong, J. (2023). Tea Harvesting and Processing Techniques and Its Effect on Phytochemical Profile and Final Quality of Black Tea: A Review. *Foods*, 12(24), 4467. <https://doi.org/10.3390/foods12244467>
- Ahmed, S., Orians, C. M., Griffin, T. S., Buckley, S., Unachukwu, U., Stratton, A. E., Stepp, J. R. Robbat AJr., Cash, S., Kennelly, E. J. (2014). Effects of water availability and pest pressures on tea (*Camellia sinensis*) growth and functional quality. *AoB Plants*, 6, plt054.
- Akuli, A., Pal, A., Bej, G., Dey, T., Ghosh, A., Tudu, B., Bhattacharyya, N. and Bandyopadhyay, R., (2016). A machine vision system for estimation of theaflavins and thearubigins in orthodox black tea. *International Journal on Smart Sensing and Intelligent Systems*, 9(2), 709.
- Apriyani, T. W., Supriyadi, S., & Gunadi, R. (2022). Improvement of antioxidant activity and sensory quality of Pagilaran's Tea clones treated by tannase. *AgriTECH*, 42(3), 218-230.
- Badan Pusat Statistik. (2024). *Statistik Teh Indonesia 2023*. Jakarta: Badan Pusat Statistik.
- Badan Standardisasi Nasional. (2016). *SNI 01-1902-1990 Teh Hitam*. Jakarta: Badan Standardisasi Nasional.

- Badan Standardisasi Nasional. (2016). *SNI 1902:2016 Teh Hitam*. Jakarta: Badan Standardisasi Nasional.
- Benti, T., Debela, A., Bekele, Y., & Suleman, S. (2023). Effect of seasonal variation on yield and leaf quality of tea clone (*Camellia sinensis* (L.) O. Kuntze) in South West Ethiopia. *Heliyon*, 9(3), e14051. <https://doi.org/10.1016/j.heliyon.2023.e14051>
- Borse, B. B., & Rao, J. M. L. (2012). Novel Bio-Chemical Profiling of Indian Black Teas with Reference to Quality Parameters. *Journal of Bioequivalence and Bioavailability*, 14, 1-16.
- Chaturvedula, V. S. P., & Prakash, I. (2011). The aroma, taste, color and bioactive constituents of tea. *Journal of Medicinal Plants Research*, 5(11), 2110-2124.
- De Costa, W. A., Mohotti, A. J., & Wijeratne, M. A. (2007). Ecophysiology of tea. *Brazilian Journal of Plant Physiology*, 19, 299-332.
- Deb, S., & Jolvis Pou, K. R. (2016). A review of withering in the processing of black tea. *Journal of Biosystems Engineering*, 41(4), 365-372.
- Dewi, S. (2019). Mutu Petik Teh (*Camellia sinensis* (L.) O. Kuntze) di Kebun Bedakah, Wonosobo, Jawa Tengah. *Buletin Agrohorti*, 7(3), 337-342.
- Effendi, M.S., Syakir, M., Yusron, M. (2017). *Budidaya dan Pascapanen Teh*. Jakarta: Pusat Penelitian dan Pengembangan Perkebunan.
- Hu, D., Chen, W., Miao, A., Qiao, X., Xia, H., & Ma, C. (2023). Analysis of volatile emission of tea (*Camellia sinensis*) shoots in response to temperature-

dependent postharvest treatments. *European Journal of Agronomy*, 146, 126821.

International Agency For Research on Cancer. (1991). *IARC Monograph on The Evaluation of Carcinogenic Risks to Humans: Volume 51 Coffee, Tea, Mate, Methylxanthines, and Methylgloxal*. France: IARC Lyon.

Jolvis Pou, K. R. (2016). Fermentation: The key step in the processing of black tea. *Journal of Biosystems Engineering*, 41(2), 85-92.

Kaur, A., Suri, R., Rana, K., Thakur, V., Dimri, P., & Mittal, N. (2019). Antioxidant levels in Indian rose, hibiscus, chrysanthemum and marigold tea and their comparison with black and green tea. *International Research Journal Of Pharmacy*, 10(10), 52–55. <https://doi.org/10.7897/2230-8407.1010298>

Kraujalytė, V., Pelvan, E., & Alasalvar, C. (2016). Volatile compounds and sensory characteristics of various instant teas produced from black tea. *Food Chemistry*, 194, 864-872.

Li, X., Luo, L., He, Y., & Xu, N. (2013). Determination of dry matter content of tea by near and middle infrared spectroscopy coupled with wavelet-based data mining algorithms. *Computers and Electronics in Agriculture*, 98, 46-53.

Li, Y., Pan, W., Feng, Z., Gu, F., Chen, J., Wang, F., Wang, J., Zhang, M., Li, C., McClements, D.J. and Yin, J., (2025). Deciphering composition-structure-taste relationship of black tea-infusion via assessments of nanoparticles by centrifugal treatment. *LWT*, 222, 117601.

- Lung J.K.S., dan Destiani D.P. (2017). Uji antioksidan vitamin A C E dengan metode DPPH. *Suplemen*, 15(1): 55-62.
- Luo, Q., Luo, L., Zhao, J., Wang, Y., & Luo, H. (2024). Biological potential and mechanisms of Tea's bioactive compounds: An Updated review. *Journal of advanced research*, 65, 345-363.
- Mei, S., Yu, Z., Chen, J., Zheng, P., Sun, B., Guo, J., & Liu, S. (2022). The Physiology of Postharvest Tea (*Camellia sinensis*) Leaves, According to Metabolic Phenotypes and Gene Expression Analysis. *Molecules (Basel, Switzerland)*, 27(5), 1708. <https://doi.org/10.3390/molecules27051708>.
- Nadiah, N. I., & Uthumporn, U. (2015). Determination of phenolic and antioxidant properties in tea and spent tea under various extraction method and determination of catechins, caffeine and gallic acid by HPLC. *International Journal on Advanced Science Engineering and Information Technology*, 5(3):158.
- Ntezimana, B., Li, Y., He, C., Yu, X., Zhou, J., Chen, Y., ... & Ni, D. (2021). Different withering times affect sensory qualities, chemical components, and nutritional characteristics of black tea. *Foods*, 10(11), 2627.
- Ölmez, H., & Yilmaz, A. (2010). Changes in chemical constituents and polyphenol oxidase activity of tea leaves with shoot maturity and cold storage. *Journal of Food Processing and Preservation*, 34, 653-665.

- Owuor, P. O., & Orchard, J. E. (1992). Effects of storage time in a two-stage withering process on the quality of seedling black tea. *Food chemistry*, 45(1), 45-49.
- Owuor, P., & Orchard, J. (1989). Changes in the biochemical constituents of green leaf and black tea to withering: a review. *Tea*, 10(1), 53-59.
- Parveen, A., Qin, C.-Y., Zhou, F., Lai, G., Long, P., Zhu, M., Ke, J., & Zhang, L. (2023). The Chemistry, Sensory Properties and Health Benefits of Aroma Compounds of Black Tea Produced by *Camellia sinensis* and *Camellia assamica*. *Horticulturae*, 9(12), 1253. <https://doi.org/10.3390/horticulturae9121253>
- Pribadi, I., Da'I, M., dan Utami, W. (2008). Uji aktivitas antiradikal Buah *Psidium guajava* L. dengan Metode DPPH (1,1-Difenil-2-Pikril Hidrazil) serta Penetapan Kadar Fenolik dan Flavonoid. *Jurnal Farmasi Indonesia Pharmaccon*. Vol. 9(2), h. 52-56
- Rohdiana, D. (2015). Teh, Karakteristik, Proses, dan Komponen Fungsionalnya. Pusat Penelitian Teh dan Kina. *Food Review Indonesia*, 10(8): 20-27.
- Setyamidjaja, D.J. (2000). *Budidaya dan Pengolahan Teh Pascapanen*. Kanisius: Yogyakarta.
- Sivapalan, K. (1971). Respiratory metabolism in detached shoot tips of the tea plant. *Tea Q.*, 42, 123-130.
- Sudaryat, Y., Kusmiyati, M., Pelangi, C. R., & Rustamsyah, A. (2015). Aktivitas antioksidan seduhan sepuluh jenis mutu teh hitam (*Camellia sinensis* (L.)

- O. Kuntze*) Indonesia antioxidant activity of ten grades of Indonesia black tea. *Jurnal Penelitian Teh dan Kina*, 18(2), 95–100.
- Syahbudin, A., Widyastuti, A., Masruri, N. W., & Meinata, A. (2019). Morphological Classification of Tea Clones (*Camellia sinensis*, *Theaceae*) at the Mount Lawu Forest, East Java, Indonesia. *IOP Conf. Ser.: Earth Environ. Sci.* 394 012014
- Tang, G. -Y., Meng, X., Gan, R. -Y., Zhao, C. -N., Liu, Q., Feng, Y. -B., Li, S., Wei, X. -L., Atanasov, A. G., Corke, H., & Li, H. -B. (2019). Health Functions and Related Molecular Mechanisms of Tea Components: An Update Review. *International Journal of Molecular Sciences*, 20(24), 6196. <https://doi.org/10.3390/ijms20246196>
- Tomlins, K. I., & Mashingaidze, A. (1997). Influence of withering, including leaf handling, on the manufacturing and quality of black teas—a review. *Food chemistry*, 60(4), 573-580.
- Tong, T., Liu, Y. J., Kang, J., Zhang, C. M., & Kang, S. G. (2019). Antioxidant activity and main chemical components of a novel fermented tea. *Molecules*, 24(16), 2917.
- Tran, P. N., Nguyen, Y. T., Nguyen, M. Q., Nguyen, T. D., & Tran, H. T. (2024). Fermented or not fermented: a comparative study of the biological activities and microbiomes of a non-fermented green tea and a five-year-old fermented tea prepared from young leaves of Shan tea. *International Journal of Food Science and Technology*, 59(7), 4693-4700.

- Turkmen, N., Sari, F., & Velioglu, Y. S. (2009). Factors affecting polyphenol content and composition of fresh and processed tea leaves. *Akademik Gida*, 7(6), 29-40.
- Ullah, M. R. (1986). A rapid procedure for estimating theaflavins and thearubigins of black tea. *Two and a Bud*, 33, 46–48.
- Wicaksono, I.N.A. & Martono, B. (2020). PENAMPILAN FENOTIPIK, KERAGAMAN, DAN HERITABILITAS SEMBILAN GENOTIPE TEH [*Camellia sinensis* (L.) O. Kuntze]. *Journal of Industrial and Beverages Crops* 7(2). 53-60.
- Winarsi, H., (2007). *Antioksidan Alami & Radikal Bebas*. Yogyakarta: Kanisius.
- Yang, J., & Liu, R. H. (2013). The phenolic profiles and antioxidant activity in different types of tea. *International Journal of Food Science and Technology*, 48(1), 163-171.
- Yao, L., Liu, X., Jiang, Y., Caffin, N., D’Arcy, B., Singanusong, R., Datta, N. and Xu, Y. (2006). Compositional analysis of teas from Australian supermarkets. *Food Chemistry*, 94(1), 115-122.
- Yan, Z., Zhong, Y., Duan, Y., Chen, Q., & Li, F. (2020). Antioxidant mechanism of tea polyphenols and its impact on health benefits. *Animal nutrition (Zhongguo xu mu shou yi xue hui)*, 6(2), 115–123. <https://doi.org/10.1016/j.aninu.2020.01.001>
- Zhang, S., Li, Q., Wen, S., Sun, L., Chen, R., Zhang, Z., Cao, J., Lai, Z., Li, Z., Lai, X. and Wu, P. (2023). Metabolomics reveals the effects of different storage

times on the acidity quality and metabolites of large-leaf black tea. *Food*

Chemistry, 426, 136601.