

## DAFTAR PUSTAKA

- Abdel-Aziz, S.A., Taha, M., Balabel, M., Hegazy, O., Salah, E., Soltan, M., *et al.* (2023). Prognostic factors determining the surgical interference of biliary atresia in Egyptian infants: single-center experience. *Egypt. Liver J.* 13.
- Anindita, A.N., Setyoboedi, B., Arief, S., & Prihaningtyas, R.A. (2024). Accuracy of 2-phase abdominal ultrasound for diagnosing biliary atresia. *Bali Med. J.* 13 : 54–58.
- Arua, H.N., Eze, C.U., & Arua, C.L. (2021). Correlation of Sonographic Measurements of the Gallbladder and somatometric variables in Primary School Age Pupils in Nigeria. *Adv. Res. Gastroenterol. Hepatol.* 17.
- Brahee, D.D., & Lampl, B.S. (2022). Neonatal diagnosis of biliary atresia: a practical review and update. *Pediatr. Radiol.* 52 : 685–692.
- Caruso, M., Cuocolo, R., Di Dato, F., Mollica, C., Vallone, G., Romeo, V., *et al.* (2020). Ultrasound, shear-wave elastography, and magnetic resonance imaging in native liver survivor patients with biliary atresia after Kasai portoenterostomy: correlation with medical outcome after treatment. *Acta radiol.* 61 : 1300–1308.
- Caruso, M., Di Dato, F., Mollica, C., Vallone, G., Romeo, V., Liuzzi, R., *et al.* (2021). Imaging prediction with ultrasound and MRI of long-term medical outcome in native liver survivor patients with biliary atresia after kasai portoenterostomy: a pilot study. *Abdom. Radiol.* 46 : 2595–2603.
- Cavallo, L., Kovar, E.M., Aqul, A., McLoughlin, L., Mittal, N.K., Rodriguez-Baez, N., *et al.* (2022). The Epidemiology of Biliary Atresia: Exploring the Role of Developmental Factors on Birth Prevalence. *J. Pediatr.* 246 : 89-94.e2.
- Cho, Y.S., Jeong, W.K., & Kim, Y. (2020). Ultrasonographic morphological diagnosis of chronic liver disease: 2-dimensional shear wave elastography as an add-on test. *Ultrasonography* 39 : 272–280.
- Choochuen, P., Kritsaneepaiboon, S., Charoonratana, V., & Sangkhathat, S. (2019a). Is “gallbladder length-to-width ratio” useful in diagnosing biliary atresia? *J. Pediatr. Surg.* 54 : 1946–1952.
- Choochuen, P., Kritsaneepaiboon, S., Charoonratana, V., & Sangkhathat, S. (2019b). Is “gallbladder length-to-width ratio” useful in diagnosing biliary atresia? *J. Pediatr. Surg.* 54 : 1946–1952.
- Chung, P.H.Y., Wong, K.K.Y., & Tam, P.K.H. (2015). Predictors for failure after Kasai operation. *J. Pediatr. Surg.* 50 : 293–296.
- Dahlan, S. (2010). Besar Sampel dan Cara Pengambilan Sampel. Jakarta : Salemba Medika.
- Degtyareva, A., Isaeva, M., Tumanova, E., Filippova, E., Sugak, A., Razumovsky, A., *et al.* (2024). Combined Predictors of Long-Term Outcomes of Kasai Surgery in Infants with Biliary Atresia 27 : 224–235.
- Di Serafino, M., Severino, R., Gioioso, M., Rossi, E., Vezzali, N., Pelliccia, P., *et al.* (2020). Paediatric liver ultrasound: a pictorial essay. *J. Ultrasound* 23 : 87–103.
- Elmanar, R., Violetta, L., & Setiawati, R. (2024). Correlation between Length-to-

width Ratio of Gallbladder and Gamma- glutamyl Transferase Value in Biliary Atresia 12 : 256–262.

Gad, E.H., Kamel, Y., Salem, T.A.H., Ali, M.A.H., & Sallam, A.N. (2021). Short- and long-term outcomes after Kasai operation for type III biliary atresia: Twenty years of experience in a single tertiary Egyptian center-A retrospective cohort study. *Ann. Med. Surg.* 62 : 302–314.

Ghazy, R.M., Adawy, N.M., Khedr, M.A., & Tahoun, M.M. (2018). Biliary atresia recent insight. *Egypt. Pediatr. Assoc. Gaz.* 66 : 1–8.

Harumatsu, T., Muraji, T., Masuya, R., Tsuruno, Y., Iwamoto, Y., Ogata, M., *et al.* (2024). Inflammation patterns in early post-operative cholangitis predict long-term outcomes in biliary atresia: a potential role of non-suppurative cholangitis. *Pediatr. Surg. Int.* 40 : 1–6.

J Richard Landis, & Koch., G.G. (1977). The Measurement of Observer Agreement for Categorical Data. *Biometrics* 33 : 159–174.

Jeon, T.Y., Yoo, S.Y., Kim, J.H., Eo, H., & Lee, S.K. (2013). Serial ultrasound findings associated with early liver transplantation after Kasai portoenterostomy in biliary atresia. *Clin. Radiol.* 68 : 588–594.

Kanegawa, K., Akasaka, Y., Kitamura, E., Nishiyama, S., Muraji, T., Nishijima, E., *et al.* (2003). Sonographic Diagnosis of Biliary Atresia in Pediatric Patients Using the “Triangular Cord” Sign Versus Gallbladder Length and Contraction. *Am. J. Roentgenol.* 181 : 1387–1390.

Kelly, D.A., & Davenport, M. (2007). Current management of biliary atresia. *Arch. Dis. Child.* 92 : 1132–1135.

Kim, H., Yoo, S.Y., Kim, J.H., Kim, M.J., Lee, S., & Jeon, T.Y. (2024). Preoperative imaging findings to predict 2-year native liver survival after the Kasai procedure in patients with biliary atresia. *Eur. Radiol.* 34 : 1493–1501.

Kim, M.Y., Cho, M.Y., Baik, S.K., Park, H.J., Jeon, H.K., Im, C.K., *et al.* (2011). Histological subclassification of cirrhosis using the Laennec fibrosis scoring system correlates with clinical stage and grade of portal hypertension. *J. Hepatol.* 55 : 1004–1009.

Kim, S.S., Kim, M.J., Lee, M.J., Yoon, C.S., Han, S.J., & Koh, H. (2014). Ultrasonographic findings of type IIIa biliary atresia. *Ultrasonography* 33 : 267–274.

Koo, T.K., & Li, M.Y. (2016). A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J. Chiropr. Med.* 15 : 155–163.

Kuebler, J.F., Madadi-Sanjani, O., Pfister, E.D., Baumann, U., Fortmann, D., Leonhardt, J., *et al.* (2021). Adjuvant therapy with budesonide post-Kasai reduces the need for liver transplantation in biliary atresia. *J. Clin. Med.* 10.

Kumar, R., Lal, B.B., Sood, V., Khanna, R., Kumar, S., Bharathy, K.G.S., *et al.* (2019). Predictors of Successful Kasai Portoenterostomy and Survival with Native Liver at 2 Years in Infants with Biliary Atresia. *J. Clin. Exp. Hepatol.* 9 : 453–459.

Lee, S.M., Cheon, J.E., Choi, Y.H., Kim, W.S., Cho, H.H., Kim, I.O., *et al.* (2015). Ultrasonographic diagnosis of biliary atresia based on a decision-making tree model. *Korean J. Radiol.* 16 : 1364–1372.

- Lefkowitz, J.H. (1998). Biliary Atresia. *Mayo Clin. Proc.* 73 : 90–95.
- Liu, J.Q., Chen, W.J., Zhou, M.J., Li, W.F., Tang, J., & Zhou, Q.C. (2021). A Nomogram Predicting the Prognosis of Children With Biliary Atresia After Hepatopertoenterostomy. *Front. Pediatr.* 9 : 1–10.
- Malik, A., Thanekar, U., Mourya, R., & Shivakumar, P. (2020). Recent developments in etiology and disease modeling of biliary atresia: a narrative review. *Dig. Med. Res.* 3.
- Mohtasib, R.S., Alshamiri, K., Jobeir, A., Ambu-Saidi, F.M., Masawi, A., Alabdulaziz, L., *et al.* (2021). Sonographic measurements for spleen size in healthy Saudi children and correlation with body parameters. *Ann. Saudi Med.* 41 : 14–23.
- Nugrahaeni, A.S. (2023). Faktor-Faktor Prediktor Kegagalan Prosedur Kasai Pada Pasien Anak Atresia Bilier. Universitas Gadjah Mada.
- Oswari, H., & Lidvihurin, A.S. (2019). Pemberian Steroid untuk Meningkatkan Bilirubin Clearance pada Pasien dengan Atresia Bilier Pasca Prosedur Kasai. *Sari Pediatr.* 20 : 249.
- Rahayatri, T.H., Siahaan, K.B.P., Ulina, R.P., Stephanie, M., & Oswari, H. (2019). Evaluasi Pasien Praoperasi Transplantasi Hati Anak di RSUPN dr. Cipto Mangunkusumo Evaluation of Preoperative Pediatric Liver Transplantation in dr. Cipto Mangunkusumo Hospital. *Desember 7* : 53–60.
- Ramakrishna, S.H., Nayak, S.P., Rao, S., D’Cruz, A.L.J., Garg, M., & Ramachandran, K. (2023). Kasai Portoenterostomy at a Slightly Delayed Age and Native Liver Survival in Children With Biliary Atresia: Single Center Experience. *Indian Pediatr.* 60 : 659–662.
- Romadhon, B.D., Supriyadi, B., Artsini, E., Wibawa, T., Choridah, L., & Utomo, B.P. (2025). Perbandingan Morfologi Vesica Felea, Contractility Index Dan Ketebalan Triangular Cord Sign Dalam Pemeriksaan Ultrasonografi Abdomen Dua Fase Terhadap Hasil Kolangiografi Intraoperatif Untuk Diagnosis Atresia Bilier. Gadjah Mada University.
- Sastroasmoro, S., & Ismael, S. (2011). Dasar-dasar metodologi penelitian klinis. Jakarta : Sagung Seto.
- Saxena, A.K., Mittal, V., & Sodhi, K.S. (2012). Triangular Cord Sign in Biliary Atresia: Does It Have Prognostic and Medical Significance? *Radiology* 263 : 621.
- Schreiber, R.A., Harpavat, S., Hulscher, J.B.F., & Wildhaber, B.E. (2022). Biliary Atresia in 2021: Epidemiology, Screening and Public Policy. *J. Clin. Med.* 11.
- Schuh, R. (2023). Coarsened hepatic echotexture [WWW Document]. *Radiopaedia*.
- Serinet, M.O., Wildhaber, B.E., Broué, P., Lachaux, A., Sarles, J., Jacquemin, E., *et al.* (2009). Impact of age at Kasai operation on its results in late childhood and adolescence: A rational basis for biliary atresia screening. *Pediatrics* 123 : 1280–1286.
- Sharma, B., & John, S. (2024). Hepatic Cirrhosis [WWW Document]. *StatPearls*. URL <https://www.ncbi.nlm.nih.gov/books/NBK482419/> (accessed 11.26.24).
- Shen, W.J., Chen, G., Wang, M., & Zheng, S. (2019). Liver fibrosis in biliary atresia. *World J. Pediatr.* 15 : 117–123.

- Sherwani, P., Bolia, R., Kaushik, A., Kumar, S., Kishore, S., & Chauhan, U. (2021). An Unusual Cause of Cholestasis in an Infant: Biliary Atresia Type IIB. *Indian J. Radiol. Imaging* 31 : 689–692.
- Siddiqui, A.I., & Ahmad, T. (2023). Biliary Atresia. [Updated 2023 Jun 26] [WWW Document]. *StatPearls*. URL <https://www.ncbi.nlm.nih.gov/books/NBK537262/%0A> (accessed 11.25.24).
- Soekersi, H., Santiana, L., & Fatmawaty, F. (2020). Comparative Study Gallbladder Contractility Index Using Ultrasound in Patients with and without Liver Cirrhosis Perbedaan Indeks Kontraktilitas Kandung Empedu Menggunakan Ultrasonografi pada Penderita Sirosis Hati dan tanpa Sirosis Hati 8 : 7–12.
- Sugiono (2007). Statistik untuk penelitian, 11th ed. Bandung : CV Alfabeta.
- Sutowo, D.W., Gunadi, & Makhmudi, A. (2019). Faktor Prognosis Kolangitis, Hipertensi Porta, dan Varises Esofagus untuk Kesintasan Pasien Atresia Bilier pasca Operasi Kasai di RSUP Dr. Sardjito. *Repos. UGM*.
- Vehmas, T., Kaukiainen, A., Luoma, K., Lohman, M., Nurminen, M., & Taskinen, H. (2004). Liver echogenicity: Measurement or visual grading? *Comput. Med. Imaging Graph.* 28 : 289–293.
- Vij, M., & Rela, M. (2020). Biliary atresia: Pathology, etiology and pathogenesis. *Futur. Sci. OA* 6.
- Wang, G., Zhang, N., Zhang, X., Zhou, W., Xie, X., & Zhou, L. (2021). Ultrasound characteristics combined with gamma-glutamyl transpeptidase for diagnosis of biliary atresia in infants less than 30 days. *Pediatr. Surg. Int.* 37 : 1175–1182.
- Wong, K.K.Y., Chung, P.H.Y., Chan, I.H.Y., Lan, L.C.L., & Tam, P.K.H. (2010). Performing kasai portoenterostomy beyond 60 days of life is not necessarily associated with a worse outcome. *J. Pediatr. Gastroenterol. Nutr.* 51 : 631–634.
- Yan, H., Liu, J., Jin, S., Du, L., Wang, Q., & Luo, Y. (2023). A novel prediction tool based on shear wave elastography, gallbladder ultrasound, and serum biomarkers for the early diagnosis of biliary atresia in infants younger than 60 days old. *Quant. Imaging Med. Surg.* 13 : 259–270.
- Yang, C.Z., Zhou, Y., Ke, M., Gao, R.Y., Ye, S.R., Diao, M., *et al.* (2022). Effects of postoperative adjuvant steroid therapy on the outcomes of biliary atresia: A systematic review and updated meta-analysis. *Front. Pharmacol.* 13 : 1–13.
- Yen, Y.H., Kuo, F.Y., Chen, C.H., Hu, T.H., Lu, S.N., Wang, J.H., *et al.* (2019). Ultrasound is highly specific in diagnosing compensated cirrhosis in chronic hepatitis C patients in real world clinical practice. *Medicine (Baltimore)*. 98 : e16270.
- Yoon, H., Ihn, K., Kim, J., Lim, H.J., Park, S., Han, S.J., *et al.* (2023). Pre-and Immediate Post-Kasai Portoenterostomy Shear Wave Elastography for Predicting Hepatic Fibrosis and Native Liver Outcomes in Patients With Biliary Atresia. *Korean J. Radiol.* 24 : 465–475.
- Zagory, J.A., Nguyen, M. V, & Wang, K.S. (2015). Recent advances in the pathogenesis and management of biliary atresia. *Curr. Opin. Pediatr.* 27 : 389–394.
- Zhang, K., Tang, Y., Zheng, Z., Tang, C., Zhu, D., Du, Q., *et al.* (2023). Value of

gallbladder length-to-width ratio for diagnosis of biliary atresia by correlation with age. *Updates Surg.* 75 : 915–920.

Zhou, W., & Zhou, L. (2022). Ultrasound for the diagnosis of biliary atresia: From conventional ultrasound to artificial intelligence. *Diagnostics* 12.