

DAFTAR PUSTAKA

- Abdurahman, A., Bandiara, R., & Supriyadi, R. (2019). MON-055 The growing burden of end stage renal disease in Indonesia: ten years of the Indonesian renal registry reports. *Kidney International Reports*, 4(7), S327.
- Administration, U. S. F. and D. (2004). FDA announces qualified health claims for omega-3 fatty acids. *FDA News Release P*, 4, 89.
- Afsar, B., & Elsurer Afsar, R. (2012). HbA1c is related with uremic pruritus in diabetic and nondiabetic hemodialysis patients. *Renal Failure*, 34(10), 1264–1269.
- Agarwal, R. (2016). Defining end-stage renal disease in clinical trials: a framework for adjudication. *Nephrology Dialysis Transplantation*, 31(6), 864–867.
- Ahuja, Rajeev B., et al. "A comparative analysis of cetirizine, gabapentin and their combination in the relief of post-burn pruritus." *Burns* 37.2 (2011): 203-207.
- Akhyani, M., Ganji, M. R., Samadi, N., Khamesan, B., & Daneshpazhooh, M. (2005). Pruritus in hemodialysis patients. *BMC Dermatology*, 5. <https://doi.org/10.1186/1471-5945-5-7>
- Aresi, G., Rayner, H. C., Hassan, L., Burton, J. O., Mitra, S., Sanders, C., & van der Veer, S. N. (2019). Reasons for underreporting of uremic pruritus in people with chronic kidney disease: a qualitative study. *Journal of Pain and Symptom Management*, 58(4), 578–586.
- Association, W. M. (2013). World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *Jama*, 310(20), 2191–2194.
- Balitbang Kemenkes, R. I. (2018). Laporan Nasional Riset Kesehatan Dasar Kementrian Kesehatan RI. *Jakarta: Kementrian Kesehatan Republik Indonesia*.
- Bautista, D. M., Wilson, S. R., & Hoon, M. A. (2014). Why we scratch an itch: the molecules, cells and circuits of itch. *Nature Neuroscience*, 17(2), 175–182.
- Begum, R., Belury, M. A., Burgess, J. R., & Peck, L. W. (2004). Supplementation with n-3 and n-6 polyunsaturated fatty acids: Effects on lipoxygenase activity and clinical symptoms of pruritus in hemodialysis patients. *Journal of Renal Nutrition*, 14(4), 233–241.
- Bergasa, N. V., Alling, D. W., Talbot, T. L., Swain, M. G., Yurdaydin, C., Turner, M. L., Schmitt, J. M., Walker, E. C., & Jones, E. A. (1995). Effects of naloxone infusions in patients with the pruritus of cholestasis: a double-blind, randomized, controlled trial. *Annals of Internal Medicine*, 123(3), 161–167.
- Bigliardi, P. L., Tobin, D. J., Gaveriaux-Ruff, C., & Bigliardi-Qi, M. (2009). Opioids and the skin—where do we stand? *Experimental Dermatology*, 18(5), 424–430.

- Blachley, J. D., Blankenship, D. M., Menter, A., Parker III, T. F., & Knochel, J. P. (1985). Uremic pruritus: skin divalent ion content and response to ultraviolet phototherapy. *American Journal of Kidney Diseases*, 5(5), 237–241.
- Borbe, J. B. C., Elvambuena, B. F., Eugenio, F. A. D. R., & Tan, R. J. M. (2024). Effect of Omega-3 Fatty Acid Supplementation in the Treatment of Uremic Pruritus among Dialytic Chronic Kidney Disease Patients: A Meta-analysis. *Acta Medica Philippina*, 58(8), 125.
- Chang, C.-H., Tseng, P.-T., Chen, N.-Y., Lin, P.-C., Lin, P.-Y., Chang, J. P.-C., Kuo, F.-Y., Lin, J., Wu, M.-C., & Su, K.-P. (2018). Safety and tolerability of prescription omega-3 fatty acids: A systematic review and meta-analysis of randomized controlled trials. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 129, 1–12.
- Chen, H.-Y., Chiu, Y.-L., Hsu, S.-P., Pai, M.-F., Lai, C.-F., Yang, J.-Y., Peng, Y.-S., Tsai, T.-J., & Wu, K.-D. (2010). Elevated C-reactive protein level in hemodialysis patients with moderate/severe uremic pruritus: a potential mediator of high overall mortality. *QJM: An International Journal of Medicine*, 103(11), 837–846.
- CL, H. (1968). Disappearance of " UREMIC" itching after subtortal parathyroid-ectomy. *N Engl J Med*, 279, 695–697.
- Combs, S. A., Teixeira, J. P., & Germain, M. J. (2015). Pruritus in kidney disease. *Seminars in Nephrology*, 35(4), 383–391.
- Cowan, A., Kehner, G. B., & Inan, S. (2015). Targeting itch with ligands selective for κ opioid receptors. *Pharmacology of Itch*, 291–314.
- Cunnane, S. C. (2003). Problems with essential fatty acids: time for a new paradigm? *Progress in Lipid Research*, 42(6), 544–568.
- Dugas-Breit, S., Schöpf, P., Dugas, M., Schiffli, H., Ruëff, F., & Przybilla, B. (2005). Baseline serum levels of mast cell tryptase are raised in hemodialysis patients and associated with severity of pruritus: Basale Mastzelltryptase-Serumspiegel sind bei Haemodialyse-Patienten erhöht und mit der Pruritus-Intensität assoziiert. *JDDG: Journal Der Deutschen Dermatologischen Gesellschaft*, 3(5), 343–347.
- Elman, S., Hynan, L. S., Gabriel, V., & Mayo, M. J. (2010). The 5-D itch scale: a new measure of pruritus. *British Journal of Dermatology*, 162(3), 587–593.
- Fallahzadeh, M. K., Roozbeh, J., Geramizadeh, B., & Namazi, M. R. (2011). Interleukin-2 serum levels are elevated in patients with uremic pruritus: a novel finding with practical implications. *Nephrology Dialysis Transplantation*, 26(10), 3338–3344.
- Fleming, G. M. (2011). Renal replacement therapy review: past, present and future. *Organogenesis*, 7(1), 2–12.
- Forouhari, A., Moghtaderi, M., Raeisi, S., Shahidi, S., Parin Hedayati, Z., Zaboliyan, J., Ani, S., Moeinzadeh, F., & Mortazavi, M. (2022). Pruritus-reducing effects of omega-3 fatty acids in hemodialysis patients: A cross-over randomized clinical trial. *Hemodialysis International*, 26(3), 408–414.

- Francos, G. C., Kauh, Y. C., Gittlen, S. D., Schulman, E. S., Besarab, A., Goyal, S., & Burke Jr, J. F. (1991). Elevated plasma histamine in chronic uremia effects of ketotifen on pruritus. *International Journal of Dermatology*, 30(12), 884–889.
- Garcia, G. G., Harden, P., & Chapman, J. (2012). The global role of kidney transplantation. *The Lancet*, 379(9820), e36–e38.
- Ghanei, E., Zeinali, J., Borghei, M., & Homayouni, M. (2012). Efficacy of omega-3 fatty acids supplementation in treatment of uremic pruritus in hemodialysis patients: a double-blind randomized controlled trial. *Iranian Red Crescent Medical Journal*, 14(9), 515.
- Gilchrest, B. A., Stern, R. S., Steinman, T. I., Brown, R. S., Arndt, K. A., & Anderson, W. W. (1982). Clinical features of pruritus among patients undergoing maintenance hemodialysis. *Archives of Dermatology*, 118(3), 154–156.
- Grochulska, K., Ofenloch, R., Mettang, T., & Weisshaar, E. (2019). Mortality of Haemodialysis Patients With and Without Chronic Itch: A Follow-up Study of the German Epidemiological Hemodialysis Itch Study (GEHIS). *Acta Dermato-Venereologica*, 99(4).
- Gunal, A. I., Ozalp, G., Yoldas, T. K., Gunal, S. Y., Kirciman, E., & Celiker, H. (2004). Gabapentin therapy for pruritus in haemodialysis patients: a randomized, placebo-controlled, double-blind trial. *Nephrology Dialysis Transplantation*, 19(12), 3137–3139.
- Heydarbaki, Mansoureh, *et al.* An investigation into the effects of omega-3 on uremic pruritus in hemodialysis patients: a randomized crossover clinical trial. *Saudi Journal of Kidney Diseases and Transplantation*, 2021, 32.6: 1562-1569.
- Hibbeln, J. R., Nieminen, L. R. G., Blasbalg, T. L., Riggs, J. A., & Lands, W. E. M. (2006). Healthy intakes of n-3 and n-6 fatty acids: estimations considering worldwide diversity. *The American Journal of Clinical Nutrition*, 83(6), 1483S-1493S.
- Hiroshige, K., Kabashima, N., Takasugi, M., & Kuroiwa, A. (1995). Optimal dialysis improves uremic pruritus. *American Journal of Kidney Diseases*, 25(3), 413–419.
- Hu, X., Sang, Y., Yang, M., Chen, X., & Tang, W. (2018). Prevalence of chronic kidney disease-associated pruritus among adult dialysis patients: a meta-analysis of cross-sectional studies. *Medicine*, 97(21), e10633.
- Indonesia, P. N. (2018). 11th Report of Indonesian renal registry 2018. *Jakarta: Perhimpunan Nefrologi Indonesia*, 1–46.
- Inker, L. A., Astor, B. C., Fox, C. H., Isakova, T., Lash, J. P., Peralta, C. A., Tamura, M. K., & Feldman, H. I. (2014). KDOQI US commentary on the 2012 KDIGO clinical practice guideline for the evaluation and management of CKD. *American Journal of Kidney Diseases*, 63(5), 713–735.
- Jeong, S. K., Park, H. J., Park, B. D., & Kim, I.-H. (2010). Effectiveness of topical chia seed oil on pruritus of end-stage renal disease (ESRD) patients and healthy volunteers. *Annals of Dermatology*, 22(2), 143–148.

- Johansson, O., Hilliges, M., & Ståhle-Bäckdahl, M. (1989). Intraepidermal neuron-specific enolase (NSE)-immunoreactive nerve fibres: evidence for sprouting in uremic patients on maintenance hemodialysis. *Neuroscience Letters*, 99(3), 281–286.
- Kim, D., & Pollock, C. (2021). Epidemiology and burden of chronic kidney disease-associated pruritus. *Clinical Kidney Journal*, 14(Supplement_3), i1–i7.
- Kimata, N., Fuller, D. S., Saito, A., Akizawa, T., Fukuhara, S., Pisoni, R. L., Robinson, B. M., & Akiba, T. (2014). Pruritus in hemodialysis patients: Results from the Japanese Dialysis Outcomes and Practice Patterns Study (JDOPPS). *Hemodialysis International*, 18(3), 657–667.
- Kimmel, M., Alscher, D. M., Dunst, R., Braun, N., Machleidt, C., Kiefer, T., Stülten, C., van der Kuip, H., Pauli-Magnus, C., & Raub, U. (2006). The role of micro-inflammation in the pathogenesis of uraemic pruritus in haemodialysis patients. *Nephrology Dialysis Transplantation*, 21(3), 749–755.
- Klein, P. A., & Clark, R. A. (1999). An evidence-based review of the efficacy of antihistamines in relieving pruritus in atopic dermatitis. *Archives of dermatology*, 135(12), 1522–1525.
- Ko, M.-J., Peng, Y.-S., Chen, H.-Y., Hsu, S.-P., Pai, M.-F., Yang, J.-Y., Wen, S.-Y., Jee, S.-H., Wu, H.-Y., & Chiu, H.-C. (2014). Interleukin-31 is associated with uremic pruritus in patients receiving hemodialysis. *Journal of the American Academy of Dermatology*, 71(6), 1151–1159.
- Ko, M.-J., Wu, H.-Y., Chen, H.-Y., Chiu, Y.-L., Hsu, S.-P., Pai, M.-F., Lai, C.-F., Lu, H.-M., Huang, S.-C., & Yang, S.-Y. (2013). Uremic pruritus, dialysis adequacy, and metabolic profiles in hemodialysis patients: a prospective 5-year cohort study. *PloS One*, 8(8), e71404.
- Krajewski, P. K., Krajewska, M., & Szepietowski, J. C. (2020). Pruritus in renal transplant recipients: Current state of knowledge. *Advances in Clinical & Experimental Medicine*, 29(6).
- Krajewski, P. K., Olczyk, P., Krajewska, M., Krajewski, W., & Szepietowski, J. C. (2021). Clinical characteristics of itch in renal transplant recipients. *Frontiers in Medicine*, 7, 615334.
- Lahiji, A. P., Mortazavi, M., Tirani, S. A., Moeinzadeh, F., Bidaki, E. Z., Naini, A. E., Faghihi, G., Toghyani, A., & Farajzadegan, Z. (2018). Omega-3 supplementation improves pruritus in continuous ambulatory peritoneal dialysis patients: a crossover randomized pilot clinical trial. *Journal of Research in Pharmacy Practice*, 7(4), 195–199.
- Lai, J.-W., Chen, H.-C., Chou, C.-Y., Yen, H.-R., Li, T.-C., Sun, M.-F., Chang, H.-H., Huang, C.-C., Tsai, F.-J., & Tschen, J. (2017). Transformation of 5-D itch scale and numerical rating scale in chronic hemodialysis patients. *BMC Nephrology*, 18, 1–5.
- Lau, T., Leung, S., & Lau, W. (2016). Gabapentin for uremic pruritus in hemodialysis patients: a qualitative systematic review. *Canadian Journal of Kidney Health and Disease*, 3, 107.

- Levey, A. S., Coresh, J., Bolton, K., Culleton, B., Harvey, K. S., Ikizler, T. A., Johnson, C. A., Kausz, A., Kimmel, P. L., & Kusek, J. (2002). K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *American Journal of Kidney Diseases*, 39(2 SUPPL. 1), i-ii+.
- Levin, A., Stevens, P. E., Bilous, R. W., Coresh, J., De Francisco, A. L. M., De Jong, P. E., Griffith, K. E., Hemmelgarn, B. R., Iseki, K., & Lamb, E. J. (2013). Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney International Supplements*, 3(1), 1–150.
- Li, J., Guo, Q., Lin, J., Yi, C., Yang, X., & Yu, X. (2015). Prevalence and associated factors of uraemic pruritus in continuous ambulatory peritoneal dialysis patients. *Internal Medicine*, 54(22), 2827–2833.
- Lu, P.-H., Yao, X.-F., Lin, Y.-S., Tzeng, I.-S., Lu, P.-H., & Kuo, K.-L. (2022). Omega-3 fatty acids for uremic pruritus: A meta-analysis of randomized controlled trials. *Tzu Chi Medical Journal*, 34(4), 394–401.
- Lydia, A., Widiana, I. G. R., Bandiara, R., Afiatin, Ali, Z., Nugroho, P., & Hidayati, E. L. (2021). Nephrology in Indonesia. *Nephrology Worldwide*, 299–312.
- Majeski, C. J., Johnson, J. A., Davison, S. N., & Lauzon, G. J. (2007). Itch Severity Scale: a self-report instrument for the measurement of pruritus severity. *British Journal of Dermatology*, 156(4), 667–673.
- Masi, C. M., & Cohen, E. P. (1992). Dialysis efficacy and itching in renal failure. *Nephron*, 62(3), 257–261.
- Mathur, V. S., Lindberg, J., Germain, M., Block, G., Tumlin, J., Smith, M., Grewal, M., McGuire, D., & Investigators, I. N. R. (2010). A longitudinal study of uremic pruritus in hemodialysis patients. *Clinical Journal of the American Society of Nephrology*, 5(8), 1410–1419.
- Mettang, T., & Kremer, A. E. (2015). Uremic pruritus. *Kidney International*, 87(4), 685–691.
- Min, J.-W., Kim, S.-H., Kim, Y. O., Jin, D. C., Song, H. C., Choi, E. J., Kim, Y.-L., Kim, Y.-S., Kang, S.-W., & Kim, N.-H. (2016). Comparison of uremic pruritus between patients undergoing hemodialysis and peritoneal dialysis. *Kidney Research and Clinical Practice*, 35(2), 107–113.
- Moeller, S., Gioberge, S., & Brown, G. (2002). ESRD patients in 2001: global overview of patients, treatment modalities and development trends. *Nephrology Dialysis Transplantation*, 17(12), 2071–2076.
- Murtagh, F. E. M., Addington-Hall, J. M., Edmonds, P. M., Donohoe, P., Carey, I., Jenkins, K., & Higginson, I. J. (2007). Symptoms in advanced renal disease: a cross-sectional survey of symptom prevalence in stage 5 chronic kidney disease managed without dialysis. *Journal of Palliative Medicine*, 10(6), 1266–1276.

- Narita, I., Alchi, B., Omori, K., Sato, F., Ajiro, J., Saga, D., Kondo, D., Skatsume, M., Maruyama, S., & Kazama, J. J. (2006). Etiology and prognostic significance of severe uremic pruritus in chronic hemodialysis patients. *Kidney International*, 69(9), 1626–1632.
- Osakwe, N., & Hashmi, M. F. (2024). Uremic pruritus evaluation and treatment. In *StatPearls [Internet]*. StatPearls Publishing.
- Oweis, A. O., Firas, A.-Q., Bodoor, K., Heis, L., Alfaqih, M. A., Almomani, R., Obeidat, M. A., & Alshelleh, S. A. (2021). Elevated interleukin 31 serum levels in hemodialysis patients are associated with uremic pruritus. *Cytokine*, 138, 155369.
- Ozen, N., Cinar, F. I., Askin, D., & Mut, D. (2018). Uremic pruritus and associated factors in hemodialysis patients: A multi-center study. *Kidney Research and Clinical Practice*, 37(2), 138.
- Panahi, Y., Dashti-Khavidaki, S., Farnood, F., Noshad, H., Lotfi, M., & Gharekhani, A. (2016). Therapeutic effects of omega-3 fatty acids on chronic kidney disease-associated pruritus: a literature review. *Advanced Pharmaceutical Bulletin*, 6(4), 509.
- Pisoni, R. L., Wikström, B., Elder, S. J., Akizawa, T., Asano, Y., Keen, M. L., Saran, R., Mendelssohn, D. C., Young, E. W., & Port, F. K. (2006). Pruritus in haemodialysis patients: International results from the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Nephrology Dialysis Transplantation*, 21(12), 3495–3505.
- Plewig, N., Ofenloch, R., Mettang, T., & Weisshaar, E. (2019). The course of chronic itch in hemodialysis patients: results of a 4-year follow-up study of GEHIS (German Epidemiological Hemodialysis Itch Study). *Journal of the European Academy of Dermatology and Venereology*, 33(7), 1429–1435.
- Rafieipoor, Alireza, *et al.* "Effectiveness of omega-3 fatty acid supplementation for pruritus in patients undergoing hemodialysis." *Frontiers in Nutrition* 11 (2024): 1328469.
- Ramakrishnan, K., Bond, T. C., Claxton, A., Sood, V. C., Kootsikis, M., Agnese, W., & Sibbel, S. (2013). Clinical characteristics and outcomes of end-stage renal disease patients with self-reported pruritus symptoms. *International Journal of Nephrology and Renovascular Disease*, 1–12.
- Rayner, H. C., Larkina, M., Wang, M., Graham-Brown, M., van der Veer, S. N., Ecker, T., Hasegawa, T., Kleophas, W., Bieber, B. A., & Tentori, F. (2017). International comparisons of prevalence, awareness, and treatment of pruritus in people on hemodialysis. *Clinical Journal of the American Society of Nephrology*, 12(12), 2000–2007.
- Rehman, I. U., Chan, K. G., Munib, S., Lee, L. H., & Khan, T. M. (2019). The association between CKD-associated pruritus and quality of life in patients undergoing hemodialysis in Pakistan: A STROBE complaint cross-sectional study. *Medicine*, 98(36), e16812.

- Rehman, I. U., Chohan, T. A., Bukhsh, A., & Khan, T. M. (2019). Impact of pruritus on sleep quality of hemodialysis patients: a systematic review and meta-analysis. *Medicina*, 55(10), 699.
- Rehman, I. U., Lai, P. S. M., Lim, S. K., Lee, L. H., & Khan, T. M. (2019). Sleep disturbance among Malaysian patients with end-stage renal disease with pruritus. *BMC Nephrology*, 20, 1–8.
- Rylance, P. B., Gordge, M. P., Saynor, R., Parsons, V., & Weston, M. J. (1986). Fish oil modifies lipids and reduces platelet aggregability in haemodialysis patients. *Nephron*, 43(3), 196–202.
- Scherer, J. S., Combs, S. A., & Brennan, F. (2017). Sleep disorders, restless legs syndrome, and uremic pruritus: diagnosis and treatment of common symptoms in dialysis patients. *American Journal of Kidney Diseases*, 69(1), 117–128.
- Shayanpour, S., Beladi Mousavi, S. S., Rizi, P. L., & Cheraghian, B. (2019). The effect of the omega-3 supplement on uremic pruritus in hemodialysis patients; a double-blind randomized controlled clinical trial. *Journal of Nephropathology*, 8(2).
- Solak, B., Acikgoz, S. B., Sipahi, S., & Erdem, T. (2016). Epidemiology and determinants of pruritus in pre-dialysis chronic kidney disease patients. *International Urology and Nephrology*, 48, 585–591.
- Ständer, S., Augustin, M., Reich, A., Blome, C., Ebata, T., Phan, N. Q., & Szepietowski, J. C. (2013). Pruritus assessment in clinical trials: consensus recommendations from the International Forum for the Study of Itch (IFSI) Special Interest Group Scoring Itch in Clinical Trials. *Acta Dermato-Venereologica*, 93(5).
- Sukul, N., Karaboyas, A., Csomor, P. A., Schaufler, T., Wen, W., Menzaghi, F., Rayner, H. C., Hasegawa, T., Al Salmi, I., & Al-Ghamdi, S. M. G. (2021). Self-reported pruritus and clinical, dialysis-related, and patient-reported outcomes in hemodialysis patients. *Kidney Medicine*, 3(1), 42–53.
- Sukul, N., Speyer, E., Tu, C., Bieber, B. A., Li, Y., Lopes, A. A., Asahi, K., Mariani, L., Laville, M., & Rayner, H. C. (2019). Pruritus and patient reported outcomes in non-dialysis CKD. *Clinical Journal of the American Society of Nephrology*, 14(5), 673–681.
- Sukul, N., Zhao, J., Pisoni, R. L., Walpen, S., Schaufler, T., Asgari, E., Guebre-Egziabher, F., Zho, L., Al-Ghonaim, M. A., & Nitta, K. (2023). Pruritus in hemodialysis patients: longitudinal associations with clinical and patient-reported outcomes. *American Journal of Kidney Diseases*, 82(6), 666–676.
- SuSeł, J., Batycka-Baran, A., Reich, A., & SzEPIETOWSKI, J. C. (2014). Uraemic pruritus markedly affects the quality of life and depressive symptoms in haemodialysis patients with end-stage renal disease. *Acta Dermato-Venereologica*, 94(3).
- Tey, H. L., & Yosipovitch, G. (2011). Targeted treatment of pruritus: a look into the future. *British Journal of Dermatology*, 165(1), 5–17.

- Ting, S.-W., Fan, P.-C., Lin, Y.-S., Lin, M.-S., Lee, C.-C., Kuo, G., & Chang, C.-H. (2020). Uremic pruritus and long-term morbidities in the dialysis population. *PLoS One*, 15(10), e0241088.
- Verduzco, H. A., & Shirazian, S. (2020). CKD-associated pruritus: new insights into diagnosis, pathogenesis, and management. *Kidney International Reports*, 5(9), 1387–1402.
- Wala-Zielińska, K., Świerczyńska-Mróz, K., Krajewski, P. K., Nowicka-Suszko, D., Krajewska, M., & Szepietowski, J. C. (2023). Endogenous opioid imbalance as a potential factor involved in the pathogenesis of chronic kidney disease-associated pruritus in dialysis patients. *Journal of Clinical Medicine*, 12(7), 2474.
- Weiss, M., Mettang, T., Tschulena, U., Passlick-Deetjen, J., & Weisshaar, E. (2015). Prevalence of chronic itch and associated factors in haemodialysis patients: a representative cross-sectional study. *Acta Dermato-Venereologica*, 95(7), 816–821.
- Weiss, M., Mettang, T., Tschulena, U., & Weisshaar, E. (2016). Health-related quality of life in haemodialysis patients suffering from chronic itch: results from GEHIS (German Epidemiology Haemodialysis Itch Study). *Quality of Life Research*, 25, 3097–3106.
- Weisshaar, E., Weiss, M., Passlick-Deetjen, J., Tschulena, U., Maleki, K., & Mettang, T. (2015). Laboratory and dialysis characteristics in hemodialysis patients suffering from chronic itch-results from a representative cross-sectional study. *BMC Nephrology*, 16, 1–8.
- Welter, E. de Q., Frainer, R. H., Maldotti, A., Losekann, A., & Weber, M. B. (2011). Evaluating the association between alterations in mineral metabolism and pruritus in hemodialysis patients. *Anais Brasileiros de Dermatologia*, 86, 31–36.
- Weng, C.-H., Hu, C.-C., Yen, T.-H., Hsu, C.-W., & Huang, W.-H. (2018). Uremic pruritus is associated with two-year cardiovascular mortality in long term hemodialysis patients. *Kidney and Blood Pressure Research*, 43(3), 1000–1009.
- Wu, H.-Y., Huang, J.-W., Tsai, W.-C., Peng, Y.-S., Chen, H.-Y., Yang, J.-Y., Hsu, S.-P., Pai, M.-F., Ko, M.-J., & Hung, K.-Y. (2018). Prognostic importance and determinants of uremic pruritus in patients receiving peritoneal dialysis: a prospective cohort study. *PLoS One*, 13(9), e0203474.
- Wu, H.-Y., Peng, Y.-S., Chen, H.-Y., Tsai, W.-C., Yang, J.-Y., Hsu, S.-P., Pai, M.-F., Lu, H.-M., Chiang, J.-F., & Ko, M.-J. (2016). A comparison of uremic pruritus in patients receiving peritoneal dialysis and hemodialysis. *Medicine*, 95(9), e2935.
- Wulandani, M. P., Dachlan, A. S., & Yusharyahya, S. N. (2018). Validity and Reliability of 5-D Itch Scale in Indonesian Language on Adult and Geriatric Patient at Dr. Cipto Mangunkusumo Hospital. *Advanced Science Letters*, 24(9), 6994–6998.
- Yamamoto, Y., Hayashino, Y., Yamazaki, S., Akiba, T., Akizawa, T., Asano, Y., Saito, A., Kurokawa, K., Miyachi, Y., & Fukuhara, S. (2009). Depressive symptoms predict the

future risk of severe pruritus in haemodialysis patients: Japan Dialysis Outcomes and Practice Patterns Study. *British Journal of Dermatology*, 161(2), 384–389.

Yatzidis, H., Digenis, P., & Fountas, P. (1975). Hypervitaminosis A accompanying advanced chronic renal failure. *British Medical Journal*, 3(5979), 352.

Yeap, C. T., Yo, T. E., Tan, Y. L. C., Liew, A., & Seng, J. J. B. (2021). Complementary and alternative medicine therapies for uremic pruritus—a systematic review of randomized controlled trials. *Complementary Therapies in Medicine*, 56, 102609.

Zakrzewska-Pniewska, B., & Jędras, M. (2001). Is pruritus in chronic uremic patients related to peripheral somatic and autonomic neuropathy?: study by RR interval variation test (RRIV) and by sympathetic skin response (SSR). *Neurophysiologie Clinique/Clinical Neurophysiology*, 31(3), 181–193.

Zhao, J.-H., Zhu, Q.-S., Li, Y.-W., & Wang, L.-L. (2021). Determinants of the intensity of uremic pruritus in patients receiving maintenance hemodialysis: a cross-sectional study. *Plos One*, 16(1), e0245370.