

## DAFTAR PUSTAKA

- Alves, F.C., Sun, J., Qureshi, A.R., Dai, L., Snaedal, S., Bárány, P., Heimbürger, O., Lindholm, B., Stenvinkel, P. (2018) The higher mortality associated with low serum albumin is dependent on systemic inflammation in end-stage kidney disease. *PLoS One*. Jan 3;13(1):e0190410. doi: 10.1371/journal.pone.0190410.
- Barbosa S.M.C., Barros A.J., Wang J., Heymsfield S.B., Pierson R.N. (2005). Bioelectrical impedance analysis: population reference values for phase angle by age and sex. *Am J Clin Nutr*. 2005 Jul;82(1):49-52. doi: 10.1093/ajcn.82.1.49. PMID: 16002799.
- Bek, K., Akman, S., Bilge, H., Topaluglu, R., Caliskan, S., Peru, H., Cengiz, N. (2009) Chronic kidney disease in children in Turkey. *Pediatr Nephrol* 24:797-806.
- Campbell, K.L., Ash, S., Bauer, J.D., Davies, P.S.. (2007) Evaluation of nutrition assessment tools compared with body cell mass for the assessment of malnutrition in chronic kidney disease. *J Ren Nutr*. May;17(3):189-95. doi: 10.1053/j.jrn.2006.12.005.
- CDC. (2008) Anthropometric Reference Data for Children and Adults: United States, 2003-2006. *National Health Statistics Reports*.
- CDC. (2023) Anthropometric Reference Data Set 1: Children 2 to 20 years, Clinical charts with 5th and 95th percentiles. *National Center for Health Statistics*.
- Chen, S., Ma, X., Zhou, X., Wang, Y., Liang, W., Zheng, L., Zang, X., Mei, X., Qi, Y., Jiang, Y., Zhang, S., Li, J., Chen, H., Shi, Y., Hu, Y., Tao, M., Zhuang, S., Liu, N. (2022) An updated clinical prediction model of protein-energy wasting for hemodialysis patients. *Front. Nutr*. 9:933745. doi: 10.3389/fnut.2022.933745.
- Edefonti, A., Picca, M., Damiani, M. F. (2016) Nutritional management of children with chronic kidney disease and on dialysis. *Int J Nephrol Renovasc Dis*, 9: 263–273. doi:10.2147/IJNRD.S97998.
- Fouque, D. A. (2018) proposed nomenclature and diagnostic criteria for protein-energy wasting in acute and chronic kidney disease. *Kidney International*, Volume 73, Issue 4, 391 – 398.
- Flynn, J.T., Mitsnefes, M., Pierce, C., Cole, S.R., Parekh, R.S., Furth, S.L., Warady, B.A. (2014) Chronic Kidney Disease in Children Study Group. Blood pressure in children with chronic kidney disease: a report from the Chronic Kidney Disease in Children study. *Hypertension*. Oct;52(4):631-7. doi: 10.1161/HYPERTENSIONAHA.108.110635.

- Goldstein, S.L., Currier, H., Watters, L., Hempe, J.M., Sheth, R.D., Silverstein, D. (2003) Acute and chronic inflammation in pediatric patients receiving hemodialysis. *J Pediatr.* Nov;143(5):653-7. doi: 10.1067/S0022-3476(03)00534-1.
- González, C.H., Pineda, M.C., Mejía, F.M. (2020) Skeletal muscle mass by bioelectrical impedance analysis and calf circumference for sarcopenia diagnosis. Jul 24, *Journal of Electrical Bioimpedance*, Volume 11.
- Guo, Y., Zhang, M., Ye, T., Wang, Z., Yao, Y. (2023) Application of Bioelectrical Impedance Analysis in Nutritional Management of Patients with Chronic Kidney Disease. *Nutrients.* 15(18):3941. doi.org/10.3390/nu15183941.
- Harambat, J., van Stralen, K.J., Kim, J.J., Tizard, E.J. (2012) Epidemiology of chronic kidney disease in children. *Pediatr Nephrol.* Mar;27(3):363-73. doi: 10.1007/s00467-011-1939-1.
- Helmy, R., Fadel, F.I., Galal, R.E., Mohammed, A.M., Sayed, S. (2025) Erythropoietin resistance among pediatric patients on chronic hemodialysis: A cross-sectional study. *Pediatr Nephrol.* Apr 28. doi: 10.1007/s00467-025-06776-4. Epub ahead of print. PMID: 40293460.
- Hill, N. R., Fatoba, S. T., Oke, J. L., Hirst, J. A., O'Callaghan, C. A., Lasserson, D. S. (2016) Global Prevalence of Chronic Kidney Disease – A Systematic Review and Meta-Analysis. *PLoS ONE* 11(7): e0158765. doi:10.1371/journal.pone.0158765
- Hyun, Y.Y., Lee, K.B., Han, S.H., Kim, Y.H., Kim, Y.S., Lee, S.W., Oh, Y.K., Chae, D.W., Ahn, C. (2017). Nutritional Status in Adults with Predialysis Chronic Kidney Disease: KNOW-CKD Study. *J Korean Med Sci.*:257-263. doi: 10.3346/jkms.2017.32.2.257.
- Iorember, F.M. (2018). Malnutrition in Chronic Kidney Disease. *Front Pediatr.* Jun 20;6:161. doi: 10.3389/fped.2018.00161.
- Iyengar, A., Raj, J.M., Vasudevan, A. (2021) Protein Energy Wasting in Children With Chronic Kidney Disease and End-Stage Kidney Disease: An Observational Study. *J Ren Nutr.* May;31(3):270-277. doi: 10.1053/j.jrn.2020.08.002
- Johansen, K.L., Young, B., Kaysen, G.A., Chertow, G.M. (2006) Association of body size with health status in patients undergoing dialysis. *Curr Opin Nephrol Hypertens*, 15(3): 210-217.
- Karakas, E., Colak, H., Gunes, F.E., Karakoyun, B. (2024) Determination of nutritional status and protein-energy wasting in patients with diabetic nephropathy. *North Clin Istanb.* Nov 19;11(6):560-568. doi: 10.14744/nci.2023.66503.
- KDOQI Work Group. (2009). KDOQI Clinical Practice Guideline for Nutrition in Children with CKD: 2008 update. Executive summary. *Am J Kidney Dis.* Mar;53(3 Suppl 2):S11-104. doi: 10.1053/j.ajkd.2008.11.017.

- Larkins, N., Kim, S., Craig, J., Hodson, E. (2016) Steroid-sensitive nephrotic syndrome: an evidence-based update of immunosuppressive treatment in children. *Arch Dis Child*. Apr;101(4):404-8. doi: 10.1136/archdischild-2015-308924.
- Madhileti, S., Selvam, S, Iyengar, A. (2023) Protein Energy Wasting And Clinical Outcomes Of Children with CKD2-5D- a Prospective Longitudinal Study, *Kidney International Reports*, Volume 8, Issue 3, Supplement. [doi:10.1016/j.ekir.2023.02.332](https://doi.org/10.1016/j.ekir.2023.02.332).
- Mastrangelo, A., Paglialonga, F., Edefonti, A. (2013). Assessment of nutritional status in children with chronic kidney disease and on dialysis. *Pediatric Nephrology*, 29(8), 1349–1358. doi:10.1007/s00467-013-2612-7.
- The Journal of nutrition, health and aging, Volume 23, Issue 1, 2019, Pages 96-101, doi.org/10.1007/s12603-018-1129-y
- Matyjek, A., Literacki, S., Niemczyk, S., Rymarz, A. (2020) Protein energy-wasting associated with nephrotic syndrome – the comparison of metabolic pattern in severe nephrosis to different stages of chronic kidney disease. *MC Nephrology* 21:346. doi:10.1186/s12882-020-02003-4.
- Mitsnefes, M., Flynn, J., Cohn, S., Samuels, J., Blydt. H. T., Saland, J., Kimball, T., Furth, S., Warady, B. (2010) CKiD Study Group: Masked hypertension associates with left ventricular hypertrophy in children with CKD. *J Am Soc Nephrol*. Jan;21(1):137-44. doi: 10.1681/ASN.2009060609.
- Norman, K., Stobäus, N., Pirlich, M., Bosy-W. A. (2012). Bioelectrical phase angle and impedance vector analysis – Clinical relevance and applicability of impedance parameters. *Clinical Nutrition*, 31(6), 854–861. doi:10.1016/j.clnu.2012.05.008
- North American Pediatric Renal Transplant Cooperative Study (NAPRTCS). (2008) Annual report. Rockville, MD: *The EMMES Corporation*
- Nourbakhsh, N., Rhee, C.M., Kalantar, K. (2014) Protein-energy wasting and uremic failure to thrive in children with chronic kidney disease: they are not small adults. *Pediatr Nephrol*. Dec;29(12):2249-52. doi: 10.1007/s00467-014-2898-0.
- Osunbor, O.A., Unuigbo, E.I., Okaka, E.I., Adejumo, O.A. (2023) Protein energy wasting in pre-dialysis chronic kidney disease patients in Benin City, Nigeria: A cross-sectional study. *PLoS ONE* 18(5): e0286075. doi:10.1371/journal.pone.0286075
- Panggabean, M. S. (2022). Nutrisi Pasien Anak dengan Chronic Kidney Disease (CKD). *Cermin Dunia Kedokteran*, 49(6), 320–326. <https://doi.org/10.55175/cdk.v49i6.240>.

- Perez, T. A., González, M. E., Valiente, S. J. B, Bajo, M. A., Celadilla, D. O, López, A. M. (2018) Protein-energy wasting syndrome in advanced chronic kidney disease: Prevalence and specific clinical characteristics. *Nefrologia*. 2018;38:141–151.
- Ramayani R., Rosmayanti., Ramayati R., Rusdidjas. (2013) Luaran pasien anak dengan gagal ginjal terminal. *Sari Pediatri*, Vol. 14, No. 5, Februari 201
- Rees, L., (2021) Protein energy wasting: what is it and what can we do to prevent it? *Pediatr Nephrol*. Feb;36(2):287-294. doi: 10.1007/s00467-019-04424-2.
- Ruperto, M., Sánchez, M. F. J., Barril, G. (2016) Predictors of protein-energy wasting in haemodialysis patients: a cross-sectional study. *J Hum Nutr Diet*. Feb;29(1):38-47. doi: 10.1111/jhn.12276.
- Sanna, C.S., Ravani, P., Corbani, V., Parodi, S., Haupt, R., Piaggio, G., Innocenti, M.L., Somenzi, D., Trivelli, A., Caridi, G., Izzi, C., Scolari, F., Mattioli, G., Allegri, L., Ghiggeri, G.M. (2009) Renal outcome in patients with congenital anomalies of the kidney and urinary tract. *Kidney Int*. 2009 Sep;76(5):528-33. doi: 10.1038/ki.2009.220.
- Santos, E.J.F., Dias, R.S.C., Lima, J.F.B., Salgado, F.N., Miranda, D.S.A. (2020) Erythropoietin Resistance in Patients with Chronic Kidney Disease: Current Perspectives. *Int J Nephrol Renovasc Dis*. Oct 8;13:231-237. doi: 10.2147/IJNRD.S239151.
- Schlesselman, J. J. (1982). *Case-Control Studies: Design, Conduct, Analysis*. Oxford University Press.
- Sheptulina, A. F., Lyusina, E. O., Mamutova, E. M., Yafarova, A. A., Kiselev, A. R., & Drapkina, O. M. (2024) Bioelectrical Impedance Analysis Demonstrates Reliable Agreement with Dual-Energy X-ray Absorptiometry in Identifying Reduced Skeletal Muscle Mass in Patients with Metabolic Dysfunction-Associated Steatotic Liver Disease and Hypertension. *Diagnostics*, 14(20), 2301. doi:10.3390/diagnostics14202301.
- Siew, E. D. dan Ikizler, T. A. (2010). Insulin Resistance and Protein Energy Metabolism in Patients with Advanced Chronic Kidney Disease. *Seminars in Dialysis*, 23(4), 378–382. doi:10.1111/j.1525-139x.2010.00763.x.
- Tan, R.S., Liang, D. H., Liu, Y., Zhong, X. S., Zhang, D. S., Ma, J. (2019) Bioelectrical Impedance Analysis–Derived Phase Angle Predicts Protein–Energy Wasting in Maintenance Hemodialysis Patients, *Journal of Renal Nutrition*, doi:10.1053/j.jrn.2018.09.001.
- Stenvinkel, P. (2006) Inflammation in end-stage renal disease: The hidden enemy (Review Article). *Nephrology*, 11: 36-41. doi.org/10.1111/j.1440-1797.2006.00541.x



- Tian, M., Yuan, J., Yu, F., He, P., Zhang, Q., Zha, Y. (2023) Decreased intracellular water is associated with sarcopenic obesity in chronic haemodialysis patients. *BMC Geriatr.* Oct 6;23(1):630. doi: 10.1186/s12877-023-04357-4. PMID: 37803331; PMCID: PMC10559522.
- Todisco, T., Ubertini, G.M., Bizzarri, C., Loche, S., Cappa, M. (2024) Chronic Kidney Disease and Growth Failure in Children. *Children*; 11(7):808. <https://doi.org/10.3390/children11070808>
- Wheeler D.C., dan Becker G.J. (2013) Summary of KDIGO guideline. What do we really know about management of blood pressure in patients with chronic kidney disease? *Kidney Int.* 83(3):377–383. doi: 10.1038/ki.2012.425.
- Windahl, K., Faxén, G.I., Almquist, T., Korkeila, M.L (2018) Prevalence and Risk of Protein-Energy Wasting Assessed by Subjective Global Assessment in Older Adults With Advanced Chronic Kidney Disease: Results From the EQUAL Study, *Journal of Renal Nutrition*, Volume 28, Issue 3. doi:10.1053/j.jrn.2017.11.002.
- Yuste, C., Barraca, D., Aragoncillo, S.I., Vega, M.A., Abad, S, Verdalles, G.Ú., Ruiz, C.C., Ampuero, J., López, G.J.M. (2013) Factors related with the progression of chronic kidney disease. *Nefrologia.* 2013;33(5):685-91. doi: 10.3265/Nefrologia.pre2013.May.11900.
- Zhao, Y., Gong, J., Ji, Y. (2023) Cross-sectional study of characteristics of body composition of 24,845 children and adolescents aged 3–17 years in Suzhou. *BMC Pediatr* 23, 358 doi: 10.1186/s12887-023-04134-7.