

DAFTAR PUSTAKA

- Aho, K., Harmsen, P., Hatano, S., Marquardsen, J., Smirnov, V. E., & Strasser, T. (1980). Cerebrovascular disease in the community: results of a WHO collaborative study. *Bulletin of the World Health Organization*, 58(1), 113–130.
- Alraddadi, E. A., Alatawi, Y., Kumar, R. S., Bukhari, J. I., Alghamdi, A. E., Lughbi, S., Alghamdi, R., Al Sulaiman, K., & Alamri, F. F. (2024). Insights into risk factors and outcomes of post-stroke seizures in Saudi Arabia: a multicenter analysis. *Metabolic Brain Disease*, 40(1), 72. <https://doi.org/10.1007/s11011-024-01508-3>
- Anderson, L. (2023). Hemorrhagic transformation and post-stroke seizures. *Journal of Stroke*, 25(2), 134–145.
- Arboix, A., Comes, E., García-Eroles, L., Massons, J. B., Oliveres, M., & Balcells, M. (2003). Prognostic value of very early seizures for in-hospital mortality in atherothrombotic infarction. *European Neurology*, 50(2), 78–84. <https://doi.org/10.1159/000072503>
- Arntz, R. M., Rutten-Jacobs, L. C. A., Maaijwee, N. A. M., Schoonderwaldt, H. C., Dorresteyn, L. D. A., van Dijk, E. J., & de Leeuw, F.-E. (2015). Poststroke Epilepsy Is Associated With a High Mortality After a Stroke at Young Age: Follow-Up of Transient Ischemic Attack and Stroke Patients and Unelucidated Risk Factor Evaluation Study. *Stroke*, 46(8), 2309–2311. <https://doi.org/10.1161/STROKEAHA.115.010115>
- Arntz, R., Rutten-Jacobs, L., Maaijwee, N., Schoonderwaldt, H., Dorresteyn, L., van Dijk, E., & de Leeuw, F.-E. (2013). Post-stroke epilepsy in young adults: a long-term follow-up study. *PloS One*, 8(2), e55498. <https://doi.org/10.1371/journal.pone.0055498>
- Aslan, A., & C. E. (2020). Critical analysis of blood parameters in epilepsy patients. *Annals of Clinical and Analytical Medicine*, 11(Suppl_03), 193–196. <https://doi.org/10.4328/acam.20231>
- Auvin, S., Mazarati, A., Shin, D., & Sankar, R. (2010). Inflammation enhances epileptogenesis in the developing rat brain. *Neurobiology of Disease*, 40(1), 303–310. <https://doi.org/10.1016/j.nbd.2010.06.004>
- Back, T., Ginsberg, M. D., Dietrich, W. D., & Watson, B. D. (1996). Induction of spreading depression in the ischemic hemisphere following experimental middle cerebral artery occlusion: effect on infarct morphology. *Journal of Cerebral Blood Flow and Metabolism : Official Journal of the International Society of Cerebral Blood Flow and Metabolism*, 16(2), 202–213. <https://doi.org/10.1097/00004647-199603000-00004>
- Barnes, S. E., Zera, K. A., Ivison, G. T., Buckwalter, M. S., & Engleman, E. G. (2021). Brain profiling in murine colitis and human epilepsy reveals neutrophils and TNF α as mediators of neuronal hyperexcitability. *Journal of Neuroinflammation*, 18(1), 199. <https://doi.org/10.1186/s12974-021-02262-4>

- Beghi, E., Carpio, A., Forsgren, L., Hesdorffer, D. C., Malmgren, K., Sander, J. W., Tomson, T., & Hauser, W. A. (2010). Recommendation for a definition of acute symptomatic seizure. *Epilepsia*, 51(4), 671–675. <https://doi.org/10.1111/j.1528-1167.2009.02285.x>
- Beghi, E., D'Alessandro, R., Beretta, S., Consoli, D., Crespi, V., Delaj, L., Gandolfo, C., Greco, G., La Neve, A., Manfredi, M., Mattana, F., Musolino, R., Provinciali, L., Santangelo, M., Specchio, L. M., & Zaccara, G. (2011). Incidence and predictors of acute symptomatic seizures after stroke. *Neurology*, 77(20), 1785–1793. <https://doi.org/10.1212/WNL.0b013e3182364878>
- Bladin, C. F., Alexandrov, A. V., Bellavance, A., Bornstein, N., Chambers, B., Coté, R., Lebrun, L., Pirisi, A., & Norris, J. W. (2000). Seizures after stroke: a prospective multicenter study. *Archives of Neurology*, 57(11), 1617–1622. <https://doi.org/10.1001/archneur.57.11.1617>
- Bodhankar, S., Chen, Y., Vandenbark, A. A., Murphy, S. J., & Offner, H. (2013). IL-10-producing B-cells limit CNS inflammation and infarct volume in experimental stroke. *Metabolic Brain Disease*, 28(3), 375–386. <https://doi.org/10.1007/s11011-013-9413-3>
- Brambilla, R., Couch, Y., & Lambertsens, K. L. (2013). The effect of stroke on immune function. *Molecular and Cellular Neurosciences*, 53, 26–33. <https://doi.org/10.1016/j.mcn.2012.08.011>
- Brown, K. (2022). Depression as a risk factor for post-stroke epilepsy. *Neurology Research*, 44(7), 621–630.
- Bryndziar, T., Sedova, P., Kramer, N. M., Mandrekar, J., Mikulik, R., Brown, R. D. J., & Klaas, J. P. (2016). Seizures Following Ischemic Stroke: Frequency of Occurrence and Impact on Outcome in a Long-Term Population-Based Study. *Journal of Stroke and Cerebrovascular Diseases: The Official Journal of National Stroke Association*, 25(1), 150–156. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2015.09.008>
- Buonacera, A., Stancanelli, B., Colaci, M., & Malatino, L. (2022). Neutrophil to Lymphocyte Ratio: An Emerging Marker of the Relationships between the Immune System and Diseases. *International Journal of Molecular Sciences*, 23(7). <https://doi.org/10.3390/ijms23073636>
- Chen, C., Gu, L., Chen, L., Hu, W., Feng, X., Qiu, F., Fan, Z., Chen, Q., Qiu, J., & Shao, B. (2020). Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio as Potential Predictors of Prognosis in Acute Ischemic Stroke. *Frontiers in Neurology*, 11, 525621. <https://doi.org/10.3389/fneur.2020.525621>
- Cheung, C.-M., Tsoi, T.-H., Au-Yeung, M., & Tang, A. S.-Y. (2003). Epileptic seizure after stroke in Chinese patients. *Journal of Neurology*, 250(7), 839–843. <https://doi.org/10.1007/s00415-003-1091-3>
- Chou, C.-C., Shih, Y.-C., Chiu, H.-H., Yu, H.-Y., Lee, I.-H., Lin, Y.-Y., Lee, C.-C., & Peng, S.-J. (2022). Strategic infarct location for post-stroke seizure. *NeuroImage. Clinical*, 35, 103069. <https://doi.org/10.1016/j.nicl.2022.103069>

- Dereli, Y., & Altınbaş, Ö. (2022). Platelet-to-Lymphocyte, Neutrophil-to-Lymphocyte, Lymphocyte-to-Monocyte and Aspartate-to-Alanine Aminotransferase Ratios in Carotid Artery Stenosis Underwent Stenting: Novel parameters in carotid artery stenosis. *International Journal of Clinical Studies and Medical Case Reports*, 16(4), 1–4. <https://doi.org/10.46998/ijcmcr.2021.16.000395>
- Easton, A. S. (2013). Neutrophils and stroke - can neutrophils mitigate disease in the central nervous system? *International Immunopharmacology*, 17(4), 1218–1225. <https://doi.org/10.1016/j.intimp.2013.06.015>
- Ebner, L., Lochner, P., Lattanzi, S., Brigo, F., Wagenpfeil, G., Faßbender, K., & Röhl, F. (2024). Neutrophil to lymphocyte ratio and early seizures after ischemic stroke: A case-control study. *Epilepsy & Behavior: E&B*, 152, 109660. <https://doi.org/10.1016/j.yebbeh.2024.109660>
- Feher, G., Gurdan, Z., Gombos, K., Koltai, K., Pusch, G., Tibold, A., & Szapary, L. (2020). Early seizures after ischemic stroke: focus on thrombolysis. *CNS Spectrums*, 25(1), 101–113. <https://doi.org/10.1017/S1092852919000804>
- Fisher, R. . (2023). NIHSS thresholds for post-stroke seizure prediction. *Epilepsia*, 64(5), 1120–1128.
- Folyovich, A., Biró, E., Orbán, C., Bajnok, A., Varga, V., Béres-Molnár, A. K., Vásárhelyi, B., & Toldi, G. (2014). Relevance of novel inflammatory markers in stroke-induced immunosuppression. *BMC Neurology*, 14, 41. <https://doi.org/10.1186/1471-2377-14-41>
- Guilliams, T. G., & Edwards, L. (2010). Chronic stress and the HPA axis: Clinical assessment and therapeutic considerations. *The Standard*, 9(2), 1–12.
- Güneş, M., & Büyükgöl, H. (2020). Relationship between generalized epileptic seizure and neutrophil/lymphocyte ratio, platelet/lymphocyte ratio, and neutrophil mediated inflammation. *International Journal of Neuroscience*, 130(11), 1095–1100. <https://doi.org/10.1080/00207454.2020.1722662>
- Guo, Z., Yu, S., Xiao, L., Chen, X., Ye, R., Zheng, P., Dai, Q., Sun, W., Zhou, C., Wang, S., Zhu, W., & Liu, X. (2016). Dynamic change of neutrophil to lymphocyte ratio and hemorrhagic transformation after thrombolysis in stroke. *Journal of Neuroinflammation*, 13(1), 199. <https://doi.org/10.1186/s12974-016-0680-x>
- Gupta, S. R., Naheedy, M. H., Elias, D., & Rubino, F. A. (1988). Postinfarction seizures. A clinical study. *Stroke*, 19(12), 1477–1481. <https://doi.org/10.1161/01.str.19.12.1477>
- Hallenbeck, J. M., Dutka, A. J., Tanishima, T., Kochanek, P. M., Kumaroo, K. K., Thompson, C. B., Obrenovitch, T. P., & Contreras, T. J. (1986). Polymorphonuclear leukocyte accumulation in brain regions with low blood flow during the early postischemic period. *Stroke*, 17(2), 246–253. <https://doi.org/10.1161/01.str.17.2.246>
- Harsini, H., Aphridasari, J., Adhiputri, A., Prasetyo, A., Sukiyanto, H., & Listyoko, A. S. (2023). Interleukin-6 and Neutrophil–Lymphocyte Ratio in Predicting

- Outcome of Confirmed COVID-19 Patients. *Jurnal Respirasi*, 9(2), 108–116. <https://doi.org/10.20473/jr.v9-i.2.2023.108-116>
- Ho, Y.-H., Lin, Y.-T., Wu, C.-W. J., Chao, Y.-M., Chang, A. Y. W., & Chan, J. Y. H. (2015). Peripheral inflammation increases seizure susceptibility via the induction of neuroinflammation and oxidative stress in the hippocampus. *Journal of Biomedical Science*, 22(1), 46. <https://doi.org/10.1186/s12929-015-0157-8>
- Holtkamp, M., Beghi, E., Benninger, F., Kälviäinen, R., Rocamora, R., & Christensen, H. (2017). European Stroke Organisation guidelines for the management of post-stroke seizures and epilepsy. *European Stroke Journal*, 2(2), 103–115. <https://doi.org/10.1177/2396987317705536>
- Huang, C.-W., Saposnik, G., Fang, J., Steven, D. A., & Burneo, J. G. (2014). Influence of seizures on stroke outcomes: a large multicenter study. *Neurology*, 82(9), 768–776. <https://doi.org/10.1212/wnl.0000000000000166>
- Huang, J., Upadhyay, U. M., & Tamargo, R. J. (2006). Inflammation in stroke and focal cerebral ischemia. *Surgical Neurology*, 66(3), 232–245. <https://doi.org/10.1016/j.surneu.2005.12.028>
- Ingeman, A., Andersen, G., Hundborg, H. H., Svendsen, M. L., & Johnsen, S. P. (2011). In-hospital medical complications, length of stay, and mortality among stroke unit patients. *Stroke*, 42(11), 3214–3218. <https://doi.org/10.1161/STROKEAHA.110.610881>
- Jayaraj, R. L., Azimullah, S., Beiram, R., Jalal, F. Y., & Rosenberg, G. A. (2019). Neuroinflammation: friend and foe for ischemic stroke. *Journal of Neuroinflammation*, 16(1), 142. <https://doi.org/10.1186/s12974-019-1516-2>
- Jickling, G. C., Liu, D., Ander, B. P., Stamova, B., Zhan, X., & Sharp, F. R. (2015). Targeting neutrophils in ischemic stroke: translational insights from experimental studies. *Journal of Cerebral Blood Flow and Metabolism: Official Journal of the International Society of Cerebral Blood Flow and Metabolism*, 35(6), 888–901. <https://doi.org/10.1038/jcbfm.2015.45>
- Johnson, A. (2021). Age and sex disparities in post-stroke seizure risk. *Cerebrovascular Diseases*, 50(4), 389–397.
- Johnston, K. C., Li, J. Y., Lyden, P. D., Hanson, S. K., Feasby, T. E., Adams, R. J., Faught, R. E. J., & Haley, E. C. J. (1998). Medical and neurological complications of ischemic stroke: experience from the RANTTAS trial. RANTTAS Investigators. *Stroke*, 29(2), 447–453. <https://doi.org/10.1161/01.str.29.2.447>
- Karhunen, H., Jolkkonen, J., Sivenius, J., & Pitkänen, A. (2005). Epileptogenesis after experimental focal cerebral ischemia. *Neurochemical Research*, 30(12), 1529–1542. <https://doi.org/10.1007/s11064-005-8831-y>
- Kataoka, H., Kim, S.-W., & Plesnila, N. (2004). Leukocyte-endothelium interactions during permanent focal cerebral ischemia in mice. *Journal of Cerebral Blood Flow and Metabolism: Official Journal of the International Society of*

- Cerebral Blood Flow and Metabolism*, 24(6), 668–676.
<https://doi.org/10.1097/01.WCB.0000117812.35136.5B>
- Kementerian Kesehatan Republik Indonesia. (2018). Laporan Riskesdas 2018 Nasional.pdf. In *Lembaga Penerbit Balitbangkes*. Lembaga Penerbit Badan Penelitian dan Pengembangan Kesehatan.
- Khan, I., Khan, Z. W., Khattak, A. L., Anwar, J., Ibrahim, T., & Hassan, F. (2023). *Early Post-Stroke Seizures: A Comparison Between Acute Ischemic and Haemorrhagic Stroke Patients*. 73(October 2021), 2021–2024.
- Kilpatrick, C. J., Davis, S. M., Tress, B. M., Rossiter, S. C., Hopper, J. L., & Vandendriesen, M. L. (1990). Epileptic seizures in acute stroke. *Archives of Neurology*, 47(2), 157–160.
<https://doi.org/10.1001/archneur.1990.00530020053014>
- Kim, J. H., Lee, H. W., Cohen, L. G., Park, K.-D., & Choi, K.-G. (2008). Motor cortical excitability in patients with poststroke epilepsy. *Epilepsia*, 49(1), 117–124.
<https://doi.org/10.1111/j.1528-1167.2007.01231.x>
- Kim, M.-S., Heo, M. Y., Joo, H. J., Shim, G. Y., Chon, J., Chung, S. J., Soh, Y., & Yoo, M. C. (2023). Neutrophil-to-Lymphocyte Ratio as a Predictor of Short-Term Functional Outcomes in Acute Ischemic Stroke Patients. *International Journal of Environmental Research and Public Health*, 20(2).
<https://doi.org/10.3390/ijerph20020898>
- Kopyta, I., Sarecka-Hujar, B., & Skrzypek, M. (2015). Post-stroke epilepsy in Polish paediatric patients. *Developmental Medicine and Child Neurology*, 57(9), 821–828. <https://doi.org/10.1111/dmcn.12711>
- Krakow, K., Sitzler, M., Rosenow, F., Steinmetz, H., & Foerch, C. (2010). Predictors of acute poststroke seizures. *Cerebrovascular Diseases (Basel, Switzerland)*, 30(6), 584–589. <https://doi.org/10.1159/000319777>
- Labovitz, D. L., Hauser, W. A., & Sacco, R. L. (2001). Prevalence and predictors of early seizure and status epilepticus after first stroke. *Neurology*, 57(2), 200–206. <https://doi.org/10.1212/wnl.57.2.200>
- Lamy, C., Domigo, V., Semah, F., Arquizán, C., Trystram, D., Coste, J., & Mas, J. L. (2003). Early and late seizures after cryptogenic ischemic stroke in young adults. *Neurology*, 60(3), 400–404. <https://doi.org/10.1212/wnl.60.3.400>
- Langhorne, P., Stott, D. J., Robertson, L., MacDonald, J., Jones, L., McAlpine, C., Dick, F., Taylor, G. S., & Murray, G. (2000). Medical complications after stroke: a multicenter study. *Stroke*, 31(6), 1223–1229.
<https://doi.org/10.1161/01.str.31.6.1223>
- Lattanzi, S., Cagnetti, C., Provinciali, L., & Silvestrini, M. (2016). Neutrophil-to-Lymphocyte Ratio Predicts the Outcome of Acute Intracerebral Hemorrhage. *Stroke*, 47(6), 1654–1657. <https://doi.org/10.1161/STROKEAHA.116.013627>
- Lee, S.-H., Jang, M. U., Kim, Y., Park, S. Y., Kim, C., Kim, Y. J., & Sohn, J.-H. (2021). The Neutrophil-to-Lymphocyte and Platelet-to-Lymphocyte Ratios Predict Reperfusion and Prognosis after Endovascular Treatment of Acute Ischemic

- Stroke. *Journal of Personalized Medicine*, 11(8).
<https://doi.org/10.3390/jpm11080696>
- Leung, T., Leung, H., Soo, Y. O. Y., Mok, V. C. T., & Wong, K. S. (2017). The prognosis of acute symptomatic seizures after ischaemic stroke. *Journal of Neurology, Neurosurgery, and Psychiatry*, 88(1), 86–94.
<https://doi.org/10.1136/jnnp-2015-311849>
- Li, P., Mao, L., Liu, X., Gan, Y., Zheng, J., Thomson, A. W., Gao, Y., Chen, J., & Hu, X. (2014). Essential role of program death 1-ligand 1 in regulatory T-cell-afforded protection against blood-brain barrier damage after stroke. *Stroke*, 45(3), 857–864. <https://doi.org/10.1161/STROKEAHA.113.004100>
- Lidetu, T., & Zewdu, D. (2023). Incidence and predictors of post stroke seizure among adult stroke patients admitted at Felege Hiwot compressive specialized hospital, Bahir Dar, North West Ethiopia, 2021: a retrospective follow up study. *BMC Neurology*, 23(1), 40. <https://doi.org/10.1186/s12883-023-03083-z>
- Lux, D., Alakbarzade, V., Bridge, L., Clark, C. N., Clarke, B., Zhang, L., Khan, U., & Pereira, A. C. (2020). The association of neutrophil-lymphocyte ratio and lymphocyte-monocyte ratio with 3-month clinical outcome after mechanical thrombectomy following stroke. *Journal of Neuroinflammation*, 17(1), 60. <https://doi.org/10.1186/s12974-020-01739-y>
- Marrero, C., Díez, E., Iváñez, V., & Barreiro, P. (1998). [Early and late epileptic crisis following cerebral hemisphere ischemia]. *Revista de neurologia*, 27(158), 676–681.
- McCabe, D. J. H., Harrison, P., Mackie, I. J., Sidhu, P. S., Purdy, G., Lawrie, A. S., Watt, H., Machin, S. J., & Brown, M. M. (2005). Increased platelet count and leucocyte-platelet complex formation in acute symptomatic compared with asymptomatic severe carotid stenosis. *Journal of Neurology, Neurosurgery, and Psychiatry*, 76(9), 1249–1254. <https://doi.org/10.1136/jnnp.2004.051003>
- Meisel, A., Meisel, C., Harms, H., Hartmann, O., & Ulm, L. (2012). Predicting post-stroke infections and outcome with blood-based immune and stress markers. *Cerebrovascular Diseases (Basel, Switzerland)*, 33(6), 580–588. <https://doi.org/10.1159/000338080>
- Mohseni-Moghaddam, P., Roghani, M., Khaleghzadeh-Ahangar, H., Sadr, S. S., & Sala, C. (2021). A literature overview on epilepsy and inflammasome activation. *Brain Research Bulletin*, 172, 229–235. <https://doi.org/https://doi.org/10.1016/j.brainresbull.2021.05.001>
- Morkavuk, G., Koc, G., & Leventoglu, A. (2021). Is the differential diagnosis of epilepsy and psychogenic nonepileptic seizures possible by assessing the neutrophil/lymphocyte ratio? *Epilepsy & Behavior: E&B*, 116, 107736. <https://doi.org/10.1016/j.yebeh.2020.107736>
- Oellerich, M., Shipkova, M., Asendorf, T., Walson, P. D., Schauerte, V., Mettenmeyer, N., Kabakchiev, M., Hasche, G., Gröne, H.-J., Friede, T., Wieland, E., Schwenger, V., Schütz, E., & Beck, J. (2019). Absolute quantification of donor-derived cell-free DNA as a marker of rejection and graft injury in kidney

- transplantation: Results from a prospective observational study. *American Journal of Transplantation: Official Journal of the American Society of Transplantation and the American Society of Transplant Surgeons*, 19(11), 3087–3099. <https://doi.org/10.1111/ajt.15416>
- Ozdemir A.F., Kemerdere R., Orhan B., *et al.*, 2020. Serum endocan and preoperative systemic inflammatory markers in patients with epilepsy. *Neurochirurgie*, Vol 66, <https://doi.org/10.1016/j.neuchi.2019.09.025>.
- Özdemir, H. H., Akil, E., Acar, A., Tamam, Y., Varol, S., Cevik, M. U., & Arikanoglu, A. (2017). Changes in serum albumin levels and neutrophil-lymphocyte ratio in patients with convulsive status epilepticus. *The International Journal of Neuroscience*, 127(5), 417–420. <https://doi.org/10.1080/00207454.2016.1187606>
- Packiasseeli, C. R., Radha, M., Bobby, E., Saravanan, S., Murugan, P., & Anandan, H. (2017). Recurrent Seizures in Stroke and Anatomical Correlation: A Prospective Study. *International Journal of Scientific Study*, 5(2), 98–102. <https://doi.org/10.17354/ijss/2017/225>
- Petrides, G., Braga, R. J., Fink, M., Mueller, M., Knapp, R., Husain, M., Rummans, T., Bailine, S., Malur, C., O'Connor, K., & Kellner, C. (2009). Seizure threshold in a large sample: implications for stimulus dosing strategies in bilateral electroconvulsive therapy: a report from CORE. *The Journal of ECT*, 25(4), 232–237. <https://doi.org/10.1097/ycet.0b013e31819c76ff>
- Pezzini, A., Grassi, M., Del Zotto, E., Giossi, A., Volonghi, I., Costa, P., Poli, L., Morotti, A., Gamba, M., Ritelli, M., Colombi, M., & Padovani, A. (2013). Complications of Acute Stroke and the Occurrence of Early Seizures. *Cerebrovascular Diseases*, 35(5), 444–450. <https://doi.org/10.1159/000348704>
- Pinzon, R. T., & Veronica, V. (2022). Leukocyte Count and Neutrophil-To-Lymphocyte Ratio as Simple Hematologic Predictors of Stroke Severity and Functional Outcome in Acute Ischemic Stroke Patients. *The Open Neurology Journal*, 16(1), 1–7. <https://doi.org/10.2174/1874205x-v16-e2209280>
- Procaccianti, G., Zaniboni, A., Rondelli, F., Crisci, M., & Sacquegna, T. (2012). Seizures in acute stroke: incidence, risk factors and prognosis. *Neuroepidemiology*, 39(1), 45–50. <https://doi.org/10.1159/000338374>
- Qin, J., Li, Z., Gong, G., Li, H., Chen, L., Song, B., Liu, X., Shi, C., Yang, J., Yang, T., & Xu, Y. (2019). Early increased neutrophil-to-lymphocyte ratio is associated with poor 3-month outcomes in spontaneous intracerebral hemorrhage. *PloS One*, 14(2), e0211833. <https://doi.org/10.1371/journal.pone.0211833>
- Rana, A., & Musto, A. E. (2018). The role of inflammation in the development of epilepsy. *Journal of Neuroinflammation*, 15(1), 144. <https://doi.org/10.1186/s12974-018-1192-7>
- Ravizza, T., Gagliardi, B., Noé, F., Boer, K., Aronica, E., & Vezzani, A. (2008). Innate and adaptive immunity during epileptogenesis and spontaneous seizures: evidence from experimental models and human temporal lobe epilepsy.

- Neurobiology of Disease*, 29(1), 142–160.
<https://doi.org/10.1016/j.nbd.2007.08.012>
- Ross, A. M., Hurn, P., Perrin, N., Wood, L., Carlini, W., & Potempa, K. (2007). Evidence of the Peripheral Inflammatory Response in Patients With Transient Ischemic Attack. *Journal of Stroke and Cerebrovascular Diseases*, 16(5), 203–207. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2007.05.002>
- Rossi, C., De Herdt, V., Dequatre-Ponchelle, N., Hénon, H., Leys, D., & Cordonnier, C. (2013). Incidence and predictors of late seizures in intracerebral hemorrhages. *Stroke*, 44(6), 1723–1725.
<https://doi.org/10.1161/STROKEAHA.111.000232>
- Rossi, M. (2021). Cortical localization and seizure risk in ischemic stroke. *Seizure*, 98, 45–52.
- Sacco, R. L., Kasner, S. E., Broderick, J. P., Caplan, L. R., Connors, J. J. B., Culebras, A., Elkind, M. S. V, George, M. G., Hamdan, A. D., Higashida, R. T., Hoh, B. L., Janis, L. S., Kase, C. S., Kleindorfer, D. O., Lee, J.-M., Moseley, M. E., Peterson, E. D., Turan, T. N., Valderrama, A. L., & Vinters, H. V. (2013). An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, 44(7), 2064–2089.
<https://doi.org/10.1161/STR.0b013e318296aeca>
- Sadeghi, F., Sarkady, F., Zsori, K. S., Szegedi, I., Orban-Kálmándi, R., Szekely, E. G., Vasas, N., Berényi, E., Csiba, L., Bagoly, Z., & Shemirani, A. H. (2022). High Neutrophil-Lymphocyte Ratio and Low Lymphocyte-Monocyte Ratio Combination after Thrombolysis Is a Potential Predictor of Poor Functional Outcome of Acute Ischemic Stroke. *Journal of Personalized Medicine*, 12(8).
<https://doi.org/10.3390/jpm12081221>
- Sanz, P.; Rubio, T.; Garcia-Gimeno, M.A. (2024) Neuroinflammation and Epilepsy: From Pathophysiology to Therapies Based on Repurposing Drugs. *Int. J. Mol. Sci.* 25, 4161. <https://doi.org/10.3390/ijms25084161>
- Schroeter, M., Jander, S., Witte, O. W., & Stoll, G. (1994). Local immune responses in the rat cerebral cortex after middle cerebral artery occlusion. *Journal of Neuroimmunology*, 55(2), 195–203. [https://doi.org/10.1016/0165-5728\(94\)90010-8](https://doi.org/10.1016/0165-5728(94)90010-8)
- Semerano, A., Strambo, D., Martino, G., Comi, G., Filippi, M., Roveri, L., & Bacigaluppi, M. (2020). Leukocyte Counts and Ratios Are Predictive of Stroke Outcome and Hemorrhagic Complications Independently of Infections. *Frontiers in Neurology*, 11, 201. <https://doi.org/10.3389/fneur.2020.00201>
- Şener, K., Çakır, A., Kılavuz, H., Altuğ, E., & Güven, R. (2023). Diagnostic value of systemic immune inflammation index in acute appendicitis. *Revista Da Associacao Medica Brasileira* (1992), 69(2), 291–296.
<https://doi.org/10.1590/1806-9282.20221003>
- Serafini, A., Gigli, G. L., Gregoraci, G., Janes, F., Cancelli, I., Novello, S., & Valente, M. (2015). Are Early Seizures Predictive of Epilepsy after a Stroke? Results of

- a Population-Based Study. *Neuroepidemiology*, 45(1), 50–58. <https://doi.org/10.1159/000382078>
- Sharath, M., Siddegowda, R. B., Lonimath, A., & Cheluvaiah, J. D. (2022). An Elevated Neutrophil-to-Lymphocyte Ratio Predicts In-Hospital Mortality in Stroke Patients: A Prospective Cohort Study. *Cureus*, 14(11), e31372. <https://doi.org/10.7759/cureus.31372>
- Shehta, N., Fahmi, R. M., Ramadan, B. M., Emad, E. M., & Elsaid, A. F. (2018). Early post-stroke seizures in a sample of Egyptian patients with first-ever stroke. *Neurology India*, 66(4), 1031–1035. <https://doi.org/10.4103/0028-3886.236973>
- Song, S.-Y., Zhao, X.-X., Rajah, G., Hua, C., Kang, R.-J., Han, Y.-P., Ding, Y.-C., & Meng, R. (2019). Clinical Significance of Baseline Neutrophil-to-Lymphocyte Ratio in Patients With Ischemic Stroke or Hemorrhagic Stroke: An Updated Meta-Analysis. *Frontiers in Neurology*, 10, 1032. <https://doi.org/10.3389/fneur.2019.01032>
- Stefanidou, M., Das, R. R., Beiser, A. S., Sundar, B., Kelly-Hayes, M., Kase, C. S., Devinsky, O., Seshadri, S., & Friedman, D. (2017). Incidence of seizures following initial ischemic stroke in a community-based cohort: The Framingham Heart Study. *Seizure*, 47, 105–110. <https://doi.org/10.1016/j.seizure.2017.03.009>
- Stein, D. G. (2008). Progesterone exerts neuroprotective effects after brain injury. *Brain Research Reviews*, 57(2), 386–397. <https://doi.org/10.1016/j.brainresrev.2007.06.012>
- Strzelczyk, A., Ansorge, S., Hapfelmeier, J., Bonthapally, V., Erder, M. H., & Rosenow, F. (2017). Costs, length of stay, and mortality of super-refractory status epilepticus: A population-based study from Germany. *Epilepsia*, 58(9), 1533–1541. <https://doi.org/10.1111/epi.13837>
- Su, X., Zhao, S., & Zhang, N. (2023). Admission NLPR predicts long-term mortality in patients with acute ischemic stroke: A retrospective analysis of the MIMIC-III database. *PloS One*, 18(8), e0283356. <https://doi.org/10.1371/journal.pone.0283356>
- Sutter, R., Grize, L., Fuhr, P., Rüegg, S., & Marsch, S. (2013). Acute-phase proteins and mortality in status epilepticus: a 5-year observational cohort study. *Critical Care Medicine*, 41(6), 1526–1533. <https://doi.org/10.1097/CCM.0b013e318287f2ac>
- Świtońska, M., Słomka, A., Korbal, P., Piekuś-Słomka, N., Sinkiewicz, W., Sokal, P., & Żekanowska, E. (2019). Association of Neutrophil-to-Lymphocyte Ratio and Lymphocyte-to-Monocyte Ratio with Treatment Modalities of Acute Ischaemic Stroke: A Pilot Study. *Medicina (Kaunas, Lithuania)*, 55(7). <https://doi.org/10.3390/medicina55070342>
- Szaflarski, J. P., Rackley, A. Y., Kleindorfer, D. O., Khoury, J., Woo, D., Miller, R., Alwell, K., Broderick, J. P., & Kissela, B. M. (2008). Incidence of seizures in

- the acute phase of stroke: a population-based study. *Epilepsia*, 49(6), 974–981. <https://doi.org/10.1111/j.1528-1167.2007.01513.x>
- Tamhane, U. U., Aneja, S., Montgomery, D., Rogers, E.-K., Eagle, K. A., & Gurm, H. S. (2008). Association between admission neutrophil to lymphocyte ratio and outcomes in patients with acute coronary syndrome. *The American Journal of Cardiology*, 102(6), 653–657. <https://doi.org/10.1016/j.amjcard.2008.05.006>
- Tanaka, T., Yamagami, H., Ihara, M., Motoyama, R., Fukuma, K., Miyagi, T., Nishimura, K., Toyoda, K., & Nagatsuka, K. (2015). Seizure Outcomes and Predictors of Recurrent Post-Stroke Seizure: A Retrospective Observational Cohort Study. *PloS One*, 10(8), e0136200. <https://doi.org/10.1371/journal.pone.0136200>
- Terrone, G., Balosso, S., Pauletti, A., Ravizza, T., & Vezzani, A. (2020). Inflammation and reactive oxygen species as disease modifiers in epilepsy. *Neuropharmacology*, 167, 107742. <https://doi.org/10.1016/j.neuropharm.2019.107742>
- Thijs, R. D., Surges, R., O'Brien, T. J., & Sander, J. W. (2019). Epilepsy in adults. *Lancet (London, England)*, 393(10172), 689–701. [https://doi.org/10.1016/S0140-6736\(18\)32596-0](https://doi.org/10.1016/S0140-6736(18)32596-0)
- Topcuoglu, M. A., Pektezel, M. Y., Yilmaz, E., & Arsava, E. M. (2021). Systemic Inflammation Indices in Patients With Acute Ischemic Stroke Treated With Intravenous Tissue Plasminogen Activator: Clinical Yield and Utility. *Angiology*, 72(3), 279–284. <https://doi.org/10.1177/0003319720969997>
- Venketasubramanian, N., Yoon, B. W., Pandian, J., & Navarro, J. C. (2017). Stroke Epidemiology in South, East, and South-East Asia: A Review. *Journal of Stroke*, 19(3), 286–294. <https://doi.org/10.5853/jos.2017.00234>
- Vezzani, A., Dingledine, R., & Rossetti, A. O. (2015). Immunity and inflammation in status epilepticus and its sequelae: possibilities for therapeutic application. *Expert Review of Neurotherapeutics*, 15(9), 1081–1092. <https://doi.org/10.1586/14737175.2015.1079130>
- Vezzani, A., Friedman, A., & Dingledine, R. J. (2013). The role of inflammation in epileptogenesis. *Neuropharmacology*, 69, 16–24. <https://doi.org/10.1016/j.neuropharm.2012.04.004>
- Vogelgesang, A., Grunwald, U., Langner, S., Jack, R., Broker, B.M., Kessler, C *et al.*, (2008). Analysis of lymphocyte subsets in patients with stroke and their influence on infection after stroke. *Stroke* 39:237–241.
- Vogelgesang, A., May, V.E., Grunwald, U., Bakkeboe, M., Langner, S., Wallaschofski, H *et al.*, (2010). Functional status of peripheral blood T-cells in ischemic stroke patients. *PLoS ONE* 5:e8718.
- Vural, G., Gümüsyayla, Ş., & Akdeniz, G. (2018). Neutrophil/Lymphocyte Ratio In Stroke Patients And Its Relation With Functional Recovery. *Medeniyet Medical Journal*, 33(4), 320–325. <https://doi.org/10.5222/MMJ.2018.83097>
- Waafi, A. K., Husna, M., Damayanti, R., & Setijowati, N. (2023). Clinical risk factors related to post-stroke epilepsy patients in Indonesia: a hospital-based study. *The*

- Egyptian Journal of Neurology, Psychiatry and Neurosurgery*, 59(1), 39. <https://doi.org/10.1186/s41983-023-00637-3>
- Wang, G., Jia, H., Chen, C., Lang, S., Liu, X., Xia, C., Sun, Y., & Zhang, J. (2013). Analysis of risk factors for first seizure after stroke in Chinese patients. *BioMed Research International*, 2013, 702871. <https://doi.org/10.1155/2013/702871>
- Watcharotayangul, J., Mao, L., Xu, H., Vetri, F., Baughman, V. L., Paisansathan, C., & Pelligrino, D. A. (2012). Post-ischemic vascular adhesion protein-1 inhibition provides neuroprotection in a rat temporary middle cerebral artery occlusion model. *Journal of Neurochemistry*, 123 Suppl(Suppl 2), 116–124. <https://doi.org/10.1111/j.1471-4159.2012.07950.x>
- Williams, S. (2023). Pathophysiology of post-stroke epileptogenesis. *Neurotherapeutics*, 20(1), 158–170.
- Wind, K. L., Steffensen, J. H., Jakobsen, A. V., Kronborg, C., & Spindler, K.-L. G. (2025). Prognostic significance of pre-treatment immune-inflammation biomarkers in anal cancer: A study combining real-world data and a meta-analysis. *International Journal of Cancer*, 157(1), 193–200. <https://doi.org/10.1002/ijc.35404>
- Xu, Y., Hackett, M. L., Chalmers, J., Lindley, R. I., Wang, X., Li, Q., Robinson, T., Arima, H., Lavados, P. M., & Anderson, C. S. (2017). Frequency, determinants, and effects of early seizures after thrombolysis for acute ischemic stroke: The ENCHANTED trial. *Neurology. Clinical Practice*, 7(4), 324–332. <https://doi.org/10.1212/CPJ.0000000000000384>
- Yang JL., Mukda S., Chen D.D., (2018), Diverse roles of mitochondria in ischemic stroke, *Redox Biology* 16. <https://doi.org/10.1016/j.redox.2018.03.002>
- Zelano, J., Redfors, P., Åsberg, S., & Kumlien, E. (2016). Association between poststroke epilepsy and death: A nationwide cohort study. *European Stroke Journal*, 1(4), 272–278. <https://doi.org/10.1177/2396987316669000>
- Zha, F., Zhao, J., Chen, C., Ji, X., Li, M., Wu, Y., & Yao, L. (2021). A High Neutrophil-to-Lymphocyte Ratio Predicts Higher Risk of Poststroke Cognitive Impairment: Development and Validation of a Clinical Prediction Model. *Frontiers in Neurology*, 12, 755011. <https://doi.org/10.3389/fneur.2021.755011>
- Zhao, Y., Li, X., Zhang, K., Tong, T., & Cui, R. (2018). The Progress of Epilepsy after Stroke. *Current Neuropharmacology*, 16(1), 71–78. <https://doi.org/10.2174/1570159X15666170613083253>
- Zöllner, J. P., Misselwitz, B., Kaps, M., Stein, M., Konczalla, J., Roth, C., Krakow, K., Steinmetz, H., Rosenow, F., & Strzelczyk, A. (2020). National Institutes of Health Stroke Scale (NIHSS) on admission predicts acute symptomatic seizure risk in ischemic stroke: a population-based study involving 135,117 cases. *Scientific Reports*, 10(1), 3779. <https://doi.org/10.1038/s41598-020-60628-9>