

DAFTAR PUSTAKA

- Abdullah, N., Abdul Murad, N. A., Attia, J., Oldmeadow, C., Mohd Haniff, E. A., Syafruddin, S.E., Abd Jalal, N., et al. (2015). Characterizing the genetic risk for Type 2 diabetes in a Malaysian multi-ethnic cohort. *Diabet Med.*, *32*(10), 1377–1384. <https://doi.org/10.1111/dme.12735>
- Abdullah, N., Attia, J., Oldmeadow, C., Scott, R. J., & Holliday, E. G. (2014). The architecture of risk for type 2 diabetes: Understanding asia in the context of global findings. *Int J Endocrinol.*, *2014*(March). <https://doi.org/10.1155/2014/593982>
- Adeva-Andany, M. M., Rañal-Muñoz, E., Vila-Altesor, M., Fernández-Fernández, C., Funcasta- Calderón, R., & Castro-Quintela, E. (2019). Dietary habits contribute to define the risk of type 2 diabetes in humans. *Clin Nutr ESPEN*, *34*, 8–17. <https://doi.org/10.1016/j.clnesp.2019.08.002>
- Ahmed, M. S., Shuvo, S. D., Paul, D. K., Karim, M. R., Kamruzzaman, M., Mahmud, N., Ferdous, M. J., & Elahi, M. T. (2021). Prevalence of dyslipidemia and associated risk factors among newly diagnosed Type-2 Diabetes Mellitus (T2DM) patients in Kushtia, Bangladesh. *PLOS global public health*, *1*(12), e0000003. <https://doi.org/10.1371/journal.pgph.0000003>
- Alam, S., Hasan, M. K., Neaz, S., Hussain, N., Hossain, M. F., & Rahman, T. (2021). Diabetes Mellitus: Insights from Epidemiology, Biochemistry, Risk Factors, Diagnosis, Complications and Comprehensive Management. *Diabetology*, *2*(2), 36–50. <https://doi.org/10.3390/diabetology2020004>
- Allibert, C. (2008). Austronesian Migration and the Establishment of the Malagasy Civilization : Contrasted Readings in Linguistics , Archaeology , Genetics and Cultural Anthropology. 7–16. <https://doi.org/10.1177/0392192108090734>
- Alsulami, S., Aji, A. S., Ariyasra, U., Sari, S. R., Tasrif, N., Yani, F. F., Lovegrove, J. A., et al. (2020). Interaction between the genetic risk score and dietary protein intake on cardiometabolic traits in Southeast Asian. *Genes Nutr*, *15*(1). <https://doi.org/10.1186/s12263-020-00678-w>
- American Association of Diabetes. (2021). ADA standards of diabetes care 2021. In *Diabetes Care* (Vol. 44, pp. S21–S226).
- An, D. B., Ann, S., Seok, S., Kang, Y., & Lee, S. (2023). *Hepatic Cdkal1 deletion regulates HDL catabolism and promotes reverse cholesterol transport*. *Atherosclerosis*, *375* (2022), 21–29. <https://doi.org/10.1016/j.atherosclerosis.2023.05.012>
- Anderson, C., Milne, G. L., Park, Y. M., Sandler, D. P., & Nichols, H. B. (2018). *Dietary Glycemic Index and Glycemic Load Are Positively Associated with Oxidative Stress among Premenopausal Women*. *The Journal of Nutrition*, *148*(1), 125–130. <https://doi.org/10.1093/jn/nxx022>

- Aris-Brosou, S. (2019). *Direct Evidence of an Increasing Mutational Load in Humans*. *Mol. Biol. Evol.*, 36(12), 2823–2829. <https://doi.org/10.1093/molbev/msz192>
- Arroyo, M. N., Green, J. A., Cnop, M., & Igoillo-Esteve, M. (2021). Trna biology in the pathogenesis of diabetes: Role of genetic and environmental factors. *Int J Mol Sci*, 22(2), 1–27. <https://doi.org/10.3390/ijms22020496>
- Arksey, H., & O'Malley, L. (2005). Scoping studies: Towards a methodological framework. *International Journal of Social Research Methodology: Theory and Practice*, 8(1), 19–32. <https://doi.org/10.1080/1364557032000119616>
- Ashcroft, F. M., & Rorsman, P. (2012). Diabetes mellitus and the β cell: The last ten years. *Cell*, 148(6), 1160–1171. <https://doi.org/10.1016/j.cell.2012.02.010>
- Badan Pusat Statistik Provinsi NTT. (2019). Ringkasan pola konsumsi penduduk provinsi Nusa Tenggara Timur Maret 2019. Kupang: Badan Pusat Statistik NTT
- Badan Pusat Statistik Kabupaten Sumba Barat. (2024). *Kabupaten Sumba Barat dalam angka 2024*. BPS Kabupaten Sumba Barat. <https://sumbaratkab.bps.go.id>
- Badan Pusat Statistik Kabupaten Sumba Barat Daya. (2024). *Kabupaten Sumba Barat Daya dalam angka 2024*. BPS Kabupaten Sumba Barat Daya. <https://sumbaratdayakab.bps.go.id>
- Badan Pusat Statistik Kabupaten Sumba Tengah (2024). *Kabupaten Sumba Tengah dalam angka 2024*. BPS Kabupaten Sumba tengah. <https://sumbatengahkab.bps.go.id>
- Badan Pusat Statistik Kota Tomohon. (2024). *Kota Tomohon dalam angka 2024* (Vol. 17). BPS Kota Tomohon. <https://tomohonkota.bps.go.id>
- Balder, H. F., Virtanen, M., Brants, H. A. M., Krogh, V., Dixon, L. B., Tan, F., et al. (2003). Common and Country-Specific Dietary Patterns in Four European Cohort Studies. *J Nutr*, 133(12), 4246–4251. <https://doi.org/10.1093/jn/133.12.4246>
- Basu, D., & Bornfeldt, K. E. (2020). Hypertriglyceridemia and atherosclerosis: Using human research to guide mechanistic studies in animal models. *Frontiers in Endocrinology*, 11, Article 504. <https://doi.org/10.3389/fendo.2020.00504>
- Beckett, E. L., Jones, P. R., Veysey, M., & Lucock, M. (2017). Nutrigenetics—Personalized Nutrition in the Genetic Age. *Explor Res Hypothesis Med*, 2(4), 1–8. <https://doi.org/10.14218/erhm.2017.00027>
- Béghin, L., Dauchet, L., De Vriendt, T., Cuenca-García, M., Manios, Y., Toti, E., et al. (2014). Influence of parental socio-economic status on diet quality of European adolescents: Results from the HELENA study. *Br J Nutr*, 111(7), 1303–1312. <https://doi.org/10.1017/S0007114513003796>
- Belfiore, A., Malaguarnera, R., Vella, V., Lawrence, M. C., Sciacca, L., Frasca, F., Morrione, A., & Vigneri, R. (2017). Insulin receptor isoforms in physiology and disease: An updated view. *Endocr Rev*, 38(5), 379–431.

<https://doi.org/10.1210/er.2017-00073>

- Bellwood, P. (2004). The origins and dispersals of agricultural communities in Southeast Asia. In I. Glover & P. Bellwood (Eds.), *Southeast Asia: From Prehistory to History* (pp. 21–40). London, UK: Routledge Curzon. DOI/URL: <http://hdl.handle.net/1885/77599>
- Bellwood, P. (1976). *Archaeological research in Minahasa and the Talaud Islands, Northeastern Indonesia. Asian Perspectives, 19*(2), 240–288. University of Hawai'i Press. Retrieved from <http://hdl.handle.net/10125/19177>
- Bergholdt, H. K. M., Nordestgaard, B. G., & Ellervik, C. (2015). Milk intake is not associated with low risk of diabetes or overweight-obesity: A Mendelian randomization study in 97,811 Danish individuals. *Am J Clin Nutr, 102*(2), 487–496. <https://doi.org/10.3945/ajcn.114.105049>
- Bijelic, R., Balaban, J., Milicevic, S., & Sipka, S. U. (2020). The Association of Obesity and Microvascular Complications with Glycemic Control in Patients with Type 2 Diabetes Mellitus. *Med Arch (Sarajevo, Bosnia and Herzegovina), 74*(1), 14–18. <https://doi.org/10.5455/medarh.2020.74.14-18>
- Black S. A. (2002). Diabetes, diversity, and disparity: what do we do with the evidence?. *American journal of public health, 92*(4), 543–548. <https://doi.org/10.2105/ajph.92.4.543>
- Blüher, M., Malhotra, A., & Bader, G. (2022). Beta-cell function in treatment-naïve patients with type 2 diabetes mellitus.pdf. *Diabetes, Obesity and Metabolism, 25*, 1403–1407. <https://doi.org/https://doi.org/10.1111/dom.14969>
- Bonsembiante, L., Targher, G., & Maffei, C. (2021). *Type 2 Diabetes and Dietary Carbohydrate Intake of Adolescents and Young Adults : What Is the Impact of Different Choices ?* 1–18.
- Botto, L. D., & Khoury, M. J. (2001). Commentary: Facing the challenge of gene–environment interaction: The two-by-four table and beyond. *American Journal of Epidemiology, 153*(10), 1016–1020. <https://doi.org/10.1093/aje/153.10.1016>
- Boucher, J., Kleinridders, A., & Kahn, C. R. (2014). Insulin receptor signaling in normal. *Cold Spring Harbor Perspectives in Biology, 6*(1), a009191. <https://doi.org/10.1101/cshperspect.a009191>
- Brookes, A. J. (1999). The essence of SNPs. *Gene, 234*(2), 177–186. [https://doi.org/10.1016/S0378-1119\(99\)00219-X](https://doi.org/10.1016/S0378-1119(99)00219-X)
- Bull, F. C., Maslin, T. S., & Armstrong, T. (2009). Global physical activity questionnaire (GPAQ): nine country reliability and validity study. *Journal of physical activity & health, 6*(6), 790–804. <https://doi.org/10.1123/jpah.6.6.790>
- Burton, P. R., Clayton, D. G., Cardon, L. R., Craddock, N., Deloukas, P., Duncanson, A., Kwiatkowski, D. P., et al. (2007). Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature, 447*(7145), 661–678.

<https://doi.org/10.1038/nature05911>

- Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, B. P. (2003). Beta-Cell Deficit and Increased Beta-Cell Apoptosis in Humans With Type 2 Diabetes. *Diabetes*, 52(1), 102–110.
<https://doi.org/10.1103/PhysRev.102.1295>
- Camhi, S. M., Bray, G. A., Bouchard, C., Greenway, F. L., Johnson, W. D., Newton, R. L., et al. (2011). The relationship of waist circumference and BMI to visceral, subcutaneous, and totalbody fat: sex and race differences. *Obesity*, 19(2), 402–408. <https://doi.org/10.1038/oby.2010.248>.
- Carlström, M., & Larsson, S. C. (2018). Coffee consumption and reduced risk of developing type 2 diabetes: A systematic review with meta-analysis. *Nutr Rev*, 76(6), 395–417. <https://doi.org/10.1093/nutrit/nuy014>
- Cauchi, S., Proença, C., Choquet, H., Gaget, S., De Graeve, F., Marre, M., et al. (2008). Analysis of novel risk loci for type 2 diabetes in a general French population: The D.E.S.I.R. study. *J Mol Med*, 86(3), 341–348.
<https://doi.org/10.1007/s00109-007-0295-x>
- Cavagnoli, G., Pimentel, A. L., Freitas, P. A. C., Gross, J. L., & Camargo, J. L. (2017). Effect of ethnicity on HbA1c levels in individuals without diabetes: Systematic review and meta-analysis. *PLoS ONE*, 12(2), 1–14.
<https://doi.org/10.1371/journal.pone.0171315>
- Cerf, M. E. (2013). *Beta cell dysfunction and insulin resistance*. 4(March), 1–12.
<https://doi.org/10.3389/fendo.2013.00037>
- Chen, Z., Pereira, M. A., Seielstad, M., Koh, W. P., Tai, E. S., Teo, Y. Y., et al. (2014). Joint effects of known type 2 diabetes susceptibility loci in genome-wide association study of Singapore Chinese: The Singapore Chinese health study. *PLoS ONE*, 9(2). <https://doi.org/10.1371/journal.pone.0087762>
- Cheon, E. J., Cha, D. H., Cho, S. K., Noh, H., Park, S., Kang, S., Gee, H. Y., & Lee, S. (2018). Novel association between CDKAL1 and cholesterol efflux capacity: Replication after GWAS-based discovery. *Atherosclerosis*, 273, 21–27. <https://doi.org/10.1016/j.atherosclerosis.2018.04.011>
- Ching, Y. P., Pang, A. S. H., Lam, W. H., Qi, R. Z., & Wang, J. H. (2002). Identification of a neuronal Cdk5 activator-binding protein as Cdk5 inhibitor. *J Biol Chem*, 277(18), 15237–15240.
<https://doi.org/10.1074/jbc.C200032200>
- Chistiakov, D. A., Potapov, V. ., Smetanina, S. ., Bel'chikoca, L. ., Suplotova, L. ., & Nosikov, V.(2011). *The carriage of risk variants of CDKAL1 impairs beta-cell function in both diabetic and non-diabetic patients and reduces response to non-sulfonylurea and sulfonylurea agonists of the pancreatic KATP channel*. 227–235. <https://doi.org/10.1007/s00592-011-0299-4>
- Chiu, S.L., Cline, H.T. (2010). Insulin receptor signaling in the development of neuronal structure and function. *Neural Dev* 5, 7 .
<https://doi.org/10.1186/1749-8104-5-7>
- Choi, W. J., Jin, H. S., Kim, S. S., & Shin, D. (2020). Dietary protein and fat intake affects diabetes risk with cdkal1 genetic variants in Korean adults. *Int J Mol*

- Sci*, 21(16), 1–14. <https://doi.org/10.3390/ijms21165607>
- Christensen, A. A., & Gannon, M. (2019). The Beta Cell in Type 2 Diabetes. *Curr Diab Rep*, 19(9). <https://doi.org/10.1007/s11892-019-1196-4>
- Cleland et al. (2014). Validity of the Global Physical Activity Questionnaire (GPAQ) in assessing levels and change in moderate-vigorous physical activity and sedentary behaviour. *BMC Public Health*. 14,1-11.
- Clemente-Suarez, Vincente Javier, Beltran-Velasco, A. I., Redondo-flprez, L., Martin-Rodriguez, A., & Tornero-aguilera, J. F. (2023). *Global Impacts of Western Diet and Its Effects on Metabolism and Health : A Narrative Review*.
- Colosia, A. D., Palencia, R., & Khan, S. (2013). Prevalence of hypertension and obesity in patients with type 2 diabetes mellitus in observational studies: A systematic literature review. *Diabetes, Metab Syndr Obes: Targets and Therapy*, 6(October 2015), 327–338. <https://doi.org/10.2147/DMSO.S51325>
- Cox, M. P., Hudjashov, G., Sim, A., Savina, O., Karafet, T. M., Sudoyo, H., & Lansing, J. S. (2016). Small Traditional Human Communities Sustain Genomic Diversity over Microgeographic Scales despite Linguistic Isolation. *Mol. Biol. Evol.*, 33(9), 2273–2284. <https://doi.org/10.1093/molbev/msw099>
- Clemente-Suárez, V. J., Beltrán-Velasco, A. I., Redondo-Flórez, L., Martín-Rodríguez, A., & Tornero-Aguilera, J. F. (2023). Global impacts of Western diet and its effects on metabolism and health: A narrative review. *Nutrients*, 15(12), 2749. <https://doi.org/10.3390/nu15122749>
- Corbett, S., Courtiol, A., Lummaa, V., Moorad, J., & Stearns, S. (2018). The transition to modernity and chronic disease : mismatch and natural selection. *Nature Reviews Genetics*, 19(July), 419–430. <https://doi.org/10.1038/s41576-018-0012-3>
- Cordell, H. J. (2001). *Sample size requirements to control for stochastic variation in magnitude and location of allele-sharing linkage statistics in affected sibling pairs*. 491–502.
- Corella, D., Asensio, E. M., Coltell, O., Sorlí, J. V., Estruch, R., Martínez-González, M. Á., Salas-Salvadó, J., et al. (2016). CLOCK gene variation is associated with incidence of type-2 diabetes and cardiovascular diseases in type-2 diabetic subjects: Dietary modulation in the PREDIMED randomized trial. *Cardiovas Diabetol*, 15(1), 1–13. <https://doi.org/10.1186/s12933-015-0327-8>
- Corella, D., Carrasco, P., Sorlí, J. V., Estruch, R., Rico-Sanz, J., Martínez-González, M. Á., et al. (2013). Mediterranean diet reduces the adverse effect of the TCF7L2-rs7903146 polymorphism on cardiovascular risk factors and stroke incidence: A randomized controlled trial in a high-cardiovascular-risk population. *Diabetes Care*, 36(11), 3803–3811. <https://doi.org/10.2337/dc13-0955>
- Cornelis, M. C., Qi, L., Kraft, P., & Hu, F. B. (2009). TCF7L2, dietary carbohydrate, and risk of type 2 diabetes in US women. *Am J Clin Nutr*,

- 89(4), 1256–1262. <https://doi.org/10.3945/ajcn.2008.27058>
- Creatore, Maria Isabella, Moineddin, R., Booth, G., Manuel, D. H., DesMeules, M., McDermott, S., & Glazier, R. H. (2010). Age- and sex-related prevalence of diabetes mellitus among immigrants to Ontario, Canada. *Cmaj*, *182*(8), 781–789. <https://doi.org/10.1503/cmaj.091551>
- Dabelea, D., Hanson, R. L., Bennett, P. H., Roumain, J., Knowler, W. C., & Pettitt, D. J. (1998). Increasing prevalence of type II diabetes in American Indian children. *Diabetologia*, *41*(8), 904–910. <https://doi.org/10.1007/s001250051006>
- Danaei, G., Finucane, M. M., Lu, Y., Singh, G. M., Cowan, M. J., Paciorek, C. J., et al. (2011). National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: Systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *The Lancet*, *378*(9785), 31–40. [https://doi.org/10.1016/S0140-6736\(11\)60679-X](https://doi.org/10.1016/S0140-6736(11)60679-X)
- Darwish, R., Abdel-Hamid, E., Gohar, N., Soliman, A., & Naguib, M. (2020). Single nucleotide polymorphisms in CDKAL1 gene are not associated with risk of type 2 diabetes mellitus in a group of Egyptian population. *Biomedical and Pharmacology Journal*, *13*(2), 529–535. <https://doi.org/10.13005/bpj/1914>
- Deforest, N., & Majithia, A. R. (2022). Genetics of Type 2 Diabetes : Implications from Large - Scale Studies. *Current Diabetes Reports*, 227–235. <https://doi.org/10.1007/s11892-022-01462-3>
- Dehwah, M. A. S., Wang, M., & Huang, Q. Y. (2010). CDKAL1 and type 2 diabetes: A global meta-analysis. *Genet Mol Res*, *9*(2), 1109–1120. <https://doi.org/10.4238/vol9-2gmr802>
- Deurenberg-Yap, M., Chew, S. K., Lin, V. F. P., Tan, B. Y., Van Staveren, W. A., & Deurenberg, P. (2001). Relationships between indices of obesity and its comorbidities in multi-ethnic Singapore. *Int J Obes*, *25*(10), 1554–1562. <https://doi.org/10.1038/sj.ijo.0801739>
- Eckel, R. H., Kahn, S. E., Ferrannini, E., Goldfine, A. B., Nathan, D. M., Schwartz, M. W., Smith, R. J., & Smith, S. R. (2011). Obesity and type 2 diabetes: What Can be unified and what needs to be individualized? *Diabetes Care*, *34*(6), 1424–1430. <https://doi.org/10.2337/dc11-0447>
- Eckhardt, B. J., Holzman, R. S., Kwan, C. K., Baghdadi, J., & Aberg, J. A. (2012). Glycated hemoglobin A 1c as screening for diabetes mellitus in HIV-infected individuals. *AIDS Patient Care STDS*, *26*(4), 197–201. <https://doi.org/10.1089/apc.2011.0379>
- Erber, E., Hopping, B. N., Grandinetti, A., Park, S. Y., Kolonel, L. N., & Maskarinec, G. (2010). Dietary patterns and risk for diabetes: The multiethnic cohort. *Diabetes Care*, *33*(3), 532– 538. <https://doi.org/10.2337/dc09-1621>
- Ericson, U., Rukh, G., Stojkovic, I., Sonestedt, E., Gullberg, B., Wirfält, E., Wallström, P., & Orho-Melander, M. (2013). Sex-specific interactions between the IRS1 polymorphism and intakes of carbohydrates and fat on

- incident type 2 diabetes. *Am J Clin Nutr*, 97(1), 208–216.
<https://doi.org/10.3945/ajcn.112.046474>
- Ericson, U., Hindy, G., Drake, I., Schulz, C. A., Brunkwall, L., Hellstrand, S., Almgren, P., & Orho-Melander, M. (2018). Dietary and genetic risk scores and incidence of type 2 diabetes. *Genes and Nutrition*, 13(1), 1–14.
<https://doi.org/10.1186/s12263-018-0599-1>
- Estadella, D., Claudia, M., Oller, P., Oyama, L. M., Ribeiro, E. B., Dâmaso, A. R., & Piano, A. De. (2013). Lipotoxicity : Effects of Dietary Saturated and Transfatty Acids. 2013. <https://doi.org/10.1155/2013/137579>
- Fahriani, I., & Sari, V. S. (2023). Burial System In Karangkelang Island, Talud Regency, North Sulawesi. *Kalpataru*, 32(2), 149–162.
<https://doi.org/https://doi.org/10.55981/kpt.2023.748>
- Fang, M., Hu, Z. F., & Feng, L. J. (2020). Association between dietary pattern and the risk of type2 diabetes mellitus in Zhejiang Province, China: A case-control study. *Asia Pac J Clin Nutr*, 29(4), 821–826.
[https://doi.org/10.6133/apjn.202012_29\(4\).0018](https://doi.org/10.6133/apjn.202012_29(4).0018)
- Fardet, A. (2010). New hypotheses for the health-protective mechanisms of whole-grain cereals: What is beyond fibre? *Nutr Res Rev*, 23(1), 65–134.
<https://doi.org/10.1017/S0954422410000041>
- Fawwad, A., Rashid, K., & Basil, A. (2017). Genetics of Type 2 Diabetes Mellitus - Asian Perspective (A Review). *Metab, Endocrinol*, 2035, 52–59. <https://doi.org/10.4274/tjem.2017-56531>
- Fisher, E., Boeing, H., Fritsche, A., Doering, F., Joost, H. G., & Schulze, M. B. (2009). Whole- grain consumption and transcription factor-7-like 2 (TCF7L2) rs7903146: Gene-diet interaction in modulating type 2 diabetes risk. *Br J Nutr*, 101(4), 478–481
<https://doi.org/10.1017/S0007114508020369>
- Food and Agriculture Organization of the United Nations. (2010). *Fats and fatty acids in human nutrition: Report of an expert consultation* (FAO Food and Nutrition Paper No. 91). Rome: FAO.
- Forouzanfar, M. H., Afshin, A., Alexander, L. T., Biryukov, S., Brauer, M., Cercey, K., Charlson, F. J., Cohen, A. J., et al. (2016). Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*, 388(10053), 1659–1724. [https://doi.org/10.1016/S0140-6736\(16\)31679-8](https://doi.org/10.1016/S0140-6736(16)31679-8)
- Franks, P. W., Pearson, E., & Florez, J. C. (2013). Gene-environment and gene-treatment interactions in type 2 diabetes: Progress, pitfalls, and prospects. *Diabetes Care*, 36(5), 1413–1421. <https://doi.org/10.2337/dc12-2211>
- Fu, Z., R. Gilbert, E., & Liu, D. (2012). Regulation of Insulin Synthesis and Secretion and Pancreatic Beta-Cell Dysfunction in Diabetes. *Curr Diabetes Rev*, 9(1), 25–53. <https://doi.org/10.2174/15733998130104>
- Garrashy, A. A. M., Idris, O. F., Salimeen, M., & Eltirafi, I. (2021). Assessment

- of C-reactive protein, lipid profile and body mass index in Type 2 diabetes mellitus in Sudanese patients. *Archives of Internal Medicine Research*, 4, 125–133. <https://doi.org/10.26502/aimr.0063>
- Garcia-Bailo, B., El-Soheemy, A., Haddad, P. S., Arora, P., Benzaied, F., Karmali, M., & Badawi, A. (2011). Vitamins D, C, and E in the prevention of type 2 diabetes mellitus: Modulation of inflammation and oxidative stress. *Biologics: Targets and Therapy*, 5, 7–19. <https://doi.org/10.2147/BTT.S14417>
- Galicia-Garcia, U., Benito-Vicente, A., Jebari, S., Larrea-Sebal, A., Siddiqi, H., Uribe, K. B., Ostolaza, H., & Martín, C. (2020). Pathophysiology of type 2 diabetes mellitus. *International Journal of Molecular Sciences*, 21(17), 1–34. <https://doi.org/10.3390/ijms21176275>
- GBD 2017 DALYs and HALE Collaborators (2018). Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet (London, England)*, 392(10159), 1859–1922. [https://doi.org/10.1016/S0140-6736\(18\)32335-3](https://doi.org/10.1016/S0140-6736(18)32335-3)
- Geng, T., & Huang, T. (2020). Gene-environment interactions and type 2 diabetes. *Asia Pacific journal of clinical nutrition*, 29(2), 220–226. [https://doi.org/10.6133/apjcn.202007_29\(2\).0002](https://doi.org/10.6133/apjcn.202007_29(2).0002)
- Ghosh, C., Das, N., Saha, S., Kundu, T., Sircar, D., & Roy, P. (2022). Involvement of Cdkal1 in the etiology of type 2 diabetes mellitus and microvascular diabetic complications : a review. *Journal of Diabetes & Metabolic Disorders*, 991–1001. <https://doi.org/10.1007/s40200-021-00953-6>
- Grarup, N., Rose, C. S., Andersson, E. A., Andersen, G., Nielsen, A. L., Clausen, J. O., Rasmussen, S. S., Jørgensen, T., Sandbæk, A., Lauritzen, T., Schmitz, O., Hansen, T., & Pedersen, O. (2007). Studies of Association of Variants Near the HHEX, CDKN2A/B, and IGF2BP2 Genes With Type 2 Diabetes and Impaired Insulin Release in 10,705 Danish Subjects Validation and Extension of Genome-Wide Association Studies. *Diabetes*, 56(December), 3105–3111. <https://doi.org/10.2337/db07-0856.N.G>
- Groenewoud, M. J., Dekker, J. M., & Fritsche, A. (2008). Variants of CDKAL1 and IGF2BP2 affect first-phase insulin secretion during hyperglycaemic clamps. 1659–1663. <https://doi.org/10.1007/s00125-008-1083-z>
- Guariguata, L., Whiting, D. R., Hambleton, I., Beagley, J., Linnenkamp, U., & Shaw, J. E. (2014). Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Res Clin Pract* 103(2), 137–149. <https://doi.org/10.1016/j.diabres.2013.11.002>
- Guasch-Ferré, M., Becerra-Tomás, N., Ruiz-Canela, M., Corella, D., Schröder, H., Estruch, R., Ros, E., Arós, F., et al. (2017). Total and subtypes of dietary fat intake and risk of type 2 diabetes mellitus in the Prevención con Dieta Mediterránea (PREDIMED) study. *The American Journal of Clinical*

Nutrition, 105(3), 723–735. <https://doi.org/10.3945/ajcn.116.142034>
PubMedSpringer Medicine

- Halban, P. A. (1994). Proinsulin processing in the regulated and the constitutive secretory pathway. *Diabetologia*, 37 (2 Supplement).
<https://doi.org/10.1007/BF00400828>
- Halban, Philippe A., Polonsky, K. S., Bowden, D. W., Hawkins, M. A., Ling, C., Mather, K. J., et al. (2014). β -Cell failure in type 2 diabetes: Postulated mechanisms and prospects for prevention and treatment. *Diabetes Care*, 37(6), 1751–1758. <https://doi.org/10.2337/dc14-0396>
- Hamilton, M. T., Hamilton, D. G., & Zderic, T. W. (2014). Sedentary behavior as a mediator of type 2 diabetes. *Med Sport Sci*, 60(225), 11–26.
<https://doi.org/10.1159/000357332>
- Han, J. C., Rutledge, M. S., Kozlosky, M., Salatia, C. G., Gustafson, J. K., Keil, M. F., Fleisch, A. F., Roberts, M. D., Ning, C., & Yanovski, J. A. (2008). Insulin Resistance, Hyperinsulinemia, and Energy Intake in Overweight Children. *The Journal of Pediatrics*, 152, 612–617.
<https://doi.org/https://doi.org/10.1016/j.jpeds.2007.12.036>
- Han, X., Luo, Y., Ren, Q., Zhang, X., Wang, F., Sun, X., Zhou, X., & Ji, L. (2010). Implication of genetic variants near SLC30A8, HHEX, CDKAL1, CDKN2A/B, IGF2BP2, FTO, TCF2, KCNQ1, and WFS1 in Type 2 Diabetes in a Chinese population. *BMC Med Genet*, 11(1), 10–15.
<https://doi.org/10.1186/1471-2350-11-81>
- Hein, R., Beckmann, L., & Ã, J. C. (2008). *Sample Size Requirements for Indirect Association Studies of Gene – environment Interactions (G Â E)*. 245, 235–245. <https://doi.org/10.1002/gepi.20298>
- Hill, C., Soares, P., Mormina, M., Macaulay, V., Clarke, D., Blumbach, P. B., Vizuete-Forster, M., et al. (2007). A mitochondrial stratigraphy for Island Southeast Asia. *Am J Hum Genet*, 80(1), 29–43.
<https://doi.org/10.1086/510412>
- Hindy, G., Sonestedt, E., Ericson, U., Jing, X. J., Zhou, Y., Hansson, O., Renström, E., Wirfält, E., & Orho-Melander, M. (2012). Role of TCF7L2 risk variant and dietary fibre intake on incident type 2 diabetes. *Diabetologia*, 55(10), 2646–2654. <https://doi.org/10.1007/s00125-012-2634-x>
- Ho, F. K., Gray, S. R., Welsh, P., Gill, J. M. R., Sattar, N., Pell, J. P., & Morales, C. C. (2022). Ethnic differences in cardiovascular risk : examining differential exposure and susceptibility to risk factors. *BMC Medicine*, 1–10.
<https://doi.org/10.1186/s12916-022-02337-w>
- Houston, D. K., Nicklas, B. J., Ding, J., Harris, T. B., Tylavsky, F. A., Newman, A. B., ... & Kritchevsky, S. B. (2008). Dietary protein intake is associated with lean mass change in older, community-dwelling adults: The Health, Aging, and Body Composition (Health ABC) Study. *The American Journal of Clinical Nutrition*, 87(1), 150–155. <https://doi.org/10.1093/ajcn/87.1.150>
- Houttu, V., Grefhorst, A., Cohn, D. M., Levels, J. H. M., Lennep, J. R. Van,

- Stroes, E. S. G., Groen, A. K., & Tromp, T. R. (2023). Severe Dyslipidemia Mimicking Familial Hypercholesterolemia Induced by High-Fat, Low-Carbohydrate Diets: A Critical Review. *Nutrients*, *15*(4), 962. <https://doi.org/https://doi.org/10.3390/nu15040962>
- Huang, C. Y., Huang, H. L., Yang, K. C., Lee, L. T., Yang, W. S., Huang, K. C., & Tseng, F. Y. (2015). Serum Triglyceride Levels Independently Contribute to the Estimation of Visceral Fat Amount Among Nondiabetic. *94*(23), 1–7. <https://doi.org/10.1097/MD.0000000000000965>
- Huang, Y., Chen, Z., Chen, B., Li, J., Yuan, X., Li, J., Wang, W., Dai, T., Chen, H., Wang, Y., Wang, R., Wang, P., Guo, J., Dong, Q., Liu, C., Wei, Q., Cao, D., & Liu, L. (2023). Dietary sugar consumption and health: umbrella review. *BMJ (Clinical research ed.)*, *381*, e071609. <https://doi.org/10.1136/bmj-2022-071609>
- Hu, F. B. (2011). Globalization of diabetes: The role of diet, lifestyle, and genes. *Diabetes Care*, *34*(6), 1249–1257. <https://doi.org/10.2337/dc11-0442>
- Hu, F. B., Li, T. Y., Colditz, G. A., Willett, W. C., & Manson, J. A. E. (2003). Television Watching and Other Sedentary Behaviors in Relation to Risk of Obesity and Type 2 Diabetes Mellitus in Women. *JAMA*, *289*(14), 1785–1791. <https://doi.org/10.1001/jama.289.14.1785>
- Hulsegge, G., Spijkerman, A. M. W., Schouw, Y. T. Van Der, Bakker, S. J. L., Gansevoort, R. T., Smit, H. A., & Verschuren, W. M. M. (2017). Trajectories of metabolic risk factors and biochemical markers prior to the onset of type 2 diabetes : the population-based longitudinal Doetinchem study. November 2016, 1–8. <https://doi.org/10.1038/nutd.2017.23>
- Hudjashov, G., Karafet, T. M., Lawson, D. J., Downey, S., Savina, O., Sudoyo, H., Lansing, J. S., et al. (2017). Complex Patterns of Admixture across the Indonesian Archipelago. *Mol Biol Evol*, *34*(10), 2439–2452. <https://doi.org/10.1093/molbev/msx196>
- Huxley, R., Lee, C. M. Y., Barzi, F., Timmermeister, L., Czernichow, S., Perkovic, V., Grobbee, D. E., Batty, D., & Woodward, M. (2009). Coffee, decaffeinated coffee, and tea consumption in relation to incident type 2 diabetes mellitus: A systematic review with meta-analysis. *Arch Intern Med*, *169*(22), 2053–2063. <https://doi.org/10.1001/archinternmed.2009.439>
- Imamura, F., O'Connor, L., Ye, Z., Mursu, J., Hayashino, Y., Bhupathiraju, S. N., & Forouhi, N.G. (2015). Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: Systematic review, meta-analysis, and estimation of population attributable fraction. *BMJ (Online)*, *351*, 1–12. <https://doi.org/10.1136/bmj.h3576>
- Inaishi, J., & Saisho, Y. (2020). Beta-cell mass in obesity and type 2 diabetes, and its relation to pancreas fat: A mini-review. *Nutrients*, *12*(12), 1–16. <https://doi.org/10.3390/nu12123846>
- Inaishi, J., Saisho, Y., Sato, S., Kou, K., Murakami, R., Watanabe, Y., Kitago, M., Kitagawa, Y., Yamada, T., & Itoh, H. (2016). Effects of obesity and

- diabetes on α - and β -cell mass in surgically resected human pancreas. *J Clin Endocrinol Metab*, 101(7), 2863–2873. <https://doi.org/10.1210/jc.2016-1374>
- Ismail, L., Materwala, H., & Al Kaabi, J. (2021). Association of risk factors with type 2 diabetes: A systematic review. *Comput Struct Biotechnol J*, 19, 1759–1785. <https://doi.org/10.1016/j.csbj.2021.03.003>
- International Diabetes Federation. (2019). *IDF Diabetes Atlas* (7th ed.). International Diabetes Federation. <https://diabetesatlas.org>
- Izaola-jáuregui, O., Primo-martín, D., López, J. J., & Luis-román, D. A. De. (2021). Endocrinología, Diabetes y Nutrición The risk variant of CDKAL1 (rs7756992) impairs fasting glucose levels and insulin resistance improvements after a partial meal-replacement hypocaloric diet. *Endocrinología y Nutrición*, 68, 68. <https://doi.org/10.1016/j.endien.2020.08.017>
- James, B., Adrienne, L., & Peter, W. F. (2000). Parental transmission of type 2 diabetes: The Framingham Offspring Study. *Diabetes*, 49(12), 2201–2207. <https://doi.org/10.2337/diabetes.49.12.2201>
- Jannasch, F., Kröger, J., & Schulze, M. B. (2017). Dietary patterns and Type 2 diabetes: A systematic literature review and meta-analysis of prospective studies. *J Nutr*, 147(6), 1174–1182. <https://doi.org/10.3945/jn.116.242552>
- Jensen, M. D. (2008). Role of Body Fat Distribution and the Metabolic Complications of Obesity. *J Clin Endocrinol Metab*, 93(November), 57–63. <https://doi.org/10.1210/jc.2008-1585>
- Jing, Y., Han, G., Hu, Y., Bi, Y., Li, L., & Zhu, D. (2009). Tea consumption and risk of type 2 diabetes: A meta-analysis of cohort studies. *J Gen Intern Med*, 24(5), 557–562. <https://doi.org/10.1007/s11606-009-0929-5>
- Joseph, G.H. (2017). Analisis pola konsumsi pangan di propinsi Sulawesi Utara. *Jurnal Ilmiah Sains* (17):2, 178-188
- Jung, U. J., & Choi, M. S. (2014). Obesity and its metabolic complications: The role of adipokines and the relationship between obesity, inflammation, insulin resistance, dyslipidemia and nonalcoholic fatty liver disease. *Int J Mol Sci* (Vol. 15, Issue 4, pp. 6184–6223). <https://doi.org/10.3390/ijms15046184>
- Jung, S. H., Ha, K. H., & Kim, D. J. (2016). Visceral Fat Mass Has Stronger Associations with Diabetes and Prediabetes than Other Anthropometric Obesity Indicators among Korean Adults. *Yonsei Med J*, 57(3), 674–680. <https://doi.org/http://dx.doi.org/10.3349/ymj.2016.57.3.674>
- Jyothi, K. U., & Reddy, B. M. (2015). Gene – gene and gene – environment interactions in the etiology of type 2 diabetes mellitus in the population of Hyderabad, India. *Gene – gene and gene – environment interactions in the etiology of type 2 diabetes mellitus in the population of Hyderabad*, November. <https://doi.org/10.1016/j.mgene.2015.05.001>
- Kalyani, R. R., Corriere, M., & Ferrucci, L. (2014). Age-related and disease-related muscle loss: the effect of diabetes, obesity, and other diseases. *Lancet Diabetes Endocrinol.*, 2(10), 819–829. <https://doi.org/10.1016/S2213->

8587(14)70034-8.

- Kandou, G.D. (2009). Kebiasaan makan makanan etnik minahasa di provinsi Sulawesi Utara. *Jurnal Kesmas Andalas*, 3(2), 53-57.
<https://doi.org/10.24893/jkma.v3i2.59>
- Kane, J. P., Pullinger, C. R., Goldfine, I. D., & Malloy, M. J. (2021). Dyslipidemia and diabetes mellitus: Role of lipoprotein species and interrelated pathways of lipid metabolism in diabetes mellitus. *Current opinion in pharmacology*, 61, 21–27.
<https://doi.org/10.1016/j.coph.2021.08.013>
- Kaneto, H. (2015). Pathophysiology of type 2 diabetes mellitus. *Nihon Rinsho. Japanese Journal of Clinical Medicine*, 73(12), 2003–2007.
<https://doi.org/10.1093/med/9780199235292.003.1336>
- Kang, J., Guan, R. C., Zhao, Y., & Chen, Y. (2020). Obesity-related loci in TMEM18, CDKAL1 and FAIM2 are associated with obesity and type 2 diabetes in Chinese Han patients. *BMC medical genetics*, 21(1), 65.
<https://doi.org/10.1186/s12881-020-00999-y>
- Kanoni, S., Nettleton, J. A., Hivert, M. F., Ye, Z., Van Rooij, F. J. A., Shungin, D., Sonestedt, E., Ngwa, J. S., et al. (2011). Total zinc intake may modify the glucose-raising effect of a zinc transporter (SLC30A8) variant: A 14-cohort meta-analysis. *Diabetes*, 60(9), 2407–2416.
<https://doi.org/10.2337/db11-0176>
- Karafet, T. M., Hallmark, B., Cox, M. P., Sudoyo, H., Downey, S., Lansing, J. S., & Hammer, M.F. (2010). Major east-west division underlies y chromosome stratification across Indonesia. *Mol Biol Evol*, 27(8), 1833–1844.
<https://doi.org/10.1093/molbev/msq063>
- Kashyap, S., & Defronzo, R. A. (2007). The insulin resistance syndrome : physiological considerations. *Diabetes & Vascular Disease Research*, 4(1), 13–19. <https://doi.org/https://doi.org/10.3132/dvdr.2007.001>
- Katzmarzyk, P. T., Bray, G. A., Greenway, F. L., Johnson, W. D., Newton, R. L., Ravussin, E., Ryan, D. H., et al. (2010). Racial differences in abdominal depot-specific adiposity in white and African American adults. *Am J Clin Nutr*, 91(1), 7–15. <https://doi.org/10.3945/ajcn.2009.28136>
- Kautzky-Willer, A., Harreiter, J., & Pacini, G. (2016). Sex and Gender Differences in Risk, Pathophysiology and Complications of Type 2 Diabetes Mellitus. *Endocrine reviews*, 37(3), 278–316.
<https://doi.org/10.1210/er.2015-1137>
- Kemenkes RI. (2007). Riset Kesehatan Dasar, RISKESDAS. Jakarta: Balitbang Kemenkes RI
- Kemenkes RI. (2013). Riset Kesehatan Dasar, RISKESDAS. Jakarta: Balitbang Kemenkes RI
- Kemenkes RI. (2019). Riset Kesehatan Dasar, RISKESDAS. Jakarta: Balitbang Kemenkes RI
- Kemenkes RI. (2013). Risesdas dalam Angka Provinsi Nusa Tenggara Timur. Jakarta: Balitbang Kemenkes RI

- Kementerian Kesehatan RI. (2013). *Riskesmas dalam Angka Provinsi Sulawesi Utara*. Jakarta: Balitbang Kemenkes RI
- Kementerian Kesehatan Republik Indonesia. (2017). *Pedoman dan standar etik penelitian dan pengembangan kesehatan nasional*. Komite Etik Penelitian dan Pengembangan Kesehatan Nasional (KEPPKN)
- Kido, Y., Nakae, J., & Accili, D. (2001). Clinical review 125: The insulin receptor and its cell targets. *J Clin Endocrinol Metab*, 86(3), 972–979. <https://doi.org/10.1210/jc.86.3.972>
- Kim, A. N., Cho, H. J., Youn, J., Jin, T., Kang, M., Sung, J., & Lee, J. E. (2020). Coffee consumption, genetic polymorphisms, and the risk of type 2 diabetes mellitus: A pooled analysis of four prospective cohort studies. *Int J Environ Res Public Health*, 17(15), 1–18. <https://doi.org/10.3390/ijerph17155379>
- Kim, C., Bullard, K. M. K., Herman, W. H., & Beckles, G. L. (2010). Association between iron deficiency and A1C levels among adults without diabetes in the National Health And Nutrition Examination Survey, 1999-2006. *Diabetes Care*, 33(4), 780–785. <https://doi.org/10.2337/dc09-0836>
- Kim, Y., Keogh, J. B., & Clifton, P. M. (2016). Consumption of red and processed meat and refined grains for 4 weeks decreases insulin sensitivity in insulin-resistant adults: A randomized crossover study. *Metabolism Clinical and Experimental*, 68, 173–183. <https://doi.org/10.1016/j.metabol.2016.12.011>
- Kirchhoff, K., Machicao, F., Haupt, A., & Schäfer, S. A. (2008). *Polymorphisms in the TCF7L2, CDKAL1 and SLC30A8 genes are associated with impaired proinsulin conversion*. 597–601. <https://doi.org/10.1007/s00125-008-0926-y>
- Kleden, D. (2017). Belis dan Harga Seorang Perempuan Sumba (Perkawinan Adat Suku Wewewa, Sumba Barat Daya, NTT). *Jurnal Studi Budaya Nusantara*, 1(1), 23–34.
- Kodama, S., Fujihara, K., Ishiguro, H., Horikawa, C., Ohara, N., Yachi, Y., Tanaka, S., et al. (2018). Quantitative Relationship Between Cumulative Risk Alleles Based on Genome-Wide Association Studies and Type 2 Diabetes Mellitus: A Systematic Review and Meta-analysis. *J Epidemiol*, 28(1), 3–18. <https://doi.org/10.2188/jea.JE20160151>
- Kolb, H., Martin, S., & Kempf, K. (2021). Coffee and Lower Risk of Type 2 Diabetes: Arguments for a Causal Relationship. *Nutrients*, 13(4), 1144. <https://doi.org/10.3390/nu13041144>
- Kooner, J. S., Saleheen, D., Sim, X., Sehmi, J., Zhang, W., Frossard, P., Been, L. F., et al (2011). Genome-wide association study in individuals of South Asian ancestry identifies six new type 2 diabetes susceptibility loci. *Nat Genet*, 43(10), 984–989. <https://doi.org/10.1038/ng.921>
- Kompas.com (2015, 23 November). Jejak pembauran Melanesia dan Austronesia. Diakses pada 6 Oktober 2020 dari <https://sains.kompas.com>
- Laksono, P.M., Anantasari, E., Nandiswara O.A. (2019). Reproduksi moda (pertukaran) pangan: menyemai daulat hidup di Sumba Barat (Daya). *Jurnal Masyarakat dan Budaya*, 21 (3), 341-353.

<https://doi.org/10.14203/jmb.v21i3.747%20>

- Lamri, A., Abi Khalil, C., Jaziri, R., Velho, G., Lantieri, O., Vol, S., Froguel, P., Balkau, B., Marre, M., & Fumeron, F. (2012). Dietary fat intake and polymorphisms at the PPARG locus modulate BMI and type 2 diabetes risk in the D.E.S.I.R. prospective study. *Int J Obes*, *36*(2), 218–224. <https://doi.org/10.1038/ijo.2011.91>
- Lansing, J. S., Cox, M. P., Downey, S. S., Gabler, B. M., Hallmark, B., Karafet, T. M., et al. (2007). Coevolution of languages and genes on the island of Sumba, eastern Indonesia. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(41), 16022–16026. <https://doi.org/10.1073/pnas.0704451104>
- Layman, D. K., Evans, E. M., Erickson, D., Seyler, J., Weber, J., Bagshaw, D., Griel, A., Psota, T., & Kris-etherton, P. (2009). A Moderate-Protein Diet Produces Sustained Weight Loss and Long-Term Changes in Body Composition and Blood Lipids in. *Journal of Nutrition*, *139*(3), 514–521. <https://doi.org/10.3945/jn.108.099440>
- Lea, A. J., Clark, A. G., Dahl, A. W., Devinsky, O., Garcia, R., Golden, C. D., Kamau, J., Kraft, T. S., Lim, Y. A. L., Perry, G. H., Martins, D. J., & Mogoi, D. (2023). *Applying an evolutionary mismatch framework to understand disease susceptibility*. 1–19. <https://doi.org/10.1371/journal.pbio.3002311>
- Lee, H. A., Park, H., & Hong, Y. S. (2022). *Sex Differences in the Effects of CDKAL1 Variants on Glycemic Control in Diabetic Patients : Findings from the Korean Genome and Epidemiology Study*. 879–889.
- Lee, J. K., Kim, K., Ahn, Y., Yang, M., & Lee, J. E. (2015). Habitual coffee intake, genetic polymorphisms, and type 2 diabetes. *Eur J Endocrinol* *172*(5), 595–601. <https://doi.org/10.1530/EJE-14-0805>
- Lee, Y. H., Kang, E. S., Kim, S. H., Han, S. J., Kim, C. H., Kim, H. J., Ahn, C. W., et al. (2008). Association between polymorphisms in SLC30A8, HHEX, CDKN2A/B, IGF2BP2, FTO, WFS1, CDKAL1, KCNQ1 and type 2 diabetes in the Korean population. *J Hum Genet*, *53*(11–12), 991–998. <https://doi.org/10.1007/s10038-008-0341-8>
- Leja, D., & Manolio, T. (2011). Genetic basis of type 1 diabetes, type 2 diabetes, obesity, and their complications. *Advances and Emerging Opportunities in Diabetes Research: A Strategic Planning Report of the DMICC*, 44–67. [http://www.niddk.nih.gov/about-niddk/strategic-plans-reports/Documents/2011-02_Advances and Emerging Opportunities in Diabetes Research_A Strategic Planning Report of the Diabetes Mellitus Inte/DSP2011_03_GeneticBasis_508.pdf](http://www.niddk.nih.gov/about-niddk/strategic-plans-reports/Documents/2011-02_Advances%20and%20Emerging%20Opportunities%20in%20Diabetes%20Research_A%20Strategic%20Planning%20Report%20of%20the%20Diabetes%20Mellitus%20Inte/DSP2011_03_GeneticBasis_508.pdf)
- Ley, S. H., Hamdy, O., Mohan, V., & Hu, F. B. (2014). Prevention and management of type 2 diabetes: Dietary components and nutritional strategies. *The Lancet*, *383*(9933), 1999–2007. [https://doi.org/10.1016/S0140-6736\(14\)60613-9](https://doi.org/10.1016/S0140-6736(14)60613-9)
- Li, S. X. (2017). *The interplay between genes and dietary factors in the aetiology of Type 2 diabetes mellitus* [Doctoral dissertation, University of

Cambridge]. University of Cambridge Repository.

<https://www.repository.cam.ac.uk/handle/1810/275094>

- Li, S., Xiao, J., Ji, L., Weng, J., Jia, W., Lu, J., Guo, X., Liu, J., Shan, Z., Zhu, D., Chen, L., Zhao, Z., Ji, Q., Ge, J., Li, Q., Lin, L., Yang, Z., He, J., Yang, W., ... Hospital, F. P. (2014). BMI and waist circumference are associated with impaired glucose metabolism and type 2 diabetes in normal weight Chinese adults. *J Diabetes Complications*, 28(4), 470–476. <https://doi.org/10.1016/j.jdiacomp.2014.03.015>.
- Li, Y. Y., Wang, L. S., Lu, X. Z., Yang, Z. J., Wang, X. M., Zhou, C. W., Xu, J., Qian, Y., & Chen, A. L. (2013). CDKAL1 gene rs7756992 A/G polymorphism and type 2 diabetes mellitus: A meta-analysis of 62,567 subjects. *Sci Rep*, 3, 1–8. <https://doi.org/10.1038/srep03131>
- Lin, C. Y., Li, J. B., Wu, F., Wang, J. J., An, H. H., Qiu, H. N., Xia, L. F., & Li, Y. S. (2023). Comparison of lipid accumulation product and visceral adiposity index with traditional obesity indices in early - onset type 2 diabetes prediction : a cross - sectional study. *Diabetology & Metabolic Syndrome*, 15(111), 1–12. <https://doi.org/10.1186/s13098-023-01056-3>
- Lipoeto, N. I., Geok Lin, K., & Angeles-Agdeppa, I. (2013). Food consumption patterns and nutrition transition in South-East Asia. *Public Health Nutr*, 16(9), 1637–1643. <https://doi.org/10.1017/S1368980012004569>
- Lipson, M., Loh, P., Patterson, N., Moorjani, P., Ko, Y., Stoneking, M., Berger, B., & Reich, D. (2014). Reconstructing Austronesian population history in Island Southeast Asia. *Nature Communications*, 5, 1–7. <https://doi.org/10.1038/ncomms5689>
- Liu, S., Manson, J. E., Stamfer, M. J., Hu, F. B., Giovannucci, E., Colditz, G. A., Hennekens, C. H., & Willett, W. C. (2000). A prospective study of whole-grain intake and risk of type 2 diabetes mellitus in US women. *Am J Public Health*, 90(9), 1409–1415. <https://doi.org/10.2105/AJPH.90.9.1409>
- Liu, X., Liang, J., Geng, H., Xu, W., Teng, F., & Yang, M. (2020). Association of the CDKAL1 polymorphism rs10946398 with type 2 diabetes mellitus in adults: A meta-analysis. *Medicine*, 99(30), e21383. <https://doi.org/10.1097/MD.00000000000021383>
- Luthfiya, L., Pibriyanti, K., Nabawiyah, H., Fathimah, F., Ummah, S. K., & Luthfiya, L. (2024). Faktor-Faktor yang Berpengaruh terhadap Lingkar Perut pada Remaja Putri Factors Affecting Abdominal Circumference in Adolescent Girls. *Amerta Nutrition*, 8(1), 74–81. <https://doi.org/10.20473/amnt.v8i1.2024.74-81>
- Ma, Q., Liu, H., Xiang, G., Shan, W., & Xing, W. (2016). Association between glycated hemoglobin A1c levels with age and gender in Chinese adults with no prior diagnosis of diabetes mellitus. *Biomed Rep*, 4(6), 737–740. <https://doi.org/10.3892/br.2016.643>
- Ma, R. C. W., & Chan, J. C. N. (2013). Type 2 diabetes in East Asians: Similarities and differences with populations in Europe and the United States. *Ann N Y Acad Sci*, 1281(1), 64–91. <https://doi.org/10.1111/nyas.12098>
- Mahajan, A., Wessel, J., Willems, S. M., Zhao, W., Robertson, N. R., Chu, A. Y.,

- Gan, W., Kitajima, H., Taliun, D., Rayner, N. W., Guo, X., Lu, Y., Li, M., Jensen, R. A., Hu, Y., Huo, S., Lohman, K. K., Zhang, W., Cook, J. P., Prins, B. P., ... McCarthy, M. I. (2018). Refining the accuracy of validated target identification through coding variant fine-mapping in type 2 diabetes. *Nature genetics*, *50*(4), 559–571. <https://doi.org/10.1038/s41588-018-0084-1>
- Maliszewska, K., Adamska-patruno, E., Goscik, J., Lipinska, D., Citko, A., Krahel, A., Miniewska, K., Fiedorczuk, J., Moroz, M., Gorska, M., & Kretowski, A. (2019). *The Role of Muscle Decline in Type 2 Diabetes Development : A 5-Year Prospective Observational*.
- Martín-Campos, J. M. (2021). Genetic Determinants of Plasma Low-Density Lipoprotein Cholesterol Levels: Monogenicity, Polygenicity, and “Missing” Heritability. *Biomedicines*, *9*(11), 1728. <https://doi.org/10.3390/biomedicines9111728>
- McCarthy, M. I., & Hattersley, A. T. (2008). Learning from molecular genetics: novel insights arising from the definition of genes for monogenic and type 2 diabetes. *Diabetes*, *57*(11), 2889–2898. <https://doi.org/10.2337/db08-0343>
- Medici, F., Hawa, M., Ianari, A., Pyke, D. A., & Leslie, R. D. G. (1999). Concordance rate for type II diabetes mellitus in monozygotic twins: Actuarial analysis. *Diabetologia*, *42*(2), 146–150. <https://doi.org/10.1007/s001250051132>
- Meilani, J., Kandou, G. D., Ottay, R. I., & Kaunang, W. P. J. (2014). *Gambaran kebiasaan makan makanan beresiko penyakit jantung koroner pada etnis Minahasa di Kecamatan Tomohon Utara Kelurahan Kakaskasen 1*. *Jurnal Kedokteran Komunitas dan Tropik*, *2*(2).
- Merino, J., Leong, A., Liu, C., Porneala, B., Walford, G. A., & Grotthuss, M. Von. (2018). Metabolomics insights into early type 2 diabetes pathogenesis and detection in individuals with normal fasting glucose. *Diabetologia*, *61*, 1315–1324. <https://doi.org/https://doi.org/10.1007/s00125-018-4599-x>
- Mihardja, L., Soetrisno, U., & Soegondo, S. (2014). Prevalence and clinical profile of diabetes mellitus in productive aged urban Indonesians. *J Diabetes Investig*, *5*(5), 507–512. <https://doi.org/10.1111/jdi.12177>
- Miyaki, K., Oo, T., Song, Y., Lwin, H., Tomita, Y., Hoshino, H., Suzuki, N., & Muramatsu, M. (2010). Association of a cyclin-dependent kinase 5 regulatory subunit-associated protein 1- like 1 (CDKAL1) polymorphism with elevated hemoglobin A1c levels and the prevalence of metabolic syndrome in Japanese men: Interaction with dietary energy intake. *Am J Epidemiol*, *172*(9), 985–991. <https://doi.org/10.1093/aje/kwq281>
- Montonen, J., Järvinen, R., Heliövaara, M., Reunanen, A., Aromaa, A., & Knekt, P. (2005). Food consumption and the incidence of type II diabetes mellitus. *Eur J Clin Nutr*, *59*(3), 441–448. <https://doi.org/10.1038/sj.ejcn.1602094>
- Montonen, J., Knekt, P., Järvinen, R., & Reunanen, A. (2004). Dietary Antioxidant Intake and Risk of Type 2 Diabetes. *Diabetes Care*, *27*(2), 362–366. <https://doi.org/10.2337/diacare.27.2.362>

- Murea, M., Ma, L., & Freedman, B. I. (2012). Genetic and environmental factors associated with type 2 diabetes and diabetic vascular complications. *Rev Diabet Stud*, 9(1), 6–22. <https://doi.org/10.1900/RDS.2012.9.6>
- Nainggolan, O., Leli, I., & Julinati, P. (2018). Kebugaran jasmani menurut instrument GPAQ dibandingkan dengan V02max pada Wanita umur 25 sampai 54 tahun. *Buletin Penelitian Sistem Kesehatan*, 21(4), 271-280
- Nair, K. S., Bigelow, M. L., Asmann, Y. W., Chow, L. S., Coenen-Schimke, J. M., Klaus, K. A., Guo, Z. K., Sreekumar, R., & Irving, B. A. (2008). Asian indians have enhanced skeletal muscle mitochondrial capacity to produce ATP in association with severe insulin resistance. *Diabetes*, 57(5), 1166–1175. <https://doi.org/10.2337/db07-1556>
- Nakanishi, N., Suzuki, K., & Tatara, K. (2003). Alcohol consumption and risk for development of impaired fasting glucose or type 2 diabetes in middle-aged Japanese men. *Diabetes care*, 26(1), 48–54. <https://doi.org/10.2337/diacare.26.1.48>
- Naser, F.H., Fadheel, H.K., Hussain M.K., et al. (2021). Association of CDKAL1 gene polymorphisms with type 2 diabetes mellitus in a sample of Iraqi population. *Gene Reports*, 25. <https://doi.org/10.1016/j.genrep.2021.101371>
- Natri, H. M., Bobowik, K. S., Kusuma, P., Darusallam, C. C., Jacobs, G. S., Hudjashov, G., Stephen Lansing, J., Sudoyo, H., et al. (2020). Genome-wide DNA methylation and gene expression patterns reflect genetic ancestry and environmental differences across the Indonesian archipelago. *PloS Genetics*, 16(5), 1-21 <https://doi.org/10.1371/journal.pgen.1008749>
- Neel, J. V. (1962). *Diabetes Mellitus : A " Thrifty " Genotype Rendered Detrimental by " Progress " ? 1949*, 353–362.
- Nettleton, J. A., McKeown, N. M., Kanoni, S., Lemaitre, R. N., Hivert, M. F., Ngwa, J., Van Rooij, F. J. A., et al. (2010). Interactions of dietary whole-grain intake with fasting glucose- and insulin-related genetic loci in individuals of European descent: A meta-analysis of 14 cohort studies. *Diabetes Care*, 33(12), 2684–2691. <https://doi.org/10.2337/dc10-1150>
- Neuenschwander, M., Ballon, A., Weber, K. S., Norat, T., Aune, D., Schwingshackl, L., & Schlesinger, S. (2019). Role of diet in type 2 diabetes incidence: Umbrella review of meta-analyses of prospective observational studies. *The BMJ*, 366. <https://doi.org/10.1136/bmj.l2368>
- Nfor, O. N., Wu, M. F., Lee, C. Te, Wang, L., Liu, W. H., Tantoh, D. M., Hsu, S. Y., Lee, K. J., Ho, C. C., Debnath, T., Lung, C. C., & Liaw, Y. P. (2018). Body mass index modulates the association between CDKAL1 rs10946398 variant and type 2 diabetes among Taiwanese women. *Scientific Reports*, 8(1), 1–8. <https://doi.org/10.1038/s41598-018-31415-4>
- Nishida, Y., Iyadomi, M., Higaki, Y., & Tanaka, H. (2011). Influence of Physical Activity Intensity and Aerobic Fitness on the Anthropometric Index and Serum Uric Acid Concentration in People with Obesity. *Internal Medicine*, 50, 2121–2128. <https://doi.org/10.2169/internalmedicine.50.5506>
- Ho, C. C., et al. (2018). Body mass index modulates the association between

- CDKAL1 rs10946398 variant and type 2 diabetes among Taiwanese women. *Sci Rep*, 8(1), 1–8. <https://doi.org/10.1038/s41598-018-31415-4>
- Ng, M. C. Y., Park, K. S., Oh, B., Tam, C. H. T., Cho, Y. M., Shin, H. D., et al. (2008). Implication of genetic variants near TCF7L2, SLC30A8, HHEX, CDKAL1, CDKN2A/B, IGF2BP2, and FTO in type 2 diabetes and obesity in 6,719 Asians. *Diabetes*, 57(8), 2226–2233. <https://doi.org/10.2337/db07-1583>
- Okada, Y., Kubo, M., Ohmiya, H., Takahashi, A., Kumasaka, N., Hosono, N., Maeda, S., Wen, W., Dorajoo, R., Go, M. J., Zheng, W., Kato, N., Wu, J. Y., Lu, Q., GIANT consortium, Tsunoda, T., Yamamoto, K., Nakamura, Y., Kamatani, N., & Tanaka, T. (2012). Common variants at CDKAL1 and KLF9 are associated with body mass index in east Asian populations. *Nature genetics*, 44(3), 302–306. <https://doi.org/10.1038/ng.1086>
- Okamura, T., Yanobu-Takanashi, R., Takeuchi, F., Isono, M., Akiyama, K., Shimizu, Y., et al. (2012). Deletion of CDKAL1 Affects High-Fat Diet-Induced Fat Accumulation and Glucose-Stimulated Insulin Secretion in Mice, Indicating Relevance to Diabetes. *PLoS ONE*, 7(11). <https://doi.org/10.1371/journal.pone.0049055>
- Ortega-Azorín, C., Sorlí, J. V., Asensio, E. M., Coltell, O., Martínez-González, M. T., Salas-Salvadó, J., Covas, M. I., et al. (2012). Associations of the FTO rs9939609 and the MC4R rs17782313 polymorphisms with type 2 diabetes are modulated by diet, being higher when adherence to the Mediterranean diet pattern is low. *Cardiovasc Diabetol* (Vol. 11). <https://doi.org/10.1186/1475-2840-11-137>
- Paniagua, J. A., Paniagua, J. A., & Resistance, I. (2016). *Nutrition, insulin resistance and dysfunctional adipose tissue determine the different components of metabolic syndrome*. 7(19), 483–514. <https://doi.org/10.4239/wjd.v7.i19.483>
- Papamichou, D., Panagiotakos, D. B., & Itsiopoulos, C. (2019). Dietary patterns and management of type 2 diabetes: A systematic review of randomised clinical trials. *Metab Cardiovasc Dis*, 29(6), 531–543. <https://doi.org/10.1016/j.numecd.2019.02.004>
- Parhofer K. G. (2015). Interaction between Glucose and Lipid Metabolism: More than Diabetic Dyslipidemia. *Diabetes & metabolism journal*, 39(5), 353–362. <https://doi.org/10.4093/dmj.2015.39.5.353>
- Pascoe, L., Tura, A., Patel, S. K., Ibrahim, I. M., Ferrannini, E., Zeggini, E., Weedon, M. N., Mari, A., Hattersley, A. T., McCarthy, M. I., Frayling, T. M., Walker, M., Consortium, R., & Diabetes, U. K. T. (2007). *Common Variants of the Novel Type 2 Diabetes Genes CDKAL1 and HHEX/IDE Are Associated With Decreased Pancreatic β -Cell Function*. 56(December), 3101–3104. <https://doi.org/10.2337/db07-0634>. Additional
- Pecioska, S., Zillikens, M. C., Henneman, P., Snijders, P. J., Oostra, B. A., Cornelia, M., Duijn, V., & Aulchenko, Y. S. (2010). *Association between*

Type 2 Diabetes Loci and Measures of Fatness. 5(1), 1–5.
<https://doi.org/10.1371/journal.pone.0008541>

- Pepa, D.G., Vetrani, C., Vitale, M., & Riccardi, G. (2018). Wholegrain intake and risk of type 2 diabetes: Evidence from epidemiological and intervention studies. *Nutrients*, 10(9). <https://doi.org/10.3390/nu10091288>
- PERKENI. (2019). Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 Dewasa di Indonesia 2019. *Perkumpulan Endokrinologi Indonesia*, 1–117.
- Pérez, M.R., & Medina-Gómez, G. (2011). Obesity, adipogenesis and insulin resistance. *Endocrinol Nutr (English Edition)*, 58(7), 360–369. <https://doi.org/10.1016/j.endoen.2011.05.004>
- Petersen, M. C., & Shulman, G. I. (2018). Mechanisms of insulin action and insulin resistance. *Physiol Rev*, 98(4), 2133–2223.
<https://doi.org/10.1152/physrev.00063.2017>
- Pham, T. M., Carpenter, J. R., Morris, T. P., Sharma, M., & Petersen, I. (2019). Ethnic differences in the prevalence of type 2 diabetes diagnoses in the uk: Cross-sectional analysis of the healthimprovement network primary care database. *Clin Epidemiol*, 11, 1081–1088.
<https://doi.org/10.2147/CLEP.S227621>
- Picón, M. J., Murri, M., Muñoz, A., Fernández-García, J. C., Gomez-Huelgas, R., & Tinahones, F. J. (2012). Hemoglobin A 1c versus oral glucose tolerance test in postpartum diabetes screening. *Diabetes Care*, 35(8), 1648–1653. <https://doi.org/10.2337/dc11-2111>
- Plengvidhya, N., Chanprasert, C., Chongjaroen, N., Yenchitsomanus, P. thai, Homsanit, M., & Tangjittipokin, W. (2018). *Impact of KCNQ1, CDKN2A/2B, CDKAL1, HHEX, MTNR1B, SLC30A8, TCF7L2, and UBE2E2 on risk of developing type 2 diabetes in Thai population*. *BMC Med Genet*, 19(1), 1–9. <https://doi.org/10.1186/s12881-018-0614-9>
- Popkin, B. M. (2006). Global nutrition dynamics: The world is shifting rapidly toward a diet linked with noncommunicable diseases. *Am J Clin Nutr*, 84(2), 289–298. <https://doi.org/10.1093/ajcn/84.2.289>
- Popkin, B. M., Adair, L. S., & Ng, S. W. (2013). Now and Then: The Global Nutrition Transition: The Pandemic of Obesity in Developing Countries. 70(1), 3–21. <https://doi.org/10.1111/j.1753-4887.2011.00456.x>
- Pravednikova, A. E., Shevchenko, S. Y., Kerchev, V. V., Skhirtladze, M. R., Larina, S. N., Kachaev, Z. M., Egorov, A. D., & Shidlovskii, Y. V. (2020). Association of uncoupling protein (Ucp) gene polymorphisms with cardiometabolic diseases. *Mol Med*, 26(1). <https://doi.org/10.1186/s10020-020-00180-4>
- Qi, Q., X, W., G, S., & T, W. (2015). Genetic Determinants of Type 2 Diabetes in Asians. *International Journal of Diabetology & Vascular Disease Research*, March 2015, 1–9. <https://doi.org/10.19070/2328-353x-si01001>
- Qi, L., Cornelis, M. C., Zhang, C., Van Dam, R. M., & Hu, F. B. (2009). Genetic predisposition, Western dietary pattern, and the risk of type 2 diabetes in

- men. *Am J Clin Nutr*, 89(5), 1453–1458.
<https://doi.org/10.3945/ajcn.2008.27249>
- Qi, Q., & Hu, F. B. (2012). Genetics of type 2 diabetes in European populations. *J Diabetes*, 4(3), 203–212. <https://doi.org/10.1111/j.1753-0407.2012.00224.x>
- Racette, S. B., Evans, E. M., Weiss, E. P., Hagberg, J. M., & Holloszy, J. O. (2006). Abdominal adiposity is a stronger predictor of insulin resistance than fitness among 50-95 year olds. *Diabetes care*, 29(3), 673–678.
<https://doi.org/10.2337/diacare.29.03.06.dc05-1605>
- Rajendiran, D., Packirisamy, S., & Gunasekaran, K. (2018). A review on role of antioxidants in diabetes. *Asian J Pharm Clin Res*, 11(2), 48–53. <https://doi.org/10.22159/ajpcr.2018.v11i2.23241>
- Rahmawati, N. T., Hastuti, J., & Suryanti, R. A. (2023). *Anthropometric characteristics of four groups of indigenous people of Wallacea in East Indonesia*. 86(2), 51–63.
- Ramachandran, A. (2012). Trends in prevalence of diabetes in Asian countries. *World J Diabetes*, 3(6), 110. <https://doi.org/10.4239/wjd.v3.i6.110>
- Ramachandran, A., Wan Ma, R. C., & Snehalatha, C. (2010). Diabetes in Asia. *The Lancet*, 375(9712), 408–418. [https://doi.org/10.1016/S0140-6736\(09\)60937-5](https://doi.org/10.1016/S0140-6736(09)60937-5)
- Ranasinghe, P., Piger, S., Galappatthy, P., Katulanda, P., & Constantine, G. R. (2015). Zinc and diabetes mellitus: Understanding molecular mechanisms and clinical implications. *DARU, J Pharm Sci*, 23(1), 1–13.
<https://doi.org/10.1186/s40199-015-0127-4>
- Rhee, E. (2015). Diabetes in Asians. *Endocrinology and Metabolism (Seoul)*, 30(3), 263–269. <https://doi.org/10.3803/EnM.2015.30.3.263>
- Richter, E. A., & Hargreaves, M. (2013). Exercise, GLUT4, and skeletal muscle glucose uptake. *Physiol Rev*, 93(3), 993–1017.
<https://doi.org/10.1152/physrev.00038.2012>
- Riste, L., Khan, F., & Cruickshank, K. (2001). High Prevalence of Type 2 Diabetes in All Ethnic Groups, Including Europeans, in a British Inner City. *Diabetes Care*, 24(8), 1377–1383.
<http://care.diabetesjournals.org/content/24/8/1377>.
- Rong, R., Hanson, R. L., Ortiz, D., Wiedrich, C., Kobes, S., Knowler, W. C., Bogardus, C., & Baier, L. J. (2009). Association analysis of variation in/near FTO, CDKAL1, SLC30A8, HHEX, EXT2, IGF2BP2, LOC387761, and CDKN2B with type 2 diabetes and related quantitative traits in Pima Indians. *Diabetes*, 58(2), 478–488. <https://doi.org/10.2337/db08-0877>
- Ryoo, H., Woo, J., Kim, Y., & Lee, C. (2011). Heterogeneity of genetic associations of CDKAL1 and HHEX with susceptibility of type 2 diabetes mellitus by gender. *Eur J Hum Genet*, 19(6), 672–675.
<https://doi.org/10.1038/ejhg.2011.6>
- Sahrani, W. A., Astuti, I., & Sadewa, A. H. (2016). Polymorphism of transcription factor 7-like 2 gene and HOMA-β level of individuals with and without type 2 diabetes mellitus family history. *Indonesian Journal of*

- Biotechnology*, 21(1), 1–10. <https://doi.org/10.22146/ijbiotech.9312>
- Sakai, K., Imamura, M., Tanaka, Y., Iwata, M., Hirose, H., Kaku, K., Maegawa, H., et al. (2013). Replication Study for the Association of 9 East Asian GWAS-Derived Loci with Susceptibility to Type 2 Diabetes in a Japanese Population. *PLoS ONE*, 8(9), 4–10. <https://doi.org/10.1371/journal.pone.0076317>
- Salas-Salvadó, J., Becerra-Tomás, N., Papandreou, C., & Bulló, M. (2019). Dietary Patterns Emphasizing the Consumption of Plant Foods in the Management of Type 2 Diabetes: A Narrative Review. *Adv Nutr*, 10, S320–S331. <https://doi.org/10.1093/advances/nmy102>
- Samuel, V. T., & Shulman, G. I. (2016). The pathogenesis of insulin resistance: Integrating signaling pathways and substrate flux. *J Clin Investigat*, 126(1), 12–22. <https://doi.org/10.1172/JCI77812>
- Sanghera, D. K., & Blakett, P. R. (2012). Type 2 Diabetes Genetics: Beyond GWAS. *Journal of diabetes & metabolism*, 3(198), 6948. <https://doi.org/10.4172/2155-6156.1000198>
- Santos, M. C. F., Anderson, C. P., Neschen, S., Zumbrennen-Bullough, K. B., Romney, S. J., Kahle-Stephan, M., Rathkolb, B., Gailus-Durner, V., et al. (2020). Irf2 regulates insulin production through iron-mediated Cdkal1-catalyzed tRNA modification. *Nature Communications*, 11(1), 1–16. <https://doi.org/10.1038/s41467-019-14004-5>
- Schulze, M. B., Manson, J. E., Willett, W. C., & Hu, F. B. (2003). Processed meat intake and incidence of Type 2 diabetes in younger and middle-aged women. *Diabetologia*, 46(11), 1465–1473. <https://doi.org/10.1007/s00125-003-1220-7>
- Scott, L. J., Mohlke, K. L., Bonnycastle, L. L., Willer, C. J., Li, Y., Duren, W. L., Erdos, M. R., Stringham, H. M., Chines, P. S., Jackson, A. U., et al. (2007). A genome-wide association study of type 2 diabetes in finns detects multiple susceptibility variants. *Science*, 316(5829), 1341–1345. <https://doi.org/10.1126/science.1142382>
- Seidelmann, S. B., Claggett, B., Cheng, S., Henglin, M., Shah, A., Steffen, L. M., Folsom, A. R., Rimm, E. B., Willett, W. C., & Solomon, S. D. (2018). Dietary carbohydrate intake and mortality: a prospective cohort study and meta-analysis. *The Lancet. Public health*, 3(9), e419–e428. [https://doi.org/10.1016/S2468-2667\(18\)30135-X](https://doi.org/10.1016/S2468-2667(18)30135-X)
- Shahid, A., Saeed, S., Rana, S., & Mahmood, S. (2012). Family history of diabetes and parental consanguinity: Important risk for impaired fasting glucose in South East Asians. *West Indian Med J*, 61(3). <https://doi.org/10.7727/wimj.2011.072>
- Shi, Z., Zhang, C., Zhou, M., Zhen, S., & Taylor, A. W. (2013). Exposure to the Chinese famine in early life and the risk of anaemia in adulthood. *BMC Public Health*, 13(1). <https://doi.org/10.1186/1471-2458-13-904>
- Shu, L., Shen, X. M., Li, C., Zhang, X. Y., & Zheng, P. F. (2017). Dietary patterns are associated with type 2 diabetes mellitus among middle-Aged adults in Zhejiang Province, China. *Nutr J*, 16(1), 1–10.

<https://doi.org/10.1186/s12937-017-0303-0>

- Sim, X., Ong, R. T. H., Suo, C., Tay, W. T., Liu, J., Ng, D. P. K., Boehnke, M., et al. (2011). Transferability of type 2 diabetes implicated loci in multi-ethnic cohorts from Southeast Asia. *PLoS Genetics*, 7(4).
<https://doi.org/10.1371/journal.pgen.1001363>
- Smedley, A., & Smedley, B. D. (2005). Race as biology is fiction, racism as a social problem is real : Anthropological and historical perspectives on the social construction of race. *Am Psychol*, 60(1), 16–26.
<https://doi.org/10.1037/0003-066X.60.1.16>
- Smith, A. D., Crippa, A., Woodcock, J., & Brage, S. (2016). Physical activity and incident type 2 diabetes mellitus: a systematic review and dose–response meta-analysis of prospective cohort studies. *Diabetologia*, 59(12), 2527–2545. <https://doi.org/10.1007/s00125-016-4079-0>
- Snieder, H., Sawtell, P. A., Ross, L., Walker, J., Spector, T. D., & Leslie, R. D. G. (2001). HbA 1c Levels Are Genetically Determined Even in. *Diabetes*, 50(December), 2858–2863. <https://doi.org/10.2337/diabetes.50.12.2858>
- Snijder, M. B., Dekker, J. M., Visser, M., Bouter, L. M., Stehouwer, C. D. A., Kostense, P. J., & Yudkin, J. S. (2003). Associations of hip and thigh circumferences independent of waist circumference with the incidence of type 2 diabetes : the Hoorn Study 1 – 3. *The American Journal of Clinical Nutrition*, 77(5), 1192–1197. <https://doi.org/10.1093/ajcn/77.5.1192>
- Soares, P., Trejaut, J. A., Loo, J. H., Hill, C., Mormina, M., Lee, C. L., Chen, Y. M., et al. (2008). Climate change and postglacial human dispersals in Southeast Asia. *Mol Biol Evol*, 25(6), 1209–1218.
<https://doi.org/10.1093/molbev/msn068>
- Sonestedt, E., Lyssenko, V., Ericson, U., Gullberg, B., Wirfält, E., Groop, L., & Orho-Melander, M. (2012). Genetic variation in the glucose-dependent insulinotropic polypeptide receptor modifies the association between carbohydrate and fat intake and risk of type 2 diabetes in the Malmo diet and Cancer cohort. *J Clin Endocrinol Metab*, 97(5), 810–818.
<https://doi.org/10.1210/jc.2011-2444>
- Sripichai, O., & Fucharoen, S. (2007). Genetic polymorphisms and implications for human diseases. *J Med Assoc Thai*, 90(2), 394–398.
- Stančáková, A., Pihlajamäki, J., Kuusisto, J., Stefan, N., Fritsche, A., Häring, H., Andreozzi, F., et al. (2008). Single-nucleotide polymorphism rs7754840 of CDKAL1 is associated with impaired insulin secretion in nondiabetic offspring of type 2 diabetic subjects and in a large sample of men with normal glucose tolerance. *J Clin Endocrinol Metab*, 93(5), 1924–1930.
<https://doi.org/10.1210/jc.2007-2218>
- Steinthorsdottir, V., Thorleifsson, G., Reynisdottir, I., Benediktsson, R., Jonsdottir, T., Walters, G. B., Stykarsdottir, U., et al. (2007). A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. *Nat Genet*, 39(6), 770–775. <https://doi.org/10.1038/ng2043>
- Strelitz, J., Ahern, A. L., Long, G. H., Hare, M. J. L., Irving, G., Boothby, C. E., Wareham, N. J., & Griffin, S. J. (2019). *Moderate weight change following*

diabetes diagnosis and 10 year incidence of cardiovascular disease and mortality. 1391–1402.

- Sun, X. F., Xiao, X. H., Zhang, Z. X., Liu, Y., Xu, T., Zhu, X. L., Zhang, Y., Wu, X. P., et al. (2015). Positive association between type 2 diabetes risk alleles near CDKAL1 and reduced birthweight in Chinese han individuals. *Chin Med J*, 128(14), 1873–1878. <https://doi.org/10.4103/0366-6999.160489>
- Sun, X., Yu, W., & Hu, C. (2014). Genetics of type 2 diabetes: Insights into the pathogenesis and its clinical application. *BioMed Res Int*, 2014. <https://doi.org/10.1155/2014/926713>
- Sunil, B., & Ashraf, A. P. (2020). Dyslipidemia in pediatric type 2 diabetes mellitus. *Current Diabetes Reports*, 20(10), 53. <https://doi.org/10.1007/s11892-020-01336-6>
- Tabák, A. G., Jokela, M., Akbaraly, T. N., Brunner, E. J., Kivimäki, M., & Witte, D. R. (2009). Trajectories of glycaemia, insulin sensitivity, and insulin secretion before diagnosis of type 2 diabetes: an analysis from the Whitehall II study. *The Lancet*, 373, 2215–2221. [https://doi.org/10.1016/S0140-6736\(09\)60619-X](https://doi.org/10.1016/S0140-6736(09)60619-X)
- Tan, J., Ng, D. P. K., Nurbaya, S., ... Tai, E.-S. (2009). Polymorphisms identified through genome-wide association studies and their associations with type 2 diabetes in Chinese, Malays, and Asian-Indians in Singapore. *Endocrine Reviews*, 30(7), 926–926. <https://doi.org/10.1210/edrv.30.7.9999>
- Tan, N. C., Sankari, U., Ng, C. E., Ling, Y., & Koh, E. (2022). Longitudinal study on the progression of muscle status among community-dwelling ambulatory older multiethnic Asians with type 2 diabetes mellitus. *BMC Geriatrics*, 1–11. <https://doi.org/10.1186/s12877-022-03098-0>
- Tay, J., Luscombe-Marsh, N. D., Thompson, C. H., Noakes, M., Buckley, J. D., Wittert, G. A., Yancy, W. S., Jr., & Brinkworth, G. D. (2014). A very low-carbohydrate, low-saturated fat diet for Type 2 diabetes management: A randomized trial. *Diabetes Care*, 37(11), 2909–2918. <https://doi.org/10.2337/dc14-0845>
- Tefu, M. O., & Karwur, F. F. (2017). Pitarah Manusia Nusa Tenggara Timur Berdasarkan Ceritera Kromosom Y. *Scientiae Educatia*, 6(2), 143. <https://doi.org/10.24235/sc.educatia.v6i2.1622>
- Tian, Y., Xu, J., Huang, T., Cui, J., Zhang, W., Song, W., Chen, H., et al. (2019). A Novel Polymorphism (rs35612982) in CDKAL1 Is a Risk Factor of Type 2 Diabetes: A Case-Control Study. *Kidney and Blood Press Res*, 44(6), 1313–1326. <https://doi.org/10.1159/000503175>
- Tian, S., Xu, Q., Jiang, R., Han, T., Sun, C., & Na, L. (2017). Dietary Protein Consumption and the Risk of Type 2 Diabetes: A Systematic Review and Meta-Analysis of Cohort Studies. *Nutrients*, 9(9), 982. <https://doi.org/10.3390/nu9090982>
- Triandhini, R. L. N. K. R., Sadewa, A. H., & Helmyati, S. (2025). Exploring genes related to impaired insulin secretion and the interaction with diet in

- type 2 diabetes mellitus: A scoping review of observational studies. *Obesity Medicine*, 55, Article 100611. <https://doi.org/10.1016/j.obmed.2025.100611>
- Tsai, F. J., Yang, C. F., Chen, C. C., Chuang, L. M., Lu, C. H., Chang, C. T., Wang, T. Y., et al (2010). A genome-wide association study identifies susceptibility variants for type 2 diabetes in Han Chinese. *PLoS Genetics*, 6(2). <https://doi.org/10.1371/journal.pgen.1000847>
- Tuerxunyiming, M., Mohemaiti, P., Wufuer, H., & Tuheti, A. (2015). Association of rs7754840 G/C polymorphisms in CDKAL1 with type 2 diabetes: A meta-analysis of 70141 subjects. *Int J Clin Exp Med*, 8(10), 17392–17405.
- Tumonggor, M. K., Karafet, T. M., Downey, S., Lansing, J. S., Norquest, P., Sudoyo, H., Hammer, M. F., & Cox, M. P. (2014). Isolation, contact and social behavior shaped genetic diversity in West Timor. *Journal of human genetics*, 59(9), 494–503. <https://doi.org/10.1038/jhg.2014.62>
- Tumonggor, M. K., Karafet, T. M., Hallmark, B., Lansing, J. S., Sudoyo, H., Hammer, M. F., & Cox, M. P. (2013). The Indonesian archipelago : an ancient genetic highway linking Asia and the Pacific. *Journal of Human Genetics*, 58(3), 165–173. <https://doi.org/10.1038/jhg.2012.154>
- Ubeda, M., Rukstalis, J. M., & Habener, J. F. (2006). Inhibition of cyclin-dependent kinase 5 activity protects pancreatic beta cells from glucotoxicity. *J Biol Chem*, 281(39), 28858– 28864. <https://doi.org/10.1074/jbc.M604690200>
- Unoki, H., Takahashi, A., Kawaguchi, T., Hara, K., Horikoshi, M., Andersen, G., Ng, D. P. K., Holmkvist, J., et al. (2008). SNPs in KCNQ1 are associated with susceptibility to type 2 diabetes in East Asian and European populations. *Nat Gene*, 40(9), 1098–1102. <https://doi.org/10.1038/ng.208>
- US Department of Health and Human Services. (2020). National Diabetes Statistics Report, 2020. *National Diabetes Statistics Report*, 2.
- Valdez, R., Yoon, P. W., Liu, T., & Khoury, M. J. (2007). Family history and prevalence of diabetes in the U.S. population: The 6-year results from the National Health and Nutrition Examination Survey (1999-2004). *Diabetes Care*, 30(10), 2517–2522. <https://doi.org/10.2337/dc07-0720>
- Van Dam, R. M., Willett, W. C., Rimm, E. B., Stampfer, M. J., & Hu, F. B. (2002). Dietary fat and meat intake in relation to risk of type 2 diabetes in men. *Diabetes Care*, 25(3), 417–424. <https://doi.org/10.2337/diacare.25.3.417>
- Van Nielen, M., Feskens, Edith, J. M., Mensink, M., Sluijs, I., Molina, E., Amiano, P., Beulens, J. W. J., & Boeing, H. (2014). Dietary Protein Intake and Incidence of Type 2 Diabetes in Europe : The EPIC-InterAct Case-Cohort Study. 37(July), 1854–1862. <https://doi.org/10.2337/dc13-2627>
- Van Tilburg, J., Van Haefen, T. W., Pearson, P., & Wijmenga, C. (2001). Defining the genetic contribution of type 2 diabetes mellitus. *J Med Gene*, 38(9), 569–578. <https://doi.org/10.1136/jmg.38.9.569>
- Villegas, R., Goodloe, R. J., McClellan, B. E., Boston, J., & Crawford, D. C. (2014). Gene-carbohydrate and gene-fiber interactions and type 2 diabetes in

- diverse populations from the National Health and Nutrition Examination Surveys (NHANES) as part of the Epidemiologic Architecture for Genes Linked to Environment (EAGLE) study. *BMC Genetics*, 15(1), 1–8. <https://doi.org/10.1186/1471-2156-15-69>
- Villegas, R., Shu, X. O., Yang, G., Matthews, C. E., Li, H., Cai, H., Gao, Y.-T., & Zheng, W. (2009). Energy balance and type 2 diabetes: A report from the Shanghai Women’s Health Study. *Nutrition, Metabolism & Cardiovascular Diseases*, 19(3), 190–197. <https://doi.org/10.1016/j.numecd.2008.06.003>
- Wake, A. D. (2021). Alcohol consumption and the incidence of type 2 diabetic mellitus: “issue that requires more attention.” *Int J Endocrinol Metab*, 8(6), 143–147. <https://doi.org/10.15406/emij.2021.08.00298>
- Wang, Y., Mi, J., Shan, X. Y., Wang, Q. J., & Ge, K. Y. (2007). Is China facing an obesity epidemic and the consequences? The trends in obesity and chronic disease in China. *Int J Obes*, 31(1), 177–188. <https://doi.org/10.1038/sj.ijo.0803354>
- Wang, J., Luben, R., Khaw, K.-T., Bingham, S., Wareham, N. J., & Forouhi, N. G. (2008). *Dietary Energy Density Predicts the Risk of Incident Type 2 Diabetes The European Prospective Investigation of Cancer (EPIC) - Norfolk Study*. 31(11). <https://doi.org/10.2337/dc08-1085>.
- Wang, S., Ma, W., Yuan, Z., Wang, S. M., Yi, X., Jia, H., & Xue, F. (2016). Association between obesity indices and type 2 diabetes mellitus among middle-aged and elderly people in Jinan, China: A cross-sectional study. *BMJ Open*, 6(11), 1–9. <https://doi.org/10.1136/bmjopen-2016-012742>
- Wei, F. Y., Suzuki, T., Watanabe, S., Kimura, S., Kaitsuka, T., Fujimura, A., Matsui, H., Atta, M., Michiue, H., Fontecave, M., Yamagata, K., Suzuki, T., & Tomizawa, K. (2011). Deficit of tRNA^{Lys} modification by Cdkal1 causes the development of type 2 diabetes in mice. *J Clin Invest*, 121(9), 3598–3608. <https://doi.org/10.1172/JCI58056>
- Wei, F. Y., & Tomizawa, K. (2011). Functional loss of Cdkal1, a novel tRNA modification enzyme, causes the development of type 2 diabetes. *Endocr J*, 58(10), 819–825. <https://doi.org/10.1507/endocrj.EJ11-0099>
- Wiardani, N. K., Hadi, H., & Huriyati, E. (2007). Pola makan dan obesitas sebagai faktor risiko diabetes mellitus tipe 2 di Rumah Sakit Sanglah Denpasar. *Jurnal Gizi Klinik Indonesia*, 4(1), 1–10. <https://doi.org/10.22146/ijcn.17456>
- Willett, W., Manson, J., & Liu, S. (2002). Glycemic index, glycemic load, and risk of type 2 diabetes 1,2,3. *The American Journal of Clinical Nutrition*, 76(1), 274S-280S. <https://doi.org/10.1093/ajcn/76.1.274S>
- Wirström, T., Hilding, A., Gu, H. F., Östenson, C. G., & Björklund, A. (2013). Consumption of whole grain reduces risk of deteriorating glucose tolerance, including progression to prediabetes. *Am J Clin Nutr*, 97(1), 179–187. <https://doi.org/10.3945/ajcn.112.045583>
- World Health Organization. Regional Office for the Western Pacific. (2000). *Obesity Task Force. International Association for the Study of Obesity. The*

Asia – Pacific perspective: redefining obesity and its treatment.

World Health Organization. (2014). *Global Physical Activity Questionnaire (GPAQ) analysis guide*

World Health Organization. (2016). *Global report on diabetes.*
<https://apps.who.int/iris/handle/10665/204871>

World Health Organization. (2019). Classification of diabetes mellitus. *Clin Lab Med* (Vol. 21, Issue 1). https://doi.org/10.5005/jp/books/12855_84

Wu, Y., Ding, Y., Tanaka, Y., & Zhang, W. (2014). Risk factors contributing to type 2 diabetes and recent advances in the treatment and prevention. *Int J Med Sci*, 11(11), 1185–1200. <https://doi.org/10.7150/ijms.10001>

Wysham, C., & Shubrook, J. (2020). Beta-cell failure in type 2 diabetes: mechanisms, markers, and clinical implications. *Postgraduate Medicine*, 132(8), 676–686. <https://doi.org/10.1080/00325481.2020.1771047>

Xie, P., Wei, F. Y., Hirata, S., Kaitsuka, T., Suzuki, T., Suzuki, T., & Tomizawa, K. (2013). Quantitative PCR measurement of tRNA 2-Methylthio modification for assessing type 2 diabetes risk. *Clin Chem*, 59(11), 1604–1612. <https://doi.org/10.1373/clinchem.2013.210401>

Xu, H., Luo, J., Huang, J., & Wen, Q. (2018). Flavonoids intake and risk of type 2 diabetes mellitus: A meta-analysis of prospective cohort studies. *Medicine (United States)*, 97(19), 1–7.
<https://doi.org/10.1097/MD.00000000000010686>

Xu, N., Zhang, T., Han, W., Yin, L., Ma, N., Shi, X., & Sun, J. (2021). *Association of CDKAL1 RS10946398 Gene Polymorphism with Susceptibility to Diabetes Mellitus Type 2 : A Meta-Analysis. 2021.*
<https://doi.org/10.1155/2021/1254968>

Xu, S., Qiao, N., Huang, J., Sun, C., Cui, Y., Liang, J., Lu, Q., & Wang, T. (2016). Gender Differences in Dietary Patterns and Their Association with the Prevalence of Metabolic Syndrome among Chinese : A Cross-Sectional Study. 1–17. <https://doi.org/10.3390/nu8040180>

Yabe, D., Seino, Y., Fukushima, M., & Seino, S. (2015). *β Cell Dysfunction Versus Insulin Resistance in the Pathogenesis of Type 2 Diabetes in East Asians.* <https://doi.org/10.1007/s11892-015-0602-9>

Yajnik, C. S. (2009). Nutrient-mediated teratogenesis and fuel-mediated teratogenesis: Two pathways of intrauterine programming of diabetes. *Int J Gynecol Obstet*, 104(SUPPL.), S27–S31.
<https://doi.org/10.1016/j.ijgo.2008.11.034>

Yamazaki, H., Tauchi, S., Haueise, T., Yamamoto, Y., Dohke, M., Hanawa, N., Kodama, Y., Katanuma, A., Stefan, N., Fritsche, A., & Birkenfeld, A. L. (2022). Fat Distribution Patterns and Future Type 2 Diabetes. 71(September), 1937–1945.

Yang, J. J., Yu, D., Wen, W., Saito, E., Rahman, S., Shu, X. O., Chen, Y., et al. (2019). Association of Diabetes with All-Cause and Cause-Specific Mortality in Asia: A Pooled Analysis of More Than 1 Million Participants. *JAMA Network Open*, 2(4). <https://doi.org/10.1001/jamanetworkopen.2019.2696>

- Yang, J., Mao, Q. X., Xu, H. X., Ma, X., & Zeng, C. Y. (2014). Tea consumption and risk of type 2 diabetes mellitus: A systematic review and meta-analysis update. *BMJ Open*, *4*(7). <https://doi.org/10.1136/bmjopen-2014-005632>
- Yasuda, K., Miyake, K., Horikawa, Y., Hara, K., Osawa, H., Furuta, H., Hirota, Y., et al. (2008). Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. *Nat Genet*, *40*(9), 1092–1097. <https://doi.org/10.1038/ng.207>
- Yazdi, K. V., Rezaei, M., Shadmani, F., & Ghorbani, A. (2020). SLC30A8, CDKAL1, TCF7L2, KCNQ1 and IGF2BP2 are associated with Type 2 diabetes mellitus in Iranian patients. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, *13*, 3497–3506. <https://doi.org/10.2147/DMSO.S276924>
- Yun, K. J., Han, K., Kim, M. K., Park, Y. M., Baek, K. H., Song, K. H., & Kwon, H. S. (2016). Insulin Resistance Distribution and Cut-Off Value in Koreans from the 2008-2010 Korean National Health and Nutrition Examination Survey. *PloS one*, *11*(4), e0154593. <https://doi.org/10.1371/journal.pone.0154593>
- Zeevi, D., Korem, T., Zmora, N., Halpern, Z., Elinav, E., Segal, E., Zeevi, D., Korem, T., Zmora, N., Israeli, D., Rothschild, D., & Weinberger, A. (2015). *Personalized Nutrition by Prediction of Glycemic Article Personalized Nutrition by Prediction of Glycemic Responses*. 1079–1094. <https://doi.org/10.1016/j.cell.2015.11.001>
- Zeggini, E., Weedon, M. N., Lindgren, C. M., Timothy, M., Elliott, K. S., Lango, H., Timpson, N.J., et al. (2007). Multiple type 2 diabetes susceptibility genes following genome-wide association scan in UK samples. *Science*, *316*(5829), 1336–1341. <https://doi.org/10.1126/science.1142364>. Multiple
- Zeng, C. P., Chen, Y. C., Lin, X., Greenbaum, J., Chen, Y. P., Peng, C., Wang, X. F., et al. (2017). Increased identification of novel variants in type 2 diabetes, birth weight and their pleiotropic loci. *J Diabetes*, *9*(10), 898–907. <https://doi.org/10.1111/1753-0407.12510>
- Zeng, H., Guo, M., Zhou, T., Tan, L., Chong, C. N., Zhang, T., Dong, X., et al. (2018). An isogenic human ESC platform for functional evaluation of genome-wide association study identified diabetes genes and drug discovery. *Cell Stem Cell*, *19*(3), 326–340. <https://doi.org/10.1016/j.stem.2016.07.002>
- Zeqollari, A., Spahiu, K., Vyshka, G., & Çakërri, L. (2014). Lipid Profile in Diabetes Mellitus Type 2 Patients in Albania and the Correlation with BMI, Hypertension, and Hepatosteatosis. *J Family Med Community Health* *1*(4): 1018
- Zheng, J. S., Arnett, D. K., Parnell, L. D., Smith, C. E., Li, D., Borecki, I. B., Tucker, K. L., Ordovás, J. M., & Lai, C. Q. (2013). Modulation by dietary fat and carbohydrate of IRS1 association with type 2 diabetes traits in two populations of different ancestries. *Diabetes Care*, *36*(9), 2621–2627. <https://doi.org/10.2337/dc12-2607>
- Zheng, J. S., Huang, T., Li, K., Chen, Y., Xie, H., Xu, D., Sun, J., & Li, D.

(2015). Modulation of the Association between the PEPD Variant and the Risk of Type 2 Diabetes by n-3 Fatty Acids in Chinese Hans. *J Nutrigenet Nutrigenomics*, 8(1), 36–43. <https://doi.org/10.1159/000381348>

Zhou, Z., Sun, B., Huang, S., Jia, W., & Yu, D. (2019). The tRNA-associated dysregulation in diabetes mellitus. *Metabolism: Clinical and Experimental*, 94, 9–17. <https://doi.org/10.1016/j.metabol.2019.01.017>