

DAFTAR PUSTAKA

- Abdullah S, Abdul Mudalip SK, Shaarani SMD, Che Pi NA. 2010. Ultrasonic Extraction of Oil from *Monopterus albus*: Effects of Different Ultrasonic Power, Solvent Volume and Sonication Time. *Journal of Applied Sciences*. 10:2713–2716
- Acosta, E. (2009). Bioavailability of nanoparticles in nutrient and nutraceutical delivery. *Current Opinion in Colloid and Interface Science*, 14(1), 3–15. <https://doi.org/10.1016/j.cocis.2008.01.002>
- Adepoju, M. A., Omitoyin, B. O., Ajani, E. K., & Asha, K. (2018). Effect of Smoking Time and Temperature on the Proximate Composition and Quality of Milkfish Steaks. *Journal of Aquatic Food Product Technology*, 27(3), 369–378. <https://doi.org/10.1080/10498850.2018.1437494>
- Agustini, T.W., Susilowati I., Subagyo., Setyati, W.A. dan Wibowo, B.A. 2011, Will Soft-Boned Milkfish- A Traditional Food Product From Semarang City. *Journal of Coastal Development*. Semarang. 14:1, 81-90
- Aissou, M., Chemat-Djenni, Z., Yara-Varón, E., Fabiano-Tixier, A. S., & Chemat, F. (2017). Limonene as an agro-chemical building block for the synthesis and extraction of bioactive compounds. *Comptes Rendus Chimie*, 20(4), 346–358. <https://doi.org/10.1016/j.crci.2016.05.018>
- Anas, A. N. M., Rahman, N. A. A., Tajuddin, R., & Hamid, N. S. A. (2020). Supercritical fluid extraction of fish oil from sardine (*Sardinella* sp.): Optimization and characterization. *Journal of Food Science and Technology*, 57(7), 2516–2525. <https://doi.org/10.1007/s13197-020-04315-z>
- Amin, H. A., Elbeltagy, A. E., Mostafa, M. M., & Khalil, A. H. (2014). Development of low fat mayonnaise containing different types and levels of hydrocolloid gum. *Journal of Agroalimentary Processes and Technologies*, 20(1), 54–63.
- AOAC. (2000). Official methods of analysis of analytical chemists. AOAC Inc.
- Arifah, N. Z., Alamsjah, M. A., & Sulmartiwi, L. (2024). Karakteristik Fisikokimia, Organoleptik, dan Kandungan Gizi Mayonnaise Minyak Ikan Lemuru (*Sardinella* sp.). *Bioscientist: Jurnal Ilmiah Biologi*, 12(1), 92. <https://doi.org/10.33394/bioscientist.v12i1.10712>

- Ashaq, B., Rasool, K., Habib, S., Bashir, I., Nisar, N., Mustafa, S., Ayaz, Q., Nayik, G. A., Uddin, J., Ramniwas, S., Mugabi, R., & Wani, S. M. (2024). Insights into chemistry, extraction and industrial application of lemon grass essential oil -A review of recent advances. *Food Chemistry: X*, 22(April). <https://doi.org/10.1016/j.fochx.2024.101521>
- Badan Pusat Statistik. (2023). Produksi perikanan budidaya menurut komoditas utama (ton). Diakses dari <https://www.bps.go.id/id/statistics-table/2/MTUxMyMy/produksi-perikanan-budidaya-menurut-komoditas-utama.html>
- Badan Pengawas Obat dan Makanan Republik Indonesia. (2019). Peraturan Kepala BPOM Nomor 11 Tahun 2019 tentang Bahan Tambahan Pangan. Badan Pengawas Obat dan Makanan Republik Indonesia
- Balakrishnan, J., Kannan, S., & Govindasamy, A. (2021). Structured form of DHA prevents neurodegenerative disorders: A better insight into the pathophysiology and the mechanism of DHA transport to the brain. *Nutrition Research*, 85, 119–134. <https://doi.org/10.1016/j.nutres.2020.12.003>
- Barizão, É. O., Boeing, J. S., Martins, A. C., Visentainer, J. V, & Almeida, V. C. (2015). Application of Response Surface Methodology for the Optimization of Ultrasound-Assisted Extraction of Pomegranate (*Punica granatum* L.) Seed Oil. *Food Analytical Methods*, 8(9), 2392–2400. <https://doi.org/10.1007/s12161-015-0135-5>
- Belarbi, E.H., Molina, E., Chisti, Y., 2000. A process for high yield and scaleable recovery of high purity eicosapentaenoic acid esters from microalgae and fish oil. *Enzyme Microb. Technol.* 26, 516–529. [https://doi.org/10.1016/S0141-0229\(99\)00191-X](https://doi.org/10.1016/S0141-0229(99)00191-X).
- Bertouche, S., Tomao, V., Hellal, A., Boutekdjiret, C., & Chemat, F. (2013). First approach on edible oil determination in oilseeds products using alpha-pinene. *Journal of Essential Oil Research*, 25(6), 439–443. <https://doi.org/10.1080/10412905.2013.782473>
- Biancarosa, I., Espe, M., Bruckner, C., Heesch, S., Liland, N., Waagbø, R., Torstensen, B., & LCK, E.-J. (2017). Amino acid composition, protein content, and nitrogen-to-protein conversion factors of 21 seaweed species from Norwegian waters. *Journal of Applied Phycology*, 29. <https://doi.org/10.1007/s10811-016-0984-3>

- Bija, S., Suseno, S. H., & Uju, U. (2017). Purification of Sardine Fish Oil Through Degumming and Neutralization. 20(1), 143–152. <https://doi.org/10.17844/JPHPI.V20I1.16501>
- Bodoira, R. M., Rodríguez Ruiz, A. C., Martínez, M. L., Velez, A. R., Ribotta, P. D., & Maestri, D. M. (2024). From by-product to natural antioxidant: Incorporation of peanut skin extract in mayonnaise and its effect on physico-chemical and sensory properties. *Food Bioscience*, 61, 104680. <https://doi.org/10.1016/j.fbio.2024.104680>
- Boran, G., & Karaçam, H. (2011). Seasonal changes in proximate composition of some fish species from the black sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 11(1), 01–05. <https://doi.org/10.4194/trjfas.2011.0101>
- Borlongan, I.G. and Benitez, L.V., 1992. Lipid and fatty acid composition of milkfish (*Chanos chanos*Forsskal) grown in freshwater and seawater. *Aquaculture*, 104: 79-89.
- Bruno, S. F., Kudre, T. G., & Bhaskar, N. (2019). Impact of pretreatment-assisted enzymatic extraction on recovery, physicochemical and rheological properties of oil from head. *Journal of Food Process Engineering*, 42(3), e12990. <https://doi.org/https://doi.org/10.1111/jfpe.12990>
- Bureau of Fisheries and Aquatic Resources (BFAR), 2001. The 2000 Philippine Fisheries Profile. Quezon City, Philippines: Department of Agriculture.
- Burlingame, B., Nishida, C., Uauy, R., & Weisell, R. (2009). Fats and fatty acids in human nutrition. *FAO Food and Nutrition Paper* 91.
- Bustamante, J., van Stempvoort, S., García-Gallarreta, M., Houghton, J. A., Briers, H. K., Budarin, V. L., Matharu, A. S., & Clark, J. H. (2016). Microwave assisted hydro-distillation of essential oils from wet citrus peel waste. *Journal of Cleaner Production*, 137, 598–605. <https://doi.org/https://doi.org/10.1016/j.jclepro.2016.07.108>
- Cai, D., Li, P., Luo, Z.F., Qin, P.Y., Chen, C.J., Wang, Y., Wang, Z., Tan, T.W., 2016. Effect of dilute alkaline pretreatment on the conversion of different parts of corn stalk to fermentable sugars and its application in acetone–butanol–ethanol fermentation. *Bioresour. Technol.* 211, 117–124 .
- Carrera, C., Pastol, J., Setyaningsih, W., Ruiz-Rodríguez, A., Ferreiro-González, M., Barbero, G. F., & Palma, M. (2021). Optimization by means of chemometric tools of an ultrasound-assisted method for the extraction of

betacyanins from red dragon fruit (*Hylocereus polyrhizus*). *Agronomy*, 11(6).
<https://doi.org/10.3390/agronomy11061053>

Cascant, M. M., Cascant, M. M., Breil, C., Garrigues, S., de la Guardia, M., Fabiano-Tixier, A. S., & Chemat, F. (2017). A green analytical chemistry approach for lipid extraction: computation methods in the selection of green solvents as alternative to hexane. *Analytical and Bioanalytical Chemistry*, 409(14), 3527–3539. <https://doi.org/10.1007/S00216-017-0323-9>

Çavdar, H. K., Bilgin, H., Fadiloğlu, S., & Yılmaz, F. M. (2022). Ultrasound- and Microwave-Assisted Extraction Facilitate Oil Recovery from Gilthead Seabream (*Sparus aurata*) by-Products and Enhance Fish Oil Quality Parameters. *European Journal of Lipid Science and Technology*, 2200089. <https://doi.org/https://doi.org/10.1002/ejlt.202200089>

Cengiz, E., Dogan, M., & Karaman, S. (2013). Characterization of rheological interactions of Gleditsia triacanthos gum with some hydrocolloids: Effect of hydration temperature. *Food Hydrocolloids*, 32(2), 453–462. <https://doi.org/https://doi.org/10.1016/j.foodhyd.2013.01.018>

Chakraborty, K., & Joseph, D. (2015). Cooking and pressing is an effective and eco-friendly technique for obtaining high quality oil from *Sardinella longiceps*. *European Journal of Lipid Science and Technology*, 117(6), 837–850. doi:10.1002/ejlt.201400539

Chemat, F., Rombaut, N., Sicaire, A. G., Meullemiestre, A., Fabiano-Tixier, A. S., & Abert-Vian, M. (2017). Ultrasound assisted extraction of food and natural products. Mechanisms, techniques, combinations, protocols and applications. A review. *Ultrasonics Sonochemistry*, 34, 540–560. <https://doi.org/10.1016/j.ultsonch.2016.06.035>

Chemat, S., Tomao, V., & Chemat, F. (2012). Limonene as Green Solvent for Extraction of Natural Products (pp. 175–186). Springer, Dordrecht. https://doi.org/10.1007/978-94-007-1712-1_5

Chemat-Djenni, Z., Ferhat, M. A., Tomao, V., & Chemat, F. (2010). Carotenoid extraction from tomato using a green solvent resulting from orange processing waste. *Journal of Essential Oil-Bearing Plants*, 13(2), 139–147. <https://doi.org/10.1080/0972060X.2010.10643803>

Ciftci, O. N., & Çavdar, H. K. (2025). Ultrasound-Assisted Extraction Improved the Extraction Rate, Omega-3, and Polyunsaturated Fatty Acid Content of

Fish Oil From Atlantic Bonito Waste. *European Journal of Lipid Science and Technology*. <https://doi.org/10.1002/ejlt.202400131>

Ciriminna, R., Meneguzzo, F., Delisi, R., & Pagliaro, M. (2019). Citrus waste as a source of limonene: Extraction and applications. *ChemistrySelect*, 4(18), 5106–5115. <https://doi.org/10.1002/slct.201900394>

Ciriminna, R., Meneguzzo, F., Delisi, R., Pagliaro, M., 2017. Enhancing and improving the extraction of omega-3 from fish oil. *Sustain. Chem. Pharm.* <https://doi.org/10.1016/j.scp.2017.03.001>.

Crexvi, V. T., Monte, M. L., Soares, L. A. de S., & Pinto, L. A. A. (2010). Production and refinement of oil from carp (*Cyprinus carpio*) viscera. *Food Chemistry*, 119(3), 945–950. <https://doi.org/https://doi.org/10.1016/j.foodchem.2009.07.050>

Cui, G., Yang, X., Liu, Z., Wei, M., Liu, T., Gu, H., & Yang, L. (2022). Potential Use of Limonene as an Alternative Solvent for Extraction of Gutta-Percha from *Eucommia ulmoides*. *ACS Sustainable Chemistry and Engineering*, 10(33), 11057–11068. <https://doi.org/10.1021/acssuschemeng.2c03898>

Darko, H. S., Ismaiel, L., Fanesi, B., Pacetti, D., & Lucci, P. (2024). Current Trends in Food Processing By-Products as Sources of High Value-Added Compounds in Food Fortification. In *Foods* (Vol. 13, Issue 17). <https://doi.org/10.3390/foods13172658>

Dave, J., Kumar, V., Kingwascharapong, P., Moula Ali, A. M., & Bavisetty, S. C. B. (2024). A bilayer emulsion containing encapsulated catfish oil, stabilized with whey protein isolates and fish gelatin for development of PUFA rich mayonnaise. *Journal of Food Measurement and Characterization*, 18(11), 9209–9222. <https://doi.org/10.1007/s11694-024-02872-5>

De Oliveira, D. A. S. B., Minozzo, M. G., Licodiedoff, S., & Waszczyński, N. (2016). Physicochemical and sensory characterization of refined and deodorized tuna (*Thunnus albacares*) by-product oil obtained by enzymatic hydrolysis. *Food Chemistry*, 207, 187–194. <https://doi.org/10.1016/j.foodchem.2016.03.069>

Demesa, A. G., Saavala, S., Pöysä, M., & Koiranen, T. (2024). Overview and Toxicity Assessment of Ultrasound-Assisted Extraction of Natural Ingredients from Plants. *Foods*, 13(19), 1–13. <https://doi.org/10.3390/foods13193066>

- Demydova, A. O., Aksonova, O. F., Molchenko, S. M., & Hladkyi, F. F. (2021). Research on a New Approach To Low-Temperature Deodorization and Its Effect on Oxidative Deterioration of Fish Oil. *Journal of Chemistry and Technologies*, 29(4), 639–649. <https://doi.org/10.15421/jchemtech.v29i4.238356>
- Depree, J.A. and G.P. Savage, 2001. Physical and flavour stability of mayonnaise. *Trends in Food Science & Technology*. 12: 157-163.
- Dickinson, E. (2008). Emulsification and emulsion stabilization with protein-polysaccharide complexes. *Gums and Stabilisers for the Food Industry* 14, 221–232.
- Din JN, Harding SA, Valerio CJ, Sarma J, Lyall K, Riemersma RA. 2008. Dietary intervention with oil rich fish reduces platelet-monocyte aggregation in man. *Atherosclerosis Journal*. 197(1):290-6. doi: 10.1016/j.atherosclerosis.2007.04.047.
- Djordjevic, M. S., Cekic, D. N., Isailovic, M. T., Milic, R. J., Vuleta, M. G., Lazic, L. M., & Savic, D. S. (2013). Nanoemulsions produced with varied type of emulsifier and oil content: An influence of formulation and process parameters on the characteristics and physical stability. *Chemical Industry*, 67(5), 795–809. <https://doi.org/10.2298/HEMIND120905005D>
- Egerton, S., Mannion, D., Culloty, S., Whooley, J., Stanton, C., & Ross, R. P. (2020). The proximate composition of three marine pelagic fish: blue whiting (*Micromesistius poutassou*), boarfish (*Capros aper*) and Atlantic herring (*Clupea harengus*). *Irish Journal of Agricultural and Food Research*, 59(1), 185–200. <https://doi.org/10.15212/ijaf-2020-0112>
- El-Bostany, A. N., Ahmed, M. G., & Salem, A. (2011). Development of light mayonnaise formula using carbohydrate-based fat replacement. *Australian Journal of Basic and Applied Sciences*, 5, 673–682.
- El-Rahman, F., Mahmoud, N., Badawy, A. E. K., & Younis, S. M. (2018). Extraction of Fish Oil from Fish Viscera. *The Egyptian Journal of Chemistry*, 61(2), 225–235. <https://doi.org/10.21608/EJCHEM.2018.2798.1230>
- El-Waseif, M., Saed, B., Fahmy, H., Sabry, A., Shaaban, H., Abdelgawad, M., Amin, A., & Farouk, A. (2022). Mayonnaise Enriched with Flaxseed Oil: Omega-3 Fatty Acids Content, Sensory Quality and Stability during the Storage. *Foods*, 11(15). <https://doi.org/10.3390/foods11152288>

- Fang, Y., Gu, S., Zhang, J., Liu, S., Ding, Y., & Liu, J. (2017). Deodorisation of fish oil by nanofiltration membrane process: focus on volatile flavour compounds and fatty acids composition. *International Journal of Food Science & Technology*, 53. <https://doi.org/10.1111/ijfs.13644>
- FAO. 2022. The state of world fisheries and aquaculture 2022. Towards Blue Transformation. FAO, Rome.
- Fennema, O.R. 1996. Food Chemistry, Thrid Edition, Marcel Dekker Inc, Newyork
- Ferreira, S. L. C., Bruns, R. E., Ferreira, H. S., Matos, G. D., David, J. M., Brandão, G. C., da Silva, E. G. P., Portugal, L. A., dos Reis, P. S., Souza, A. S., & dos Santos, W. N. L. (2007). Box-Behnken design: An alternative for the optimization of analytical methods. *Analytica Chimica Acta*, 597(2), 179–186. <https://doi.org/https://doi.org/10.1016/j.aca.2007.07.011>
- Flamminii, F., Daniela, C., Mattia, D., Sacchetti, G., Neri, L., Mastrocola, D., & Pittia, P. (2020). Physical and sensory properties of mayonnaise enriched with encapsulated olive leaf phenolic extracts. *Foods*, 9(997), 1–12.
- Fournier, V., Juanéda, P., Destailats, F., Dionisi, F., Lambelet, P., Sébédio, J. L., & Berdeaux, O. (2006). Analysis of eicosapentaenoic and docosahexaenoic acid geometrical isomers formed during fish oil deodorization. *Journal of Chromatography*, 1129(1), 21–28. <https://doi.org/10.1016/j.chroma.2006.06.089>
- García-Márquez, E., Higuera-Ciapara, I., & Espinosa-Andrews, H. (2017). Design of fish oil-in-water nanoemulsion by microfluidization. *Innovative Food Science and Emerging Technologies*, 40, 87–91. <https://doi.org/10.1016/j.ifset.2016.11.007>
- Gharibzahedi, S. M. T., & Jafari, S. M. (2018). Fabrication of Nanoemulsions by Ultrasonication (pp. 233–285). Academic Press. <https://doi.org/10.1016/B978-0-12-811838-2.00009-6>
- Ghelichi, S., Hajfathalian, M., García-Moreno, P. J., Yesiltas, B., Moltke-Sørensen, A.-D., & Jacobsen, C. (2021). Chapter 19 - Food enrichment with omega-3 polyunsaturated fatty acids (P. J. García-Moreno, C. Jacobsen, A.-D. Moltke Sørensen, & B. B. T.-O.-3 D. S. Yesiltas (eds.); pp. 395–425). Academic Press. <https://doi.org/https://doi.org/10.1016/B978-0-12-821391-9.00020-X>
- Giacometti, J., Bursać Kovačević, D., Putnik, P., Gabrić, D., Bilušić, T., Krešić, G., Stulić, V., Barba, F. J., Chemat, F., Barbosa-Cánovas, G., & Režek Jambrak, A. (2018). Extraction of bioactive compounds and essential oils from

mediterranean herbs by conventional and green innovative techniques: A review. *Food Research International*, 245–262. <https://doi.org/10.1016/j.foodres.2018.06.036>

Glatzer, H. J., & Doraiswamy, L. K. (2000). Rate enhancements due to autocatalysis and heterogenization in phase transfer catalysis: a comparative study. *Chemical Engineering Science*, 55(21), 5149–5160. [https://doi.org/https://doi.org/10.1016/S0009-2509\(00\)00139-1](https://doi.org/https://doi.org/10.1016/S0009-2509(00)00139-1)

Goel, A., Pothineni, N. V., Singhal, M., Paydak, H., Saldeen, T., & Mehta, J. L. (2018). Fish, Fish Oils and Cardioprotection: Promise or Fish Tale? *International Journal of Molecular Sciences*, 19(12), 3703. <https://doi.org/10.3390/IJMS19123703>

Golmakani, M. T., Rezaei, K., & Mazidi, S. (2014). Optimization of the supercritical fluid extraction of lipids from marine microorganisms using response surface methodology. *The Journal of Supercritical Fluids*, 92, 263–269. <https://doi.org/10.1016/j.supflu.2014.06.014>

Gomes, A., Costa, A.L.R., Cardoso, D.D., Náthia-Neves, G., Meireles, M.A.A., Cunha, R.L., 2021. Interactions of β -carotene with WPI/*Tween* 80 mixture and oil phase: Effect on the behavior of O/W emulsions during in vitro digestion. *Food Chem.* 341, 128155. <https://doi.org/10.1016/j.foodchem.2020.128155>

Gorji SG, Smyth HE, Sharma M, Fitzgerald M. 2016. Lipid oxidation in mayonnaise and the role of natural antioxidants: a review. *Trend in Food Sci. Technol.*,56, 88-102.

Goswami, M. J., Dutta, U., & Kakati, D. (2024). *Ultrasound-Assisted Extraction for Food, Pharmacy, and Biotech Industries* (pp. 103–128). Springer Nature. https://doi.org/10.1007/978-1-0716-3601-5_5

Guil-Guerrero, J.L., Venegas-Venegas, E., Rincón-Cervera, M.Á., Suárez, M.D., 2011. Fatty acid profiles of livers from selected marine fish species. *J. Food Compos. Anal.* 24, 217–222. <https://doi.org/10.1016/j.jfca.2010.07.011>.

Gulzar, S., & Benjakul, S. (2020). Impact of pretreatment and atmosphere on quality of lipids extracted from cephalothorax of Pacific white shrimp by ultrasonic assisted process. *Food Chemistry*, 309, 125732. <https://doi.org/https://doi.org/10.1016/j.foodchem.2019.125732>

Gupta, A., Eral, H. B., Hatton, T. A., & Doyle, P. S. (2016). Nanoemulsions: formation, properties, and applications. *Soft Matter*, 12(11), 2826-2841.

- Hafiludin, 2015. Analisis Kandungan Gizi Pada Ikan Bandeng Yang Berasal Dari Habitat Yang Berbeda. *Jurnal Kelautan*, 8: 1
- Hamed, S., Soliman, T., Hassan, L., & Abo-Elwafa, G. (2019). Preparation of Functional Yogurt Fortified with Fish Oil-In-Water Nanoemulsion. *Egyptian Journal of Chemistry*, 62(1), 6–7. <https://doi.org/10.21608/ejchem.2019.18621.2149>
- Hamzah, N. S. A., Mohamad, A., & Maskat, M. Y. (2024). Ultrasound-assisted extraction of fish oil: Effect on fatty acid profile and quality. *Marine Drugs*, 22(1), Article 15. <https://doi.org/10.3390/md22010015>
- Hashemi, H., Christensen, J., Gersen, S., & Glarborg, P. (2014). Hydrogen oxidation at high pressure and intermediate temperatures: Experiments and kinetic modeling. *Proceedings of the Combustion Institute*, 35. <https://doi.org/10.1016/j.proci.2014.05.101>
- Hashim, N. A., Mazilan, M. S. R., Man, R. C., Arshad, Z. I. M., & Mudalip, S. K. A. (2022). Recovery of omega-3 fish oil from *Monopterus albus* using microwave assisted extraction process. *AIP Conference Proceedings*, 2610(1), 60015. <https://doi.org/10.1063/5.0099547>
- Hassanzadeh, H., Rahbari, M., Galali, Y., Hosseini, M., & Ghanbarzadeh, B. (2023). The garlic extract-loaded nanoemulsion: Study of physicochemical, rheological, and antimicrobial properties and its application in mayonnaise. *Food Science and Nutrition*, 11(7), 3799–3810. <https://doi.org/10.1002/fsn3.3365>
- Helland IB, Smith L, Saarem K, Saugstad OD, Drevon CA. 2003. Maternal supplementation with very-long-chain n-3 fatty acids during pregnancy and lactation augments children's IQ at 4 years of age. *Pediatrics*.;111(1):e39-e44
- Herh, P., Colo, S., Roye, N., & Hedman, K. (2000). Rheological of foods: New techniques, capabilities and instruments. *American Laboratory*, 32, 16–20.
- Hernandez-Hernandez, O., Ferreira-Lazarte, A., Tiwari, B., & Villamiel, M. (2024). Update of high-intensity ultrasound applications for the extraction of pectin from agri-food by-products. *TrAC - Trends in Analytical Chemistry*, 175(April), 117728. <https://doi.org/10.1016/j.trac.2024.117728>
- Hidayah, N., Rohman, A., Mustafidah, M., & Irnawati. (2022). Physicochemical characterization and fatty acid profiles of fish oil from milkfish (*Chanos*

chanos F.). Food Research, 6(2), 265–270.
[https://doi.org/10.26656/fr.2017.6\(2\).222](https://doi.org/10.26656/fr.2017.6(2).222)

Holub, D. J., & Holub, B. J. (2004). Omega-3 fatty acids from fish oils and cardiovascular disease. *Molecular and Cellular Biochemistry*, 263(1), 217–225. <https://doi.org/10.1023/B:MCBI.0000041863.11248.8D>

Honary, S., & Zahir, F. (2013). Effect of zeta potential on the properties of nano-drug delivery systems - A review. *Tropical Journal of Pharmaceutical Research*, 12(2), 255-264. <https://doi.org/10.1016/j.foodchem.2006.08.009>.
<https://doi.org/10.1365/s10337-008-0696-1>

Iberahim, N., Sulaiman, R., & Ibrahim, N. (2020). Optimization of omega-3 fish oil extraction from salmon using hexane-isopropanol solvent system via response surface methodology. *Food Research*, 4(5), 1535–1543. [https://doi.org/10.26656/fr.2017.4\(5\).538](https://doi.org/10.26656/fr.2017.4(5).538)

Ikhsan, A. N., Syifa, F., Mustafidah, M., & Rohman, A. (2021). Implementation of chemometrics as a Solution to detecting and preventing falsification of herbal medicines in Southeast Asia: A review. *Journal of Applied Pharmaceutical Science*, 11(9), 139–148. <https://doi.org/10.7324/JAPS.2021.110917>

Ismail, S., Ahmed-Farid, O. A., Metwally, G. F., Mansour, D., & Khattab, A. (2024). Anti-aging effect of mustard and flax seed oils based nanoemulsion gel on aged rat skin through antioxidant and anti-inflammatory activity. *Scientific African*, 26, e02369. <https://doi.org/10.1016/j.sciaf.2024.e02369>

Jacobsen, C. (2010). Enrichment of foods with omega-3 fatty acids: a multidisciplinary challenge. *Annals of the New York Academy of Sciences*, 1190(1), 141–150. <https://doi.org/https://doi.org/10.1111/j.1749-6632.2009.05263.x>

Jafari, S., & McClements, D. J. (Eds.). (2018). *Nanoemulsions: Formulation, Applications, and Characterization*. Academic Press. <https://doi.org/10.1016/B978-0-12-811838-2.00001-1>

Jaiswal, M., Dudhe, R., & Sharma, P. K. (2015). Nanoemulsion: an advanced mode of drug delivery system. *Biotech*, 5(2), 123-127.

Jamshidi, A., Cao, H., Xiao, J., & Simal-Gandara, J. (2020). Advantages of techniques to fortify food products with the benefits of fish oil. *Food Research International*, 109353. <https://doi.org/10.1016/j.foodres.2020.109353>

- Jasmina, H., Orman, D., Elezović, A., Vranić, E., & Rahić, O. (2017). Preparation of nanoemulsions by high-energy and lowenergyemulsification methods (pp. 317–322). Springer, Singapore. https://doi.org/10.1007/978-981-10-4166-2_48
- Ji, H. (2006). Refining Process of Tuna Oil and Its Effect on Fatty Acid Composition. *Journal of Zhanjiang Ocean University*.
- Jo, Y., & Kwon, Y. (2014). Characterization of β -carotene nanoemulsions prepared by microfluidization technique. *Food Science and Biotechnology*, 23(1), 107–113.
- Julio, L. M., Copado, C. N., Diehl, B. W. K., Tomás, M. C., & Y., I. (2024). Development of chia oil-in-water nanoemulsions using different homogenization technologies and the layer-by-layer technique. *Exploration of Foods and Foodomics*, 2(2), 107–124. <https://doi.org/10.37349/eff.2024.00029>
- Kale, S. N., & Deore, S. L. (2017). Emulsion, microemulsion, and nanoemulsion: A review. *Systematic Reviews in Pharmacy*, 8(1), 39–47. <https://doi.org/10.5530/srp.2017.1.8>
- Karupaiah, T., Chuah, K.-A., Chinna, K., Matsuoka, R., Masuda, Y., Sundram, K., & Sugano, M. (2016). Comparing effects of soybean oil- and palm olein-based mayonnaise consumption on the plasma lipid and lipoprotein profiles in human subjects: a double-blind randomized controlled trial with cross-over design. *Lipids in Health and Disease*, 15(1), 131. <https://doi.org/10.1186/s12944-016-0301-9>
- Kementerian Kelautan dan Perikanan. (2023). Produksi ikan bandeng di Indonesia 2023. Diakses dari <https://portaldata.kkp.go.id>.
- Khan, M. K., Abert-Vian, M., Fabiano-Tixier, A. S., Dangles, O., & Chemat, F. (2010). Ultrasound-assisted extraction of polyphenols (flavanone glycosides) from orange (*Citrus sinensis* L.) peel. *Food Chemistry*, 119(2), 851–858.
- Kolanowski, W., & Laufenberg, G. (2006). Enrichment of food products with polyunsaturated fatty acids by fish oil addition. *European Food Research Technology*, 222, 472-477
- Komaiko, J. S., & McClements, D. J. (2016). Formation of Food-Grade Nanoemulsions Using Low-Energy Preparation Methods: A Review of Available Methods. *Comprehensive reviews in food science and food safety*, 15(2), 331–352. <https://doi.org/10.1111/1541-4337.12189>

- Kosasih, W., Rosmalina, R. T., Risdian, C., Priatni, S., & Saepudin, E. (2021). Production of omega-3 fatty acids by enzymatic hydrolysis from lemuru fish by-products. *Sains Malaysiana*, 50(8), 2271–2282. <https://doi.org/10.17576/jsm-2021-5008-11>
- Kris-Etherton, P. M., Grieger, J. A., & Etherton, T. D. (2009). Dietary reference intakes for DHA and EPA. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 81(2), 99–104. <https://doi.org/https://doi.org/10.1016/j.plefa.2009.05.011>
- Kumar, A., Lindley, M., & Mastana, S. (2014). A time efficient adaptation of GC-FID method for the analysis of PBMC lipid composition. *Journal of Biochemical Technology*, 5, 760–764.
- Laca, A., M. C. Sáenz, B. Paredes, & M. Díaz. 2010. Rheological properties, stability and sensory evaluation of low-cholesterol mayonnaises prepared using egg yolk granules as emulsifying agent. *Journal of Food Engineering* 97:243-252
- Larqué, E., Zamora, S., & Gil, A. (2001). Dietary trans fatty acids in early life: a review. *Early Human Development*, 65 Suppl, S31-41. [https://doi.org/10.1016/s0378-3782\(01\)00201-8](https://doi.org/10.1016/s0378-3782(01)00201-8)
- Lee, S. J., & McClements, D. J. (2010). Fabrication of protein-stabilized nanoemulsions using a combined homogenization and amphiphilic solvent dissolution/evaporation approach. *Food Hydrocolloids*, 24(6–7), 560–569. <https://doi.org/10.1016/j.foodhyd.2010.02.002>
- Li, A., Ha, Y., Wang, F., Li, W., & Li, Q. (2012). Determination of thermally induced trans fatty acids in soybean oil by attenuated total reflectance Fourier transform infrared spectroscopy and gas chromatography analysis. *Journal of Agricultural and Food Chemistry*, 60(42), 10709–10713. <https://doi.org/10.1021/jf3033599>
- Li, X., Yang, X., Deng, H., Guo, Y., & Xue, J. (2020). Gelatin films incorporated with thymol nanoemulsions: Physical properties and antimicrobial activities. *International Journal of Biological Macromolecules*, 150, 161–168. <https://doi.org/https://doi.org/10.1016/j.ijbiomac.2020.02.066>
- Lilian, L., & Salam, Z. (2022). Optimization of soybean oil extraction using d-limonene as a green solvent. *International Journal of Food Engineering and Technology*, 6(2), 42–48.

- Liu, F., Zhu, Z., Ma, C., Luo, X., Bai, L., Decker, E. A., Gao, Y., & McClements, D. J. (2016). Fabrication of concentrated fish oil emulsions using dual-channel microfluidization: Impact of droplet concentration on physical properties and lipid oxidation. *Journal of Agricultural and Food Chemistry*, 64(50), 9532–9541. <https://doi.org/10.1021/acs.jafc.6b04413>
- Liu, S. X., & Mamidipally, P. K. (2005). Quality Comparison of Rice Bran Oil Extracted with d-limonenand Hexane. *Cereal Chemistry*, 82(2), 209–215. <https://doi.org/https://doi.org/10.1094/CC-82-0209>
- LK Low and CS Ng. Preparation of methyl esters by Boron Trifluoride Method. In: H Hasegawa (Ed.). *Laboratory Manual on Analytical Methods and Procedures for Fish and Fish Products*. Marine Fisheries Research Department, Southeast Asian Fisheries Development Center, Singapore, 1987, p.C-11.1-C-11.2.
- Luz, A. M., Barbosa, G. D., Manske, C., & Tavares, F. W. (2023). Tween-80 on Water/Oil Interface: Structure and Interfacial Tension by Molecular Dynamics Simulations. *Langmuir*, 39(9), 3255–3265. <https://doi.org/10.1021/acs.langmuir.2c03001>
- Mahmud, J., Muranyi, P., Shankar, S., Sarmast, E., Salmieri, S., & Lacroix, M. (2024). Physiological and antimicrobial properties of a novel nanoemulsion formulation containing mixed surfactant and essential oils: Optimization modeling by response surface methodology. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 686. dis133405. <https://doi.org/10.1016/j.colsurfa.2024.133405>
- Marhamati, M., Kakhaki, K. K., & Rezaie, Z. (2020). Advance in Ultrasound-Assisted Extraction of Edible Oils: A Review. *J Nutr Fast Health*, 8(4), 220–230. <https://doi.org/10.22038/JNFH.2020.51138.1288>.
- Mason, T. G., Wilking, J. N., Meleson, K., Chang, C. B., & Graves, S. M. (2006). Nanoemulsions: Formation, structure, and physical properties. *Journal of Physics: Condensed Matter*, 18(41), R635-R666.
- McClements, D. J. (2012). Nanoemulsions versus microemulsions: Terminology, differences, and similarities. *Soft Matter*, 8(6), 1719-1729.
- McClements, D. J. (2015). *Food emulsions: Principles, practices, and techniques* (3rd ed.). CRC Press.

- McClements, D. J., Decker, E. A., & Weiss, J. (2007). Emulsion-based delivery systems for lipophilic bioactive components. *Journal of food science*, 72(8), R109–R124. <https://doi.org/10.1111/j.1750-3841.2007.00507.x>
- Medina-Torres, N., Ayora-Talavera, T., Espinosa-Andrews, H., Sanchez-Contreras, A., Pacheco, N., 2017. Ultrasound assisted extraction for the recovery of phenolic compounds from vegetable sources. *Agronomy* 7 (3), 47.
- Mehmood, T., Ahmed, A., Ahmed, Z., Ahmad, M.S., 2019. Optimization of soya lecithin and *Tween* 80 based novel vitamin D nanoemulsions prepared by ultrasonication using response surface methodology. *Food Chem.* 289, 664–670. <https://doi.org/10.1016/j.foodchem.2019.03.112>
- Mendoza, L., and Taruc, N. 2003. Microbial fermentation of milkfish solid wastes. Terminal Report of Research on Utilization of Full Grown Milkfish, Philippine Council for Industry and Energy Research and Development, UP Visayas, Miagao, Iloilo, Philippines
- Menegazzo, M. L., Petenuci, M. E., & Fonseca, G. G. (2014). Production and characterization of crude and refined oils obtained from the co-products of Nile tilapia and hybrid sorubim processing. *Food Chemistry*, 157, 100–104. <https://doi.org/10.1016/j.foodchem.2014.01.121>
- Menezes, R. C. F. de, Gomes, Q. C. de C., Almeida, B. S. de, Matos, M. F. R. de, & Pinto, L. C. (2022). Plant-based mayonnaise: Trending ingredients for innovative products. *International Journal of Gastronomy and Food Science*, 30, 100599. <https://doi.org/https://doi.org/10.1016/j.ijgfs.2022.100599>
- Merkle, S., Giese, E., Rohn, S., Karl, H., Lehmann, I., Wohltmann, A., & Fritsche, J. (2017). Impact of fish species and processing technology on minor fish oil components. *Food Control*, 73, 1379–1387. <https://doi.org/10.1016/j.foodcont.2016.11.003>
- Minasian, D., Pirouzifard, M., Almasi, H., & Hassanzadeh, H. (2024). Development of Prangos ferulacea essential oil based nanoemulsions/pickering emulsion and examination of their performance in mayonnaise shelf life. *Journal of Agriculture and Food Research*, 18, 101335. <https://doi.org/https://doi.org/10.1016/j.jafr.2024.101335>
- Misran, E., Xun, T. Y., Husin, F., Nor, N. F. A., Zubairi, S. I., & Yaakob, H. (2025). Optimization of Deodorization Parameters Using Liquid–Liquid Extraction from Edible Oil for High Antioxidants: Taguchi Optimization Approach

Technique. Food Analytical Methods, 0123456789.
<https://doi.org/10.1007/s12161-025-02816-9>

Mohamad said, K. anwar, & Afizal, M. (2016). Overview on the Response Surface Methodology (RSM) in Extraction Processes. *Journal of Applied Science & Process Engineering*, 2. <https://doi.org/10.33736/jaspe.161.2015>

Mohammadi, S., Alimi, M., Shahidi, S.-A., & Shokoohi, S. (2024). Investigating the physicochemical, rheological, and sensory properties of low-fat mayonnaise prepared with amaranth protein as an egg yolk replacer. *Food Science & Nutrition*, 12, 5147–5161. <https://doi.org/10.1002/fsn3.4163>

Mohammadpour, H., Sadrameli, S., Eslami, F., & Asoodeh, A. (2019). Optimization of ultrasound-assisted extraction of *Moringa peregrina* oil with response surface methodology and comparison with Soxhlet method. *Industrial Crops and Products*, 131. <https://doi.org/10.1016/j.indcrop.2019.01.030>

Morley, W. G. (2016). Mayonnaise (B. Caballero, P. M. Finglas, & F. B. T.-E. of F. and H. Toldrá (eds.); pp. 669–676). Academic Press. <https://doi.org/https://doi.org/10.1016/B978-0-12-384947-2.00449-9>

Murthy, L., Padiyar, P., Madhusudana Rao, B., Asha, K., Jesmi, D., Girija, P., Prasad, M., & Ravishankar, C. (2018). Nutritional Profile and Heavy Metal Content of Cultured Milkfish (*Chanos chanos*). 53(April), 245–249.

Myers RH, Montgomery DC, Anderson-Cook CM. 2009. Response surface methodology: Process and product optimization using designed experiments. 3rd ed. John Wiley & Sons; Hoboken, NJ, USA

Nejadmansouri, M., Hosseini, S. M. H., Niakosari, M., Yousefi, G. H., & Golmakani, M. T. (2016). Physicochemical properties and storage stability of ultrasound-mediated WPI-stabilized fish oil nanoemulsions. *Food Hydrocolloids*, 61, 801–811. <https://doi.org/10.1016/j.foodhyd.2016.07.011>

Nethaji, D. K., & Parambil, K. A. (2018). Development and Applications of Nano emulsion in Food Technology. *International Journal of Scientific Research in Science and Technology*, 4(2), 46–49. <https://doi.org/10.32628/IJSRST18417>

Nichols, P. D., McManus, A., Krail, K., Sinclair, A. J., & Miller, M. (2014). Recent advances in omega-3: Health benefits, Sources, Products and bioavailability. *Nutrients*, 6(9), 3727–3733. <https://doi.org/10.3390/nu6093727>

- Nieva-Echevarría, B., Goicoechea, E., & Guillén, M. D. (2017). Polyunsaturated lipids and vitamin A oxidation during cod liver oil in vitro gastrointestinal digestion. Antioxidant effect of added BHT. *Food Chemistry*, 232, 733–743. <https://doi.org/10.1016/j.foodchem.2017.04.057>
- Nikzade, V., Tehrani, M., & Saadatmand, M. (2012). Optimization of low-cholesterol–low-fat mayonnaise formulation: Effect of using soy milk and some stabilizer by a mixture design approach. *Food Hydrocolloids*, 28, 344–352. <https://doi.org/10.1016/j.foodhyd.2011.12.023>
- Ningsih, Z., Lestari, M. L. A. D., & Maharin, S. A. R. (2021). Preparation and Characterization of Curcumin Nanoemulsion in Olive Oil-*Tween* 80 System using Wet Ball Milling Method. 1(1), 16. <https://doi.org/10.34311/ICSPC.2021.1.1.16>
- Nithya, S., Krishnan, R.R., Rao, N.R., Naik, K., Praveen, N., Vasantha, V.L. (2023). Microwave-Assisted Extraction of Phytochemicals. In: Cruz, J.N. (eds) *Drug Discovery and Design Using Natural Products*. Springer, Cham. https://doi.org/10.1007/978-3-031-35205-8_8
- Noriega-Rodríguez, J. A., Ortega-García, J., Angulo-Guerrero, O., García, H. S., Medina-Juárez, L. A., & Gámez-Meza, N. (2009). Oil production from sardine (*Sardinops sagax caerulea*) Producción de aceite a partir de sardina (*Sardinops sagax caerulea*). *CYTA - Journal of Food*, 7(3), 173–179. <https://doi.org/10.1080/19476330903010243>
- Norziah, M. H., Noor, F. A., & Osman, A. (2009). Extraction, composition and antioxidant activity of fish oil from the waste of sardine and mackerel. *International Journal of Food Science and Nutrition*, 60(S7), 106–113. <https://doi.org/10.1080/09637480903190599>
- Nursid, M., Permatasari, A., Syafitri, U. D., & Batubara, I. (2022). Application of Box-Behnken Design for the Extraction of *Padina australis*. *Molekul: Jurnal Ilmiah Kimia*, 17(2), 270. <https://doi.org/10.20884/1.jm.2022.17.2.6359>
- O'Brien, R.D., 2003. *Fats and Oils: Formulating and Processing for Applications*, 2nd Edition. ed. CRC Press, New York
- Ohnishi, H., & Saito, Y. (2013). Eicosapentaenoic acid (EPA) reduces cardiovascular events: relationship with the EPA/arachidonic acid ratio. *Journal of Atherosclerosis and Thrombosis*, 20(12), 861–877. <https://doi.org/10.5551/jat.18002>

- Ojha, K. S., Aznar, R., O'Donnell, C., & Tiwari, B. K. (2020). Ultrasound technology for the extraction of biologically active molecules from plant, animal and marine sources. *Trends in Analytical Chemistry*, 122. <https://doi.org/10.1016/j.trac.2019.115663>
- Olosho, A. I., Nair, S. K., Ambade, A. V., & Adekola, F. (2023). Eco-Friendly Extraction of Thevetia peruviana Oil: A Synergistic Approach of d-Limonene and Ultrasound Activation. <https://doi.org/10.1021/acsagscitech.3c00208>
- Özogul, Y., Özogul, F., Alagoz, S., 2007. Fatty Acid Profiles and Fat Contents of Commercially Important Seawater and Freshwater Fish Species of Turkey: A comparative study. *Food Chem.* 103, 217–223.
- Ozturk, B., & McClements, D. (2015). Progress in natural emulsifiers for utilization in food emulsions. *Current Opinion in Food Science*, 7. <https://doi.org/10.1016/j.cofs.2015.07.008>
- Pagliaro, M., Fabiano-Tixier, A. S., & Ciriminna, R. (2023). Limonene as a natural product extraction solvent. *Green Chemistry*, 25(16), 6108–6119. <https://doi.org/10.1039/d3gc02068a>
- Parker, J., Schellenberger, A. N., Roe, A. L., Oketch-Rabah, H., & Calderón, A. I. (2018). Therapeutic Perspectives on Chia Seed and Its Oil: A Review. *Planta Medica*, 84(9–10), 606–612. <https://doi.org/10.1055/a-0586-4711>
- Piorkowski DT, McClements DJ. 2014. Beverage emulsions: recent developments in formulation, 174 production, and applications. *Food Hydrocolloid* 5-41. DOI: 10.1016/j.foodhyd.2013.07.009.
- Prasetyaningrum, A., Widayat, W., Jos, B., Ratnawati, R., Riyanto, T., Prinanda, G. R., Le Monde, B. U., & Susanto, E. E. (2023). Optimization of Sequential Microwave-Ultrasonic-Assisted Extraction of Flavonoid Compounds from *Moringa oleifera*. *Trends in Sciences*, 20(1), 1–14. <https://doi.org/10.48048/tis.2023.6401>
- Purnomowati, I., Hidayati, D., dan Saparinto, C. 2007. *Ragam Olahan Bandeng Kanisius*. Yogyakarta.
- R.A., R., R.J., D., 2007. Sources of the very-long-chain unsaturated omega-3 fatty acids: Eicosapentaenoic acid and docosahexaenoic acid. *Curr. Opin. Clin. Nutr. Metab. Care* 10, 123–128.
- Rachmansyah, 2004. *Analisis Daya Dukung Lingkungan Perairan Teluk Awarange Kabupaten Barru, Sulawesi Selatan bagi Pengembangan Budidaya Bandeng*

dalam Keramba Jaring Apung. [Disertasi]. Bogor (ID):Institut Pertanian Bogor

Rahman, M. S., & Rahman, R. T. (2007). pH in Food Preservation (pp. 323–332). Routledge Handbooks Online. <https://doi.org/10.1201/9780429091483-23>

Rahman, N., Hashem, S., Akther, S., & Jothi, J. S. (2023). Impact of various extraction methods on fatty acid profile, physicochemical properties, and nutritional quality index of Pangus fish oil. *Food Science & Nutrition*, 11(8), 4688–4699. <https://doi.org/10.1002/fsn3.3431>

Rahmani-Manglano, N. E., González-Sánchez, I., García-Moreno, P. J., Espejo-Carpio, F. J., Jacobsen, C., & Guadix, E. M. (2020). Development of fish oil-loaded microcapsules containing whey protein hydrolysate as film-forming material for fortification of low-fat mayonnaise. *Foods*, 9(5). <https://doi.org/10.3390/foods9050545>

Rai, A. K., Swapna, H. C., Bhaskar, N., Halami, P. M., & Sachindra, N. M. (2010). Effect of fermentation ensilaging on recovery of oil from fresh water fish viscera. *Enzyme and Microbial Technology*, 46(1), 9–13. <https://doi.org/10.1016/j.enzmictec.2009.11.001>

Raikos, V., Hayes, H., & Ni, H. (2020). Aquafaba from commercially canned chickpeas as potential egg replacer for the development of vegan mayonnaise: recipe optimisation and storage stability. *International Journal of Food Science & Technology*, 55(5), 1935–1942. <https://doi.org/https://doi.org/10.1111/ijfs.14427>

Rajasekaran, B., Singh, A., Zhang, B., Hong, H., Prodpran, T., & Benjakul, S. (2024). Shrimp Oil-Enriched Mayonnaise Prepared Using Fish Myofibrillar Protein as a Substitute for Egg Yolk: Physical, Rheological, and Sensory Properties. *Colloids and Interfaces*, 8(2). <https://doi.org/10.3390/colloids8020022>

Rao, J. dan McClements, D.J. (2011) Formation of flavor oil microemulsions, nanoemulsions and emulsions: influence of composition and preparation method. *Journal of Agricultural and Food Chemistry* 59: 5026-5035

Rezvankhah, A., Emam-Djomeh, Z., Safari, M., Askari, G., & Salami, M. (2019). Microwave-assisted extraction of hempseed oil: studying and comparing of fatty acid composition, antioxidant activity, physiochemical and thermal properties with Soxhlet extraction. *Journal of Food Science and Technology*, 56(9), 4198–4210. <https://doi.org/10.1007/s13197-019-03890-8>

- Rhinard Valery Goethe Randang, & Pande Made Nova Armita Sari. (2023). Review Metode Deodorisasi Pada Minyak Ikan. *Journal Scientific Of Mandalika (JSM)* 2809-0543, 4(2), 23–27. <https://doi.org/10.36312/10.36312/vol4iss2pp23-27>
- Rodrigues, S., & Fernandes, F. A. N. (2017). Extraction processes assisted by ultrasound. In D. Bermúdez-Aguirre (Ed.), *Ultrasound: Advances for food processing and preservation* (pp. 351–368). Academic Press. <https://doi.org/10.1016/B978-0-12-804581-7.00014-2>
- Ruiz-Álvarez, J. M., del Castillo-Santaella, T., Maldonado-Valderrama, J., Guadix, A., Guadix, E. M., & García-Moreno, P. J. (2022). PH influences the interfacial properties of blue whiting (*M. poutassou*) and whey protein hydrolysates determining the physical stability of fish oil-in-water emulsions. *Food Hydrocolloids*, 122, 107075
- Ruxton, C. H. S., Reed, S. C., Simpson, M. J. A., & Millington, K. J. (2004). The health benefits of omega-3 polyunsaturated fatty acids: A review of the evidence. *Journal of Human Nutrition and Dietetics*, 17(5), 449–459. <https://doi.org/10.1111/j.1365-277X.2004.00552.x>
- Saberi, A.H., Fang, Y., dan McClements, D.J. 2013. Effect of glycerol on formation, stability, and properties of vitamin-E enriched nanoemulsions produced using spontaneous emulsification. *Journal of Colloid And Interface Science* 411: 105–113.
- Saleh, R., Fattah, N., Sahriawati, 2019. *Minyak Ikan Bandeng (Chanos chanos F)*, 1st ed, 1. Penerbit Samudra Biru (Anggota IKAPI).
- Salvia-Trujillo, L., McClements, D. J., & Martín-Belloso, O. (2021). Nanoemulsion design for the delivery of omega-3 fatty acids: formation, oxidative stability, and digestibility (pp. 295–319). Academic Press. <https://doi.org/10.1016/B978-0-12-821391-9.00016-8>
- Samaram, S., Mirhosseini, H., Tan, C. P., Ghazali, H. M., Bordbar, S., & Serjouie, A. (2015). Optimisation of ultrasound-assisted extraction of oil from papaya seed by response surface methodology: oil recovery, radical scavenging antioxidant activity, and oxidation stability. *Food Chemistry*, 172, 7–17. <https://doi.org/10.1016/j.foodchem.2014.08.068>
- Santos, M., Destura, I., & Ordoñez, J. (2014). Milkfish and Tilapia as Biofactories: Potentials and Opportunities. *Fish for the People*, 12, 21–31.

- Sarker, S. (2020). By-products of fish-oil refinery as potential substrates for biogas production in Norway: A preliminary study. *Results in Engineering*, 6, 100137. <https://doi.org/10.1016/j.rineng.2020.100137>
- Sasongko, H., Indah, I., Nurrochmad, A., Nugroho, A. E., & Rohman, A. (2025). Effects of wet rendering extraction on the fatty acid and physicochemical profiles of catfish (*Pangasius micronema* Blkr.), milkfish (*Chanos chanos* Forsskal.) and snakehead fish (*Chana striata* Bloch). *Food Research*, 9(1), 13–18. [https://doi.org/10.26656/fr.2017.9\(1\).518](https://doi.org/10.26656/fr.2017.9(1).518)
- Scurria, A., Fabiano Tixier, A. S., Lino, C., Pagliaro, M., D'Agostino, F., Avellone, G., Chemat, F., & Ciriminna, R. (2020). High Yields of Shrimp Oil Rich in Omega-3 and Natural Astaxanthin from Shrimp Waste. *ACS Omega*, 5(28), 17500–17505. <https://doi.org/10.1021/acsomega.0c01978>
- Sekhon, B. S. (2010). Food nanotechnology- an overview. *Nanotechnology: Science and Applications*, 3, 1–15. <https://doi.org/10.2147/NSA.S12947>
- Shahidi, F., & Ambigaipalan, P. (2018). Omega-3 Polyunsaturated Fatty Acids and Their Health Benefits. *Annual Review of Food Science and Technology*, 9(January), 345–381. <https://doi.org/10.1146/annurev-food-111317-095850>
- Shahidi, F., & Zhong, Y. (2010). Lipid oxidation and improving the oxidative stability. *Chemical Society reviews*, 39(11), 4067–4079. <https://doi.org/10.1039/b922183m>
- Sharif, H. R., Williams, P. A., Sharif, M. K., Abbas, S., Majeed, H., Masamba, K. G., Safdar, W., & Zhong, F. (2018). Current progress in the utilization of native and modified legume proteins as emulsifiers and encapsulants – A review. *Food Hydrocolloids*, 76, 2–16. <https://doi.org/10.1016/j.foodhyd.2017.01.002>
- Shen, L., Pang, S., Zhong, M., Sun, Y., Qayum, A., Liu, Y., Rashid, A., Xu, B., Liang, Q., Ma, H., & Ren, X. (2023). A comprehensive review of ultrasonic assisted extraction (UAE) for bioactive components: Principles, advantages, equipment, and combined technologies. *Ultrasonics Sonochemistry*, 106646. <https://doi.org/10.1016/j.ultsonch.2023.106646>
- Siddiqui, S. A., Pahmeyer, M. J., Assadpour, E., & Jafari, S. M. (2022). Extraction and purification of d-limonen from orange peel wastes: Recent advances. *Industrial Crops and Products*, 177(September 2021), 114484. <https://doi.org/10.1016/j.indcrop.2021.114484>

- Silva, H. D., Cerqueira, M. A., Souza, B. W. S., Ribeiro, C., Avides, M. C., Quintas, M. A. C., Coimbra, J. S. R., Carneiro-da-Cunha, M. G., & Vicente, A. A. (2011). Nanoemulsions of β -carotene using a high-energy emulsification–evaporation technique. *Journal of Food Engineering*, 102(2), 130–135. <https://doi.org/https://doi.org/10.1016/j.jfoodeng.2010.08.005>
- Silva, J. P. L. D. and B. D. G. D. M. Franco. 2012. Application of Oregano Essential Oil Against Salmonella Enteritidis in Mayonnaise Salad. *International Journal of Food Science and Nutrition Engineering*. 2 (5):70-75.
- Simat, V., Vlahovic, J., Soldo, B., Skroza, D., & Ljubenkovic, I. (2019). Production and refinement of omega-3 rich oils from processing by-products of farmed fish species. *Foods*, 16(8), 125. <https://doi.org/10.3390/foods8040125>
- Siswanti, Herawati, N. C. A., Anandito, R. B. K., Purnamayati, L., & Maghfira, L. L. (2023). The effect of catfish oil (*Pangasius* sp.) addition on physical and chemical characteristics of rice bran oil mayonnaise. *Food Research*, 7, 21–27. [https://doi.org/10.26656/fr.2017.7\(S3\).3](https://doi.org/10.26656/fr.2017.7(S3).3)
- Solans, C., & Sole, I. (2012). Nanoemulsions: Formation by low-energy methods. *Current Opinion in Colloid & Interface Science*, 17, 246–254. <https://doi.org/10.1016/j.cocis.2012.07.003>
- Sonam, Hema Chaudhary, Vimal Arora, Kanchan Kholi & Vikash Kumar (2013) Effect of Physicochemical Properties of Biodegradable Polymers on Nano Drug Delivery, *Polymer Reviews*, 53:4, 546-567
- Song, G., Zhang, M., Peng, X., Yu, X., Dai, Z., & Shen, Q. (2018). Effect of deodorization method on the chemical and nutritional properties of fish oil during refining. *LWT – Food Science and Technology*, 96, 560–567. <https://doi.org/10.1016/j.lwt.2018.06.004>
- Sudrajat, A., 2008. *Buku Budidaya 23 Komoditas Laut Menguntungkan*. Penebar Swadaya. Penebar Swadaya.
- Sun, M., Wang, C., Wang, H., & Tan, M. (2024). Fish oil-based bigels with outstanding sensory and antioxidant properties: Application in low-fat mayonnaise. *Food Hydrocolloids*, 159, 110673. <https://doi.org/https://doi.org/10.1016/j.foodhyd.2024.110>
- Sun, X., Pan, C., Ying, Z., Yu, D., Duan, X., Huang, F., Ling, J., Ouyang, X. kun, 2020. Stabilization of zein nanoparticles with k-carrageenan and *Tween* 80 for encapsulation of curcumin. *Int. J. Biol. Macromol.* 146, 549–559. <https://doi.org/10.1016/j.ijbiomac.2020.01.053>

- Suseno, S. H., Tajul, A. Y., Nadiah, W. A., & Noor, A. F. (2012). Improved color properties of sardinella lemuru oil during adsorbent refining using Magnesol XL. *International Food Research Journal*, 19(4), 1383–1386.
- Szpisják-Gulyás, N., Al-Tayawi, A. N., Horváth, Z. H., László, Z., Kertész, S., & Hodúr, C. (2023). Methods for experimental design, central composite design and the Box–Behnken design, to optimise operational parameters: A review. *Acta Alimentaria*, 52(4), 521–537. <https://doi.org/10.1556/066.2023.00235>
- Tadros, T., Izquierdo, P., Esquena, J., & Solans, C. (2004). Formation and stability of nanoemulsions. *Advances in Colloid and Interface Science*, 108-109, 303-318.
- Tak, J. W., Gupta, B., Thapa, R. K., Woo, K. B., Kim, S. Y., Go, T. G., Choi, Y., Choi, J. Y., Jeong, J.-H., Choi, H.-G., Yong, C. S., & Kim, J. O. (2017). Preparation and Optimization of Immediate Release/Sustained Release Bilayered Tablets of Loxoprofen Using Box-Behnken Design. *AAPS PharmSciTech*, 18(4), 1125–1134. <https://doi.org/10.1208/s12249-016-0580-5>
- Tanzi, C. D., Vian, M. A., Ginies, C., Elmaataoui, M., & Chemat, F. (2012). Terpenes as green solvents for extraction of oil from microalgae. *Molecules*, 17(7), 8196–8205. <https://doi.org/10.3390/molecules17078196>
- Tengku-Rozaina, T. M., & Birch, E. J. (2013). Physicochemical characterisation and oxidative stability of refined hoki oil, unrefined hoki oil and unrefined tuna oil. *International Journal of Food Science & Technology*, 48(11), 2331–2339. <https://doi.org/https://doi.org/10.1111/ijfs.12222>
- Topuz, O. K., Kaya, A., & Alp, A. C. (2016). Effect Of Extraction Variables On The Omega-3 Eicosapentaenoic Acid (Epa) Content Of (*Nannochloropsis Oculata*) Microalga Oil. *Scientific Bulletin. Series F. Biotechnologies*, XX, 172–177.
- Torrejon, C., Jung, U. J., & Deckelbaum, R. J. (2007). n-3 Fatty acids and cardiovascular disease: Actions and molecular mechanisms. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 77(5), 319–326. <https://doi.org/https://doi.org/10.1016/j.plefa.2007.10.014>
- Ullah, M. R., Rahman, M. A., Haque, M. N., Sharker, M. R., Islam, M. M., & Alam, M. A. (2022). Nutritional profiling of some selected commercially important freshwater and marine water fishes of Bangladesh. *Heliyon*, 8(10). <https://doi.org/10.1016/j.heliyon.2022.e10825>

- U.S. Department of Agriculture (USDA), and Department of Health and Human Services. (2010). Dietary guidelines for Americans 2010 (7th ed.). U.S. Government Printing Office.
- Vaisali, Ch., Charanyaa, S., Belur, P. D., & Regupathi, I. (2015). Refining of edible oils: A critical appraisal of current and potential technologies. *International Journal of Food Science & Technology*, 50, 13–23. <https://doi.org/10.1111/ijfs.12657>
- Valdivia-López, M. Á., & Tecante, A. (2015). Chapter Two - Chia (*Salvia hispanica*): A Review of Native Mexican Seed and its Nutritional and Functional Properties (J. B. T.-A. in F. and N. R. Henry (ed.); Vol. 75, pp. 53–75). Academic Press. <https://doi.org/https://doi.org/10.1016/bs.afnr.2015.06.002>
- van den Elsen, L. W. J., van Esch, B. C. A. M., Hofman, G. A., Kant, J., van de Heijning, B. J. M., Garssen, J., & Willemsen, L. E. M. (2013). Dietary long chain n-3 polyunsaturated fatty acids prevent allergic sensitization to cow's milk protein in mice. *Clinical and Experimental Allergy: Journal of the British Society for Allergy and Clinical Immunology*, 43(7), 798–810. <https://doi.org/10.1111/cea.12111>
- Vazquez-Espinosa, M., Velasco González de Peredo, A., Ferreiro-González, M., Carrera, C., Palma, M., Barbero, G., & Espada-Bellido, E. (2019). Assessment of Ultrasound Assisted Extraction as an Alternative Method for the Extraction of Anthocyanins and Total Phenolic Compounds from Maqui Berries (*Aristotelia chilensis* (Mol.) Stuntz). *Agronomy*, 9, 148. <https://doi.org/10.3390/agronomy9030148>
- Virost, M., Tomao, V., Ginies, C., & Chemat, F. (2008). Total Lipid Extraction of Food Using d-limonenes an Alternative to n-Hexane. *Chromatographia*, 68(3), 311–313. <https://doi.org/10.1365/s10337-008-0696-1>
- Walker, R., Decker, E. A., & McClements, D. J. (2015). Development of food-grade nanoemulsions and emulsions for delivery of omega-3 fatty acids: Opportunities and obstacles in the food industry. *Food and Function*, 6(1), 42–55. <https://doi.org/10.1039/c4fo00723a>
- Willett, S. A., & Akoh, C. C. (2019). Physicochemical characterization of organogels prepared from menhaden oil or structured lipid with phytosterol blend or sucrose stearate/ascorbyl palmitate blend. *Food & Function*, 10(1), 180–190. <https://doi.org/10.1039/C8FO01725E>.

- Worrasinchai, S., M. Suphantharika, S. Pinjai, & P. Jamnong, 2006. β -Glucan prepared from spent brewer's yeast as a fat replacer in mayonnaise, *Food Hydrocolloids*, 20, 68-78
- Xing, Z., Xu, Y., Feng, X., Gao, C., Wu, D., Cheng, W., Meng, L., Wang, Z., Xu, T., & Tang, X. (2024). Fabrication of cinnamon essential oil nanoemulsions with high antibacterial activities via microfluidization. *Food Chemistry*, 456(June), 1–10. <https://doi.org/10.1016/j.foodchem.2024.139969>
- Yesiltas, B., García-Moreno, P. J., Mikkelsen, R. K., Echers, S. G., Hansen, D. K., Greve-Poulsen, M., Hyldig, G., Hansen, E. B., & Jacobsen, C. (2023). Physical and Oxidative Stability of Emulsions Stabilized with Fractionated Potato Protein Hydrolysates Obtained from Starch Production Side Stream. *Antioxidants*, 12(8). <https://doi.org/10.3390/antiox12081622>
- Yesiltas, B., García-Moreno, P. J., Sørensen, A. D. M., Soria Caindec, A. M., Hyldig, G., Anankanbil, S., Guo, Z., & Jacobsen, C. (2021). Enrichment of mayonnaise with a high fat fish oil-in-water emulsion stabilized with modified DATEM C14 enhances oxidative stability. *Food Chemistry*, 341, 128141. <https://doi.org/10.1016/j.foodchem.2020.128141>
- Yuan, Y., Gao, Y., Zhao, J., & Mao, L. (2008). Characterization and stability evaluation of β -carotene nanoemulsions prepared by high pressure homogenization under various emulsifying conditions. *Food Research International*, 41(1), 61–68. <https://doi.org/10.1016/j.foodres.2007.09.006>
- Zahir, E., Saeed, R., Hameed, M. A., & Yousuf, A. (2017). Study of physicochemical properties of edible oil and evaluation of frying oil quality by Fourier transform-infrared (FT-IR) spectroscopy. *Arabian Journal of Chemistry*, 10, S3870–S3876. <https://doi.org/10.1016/j.arabjc.2014.05.025>
- Zhang, J., Jia, G., Wanbin, Z., Minghao, J., Wei, Y., Hao, J., Liu, X., Gan, Z., & Sun, A. (2021). Nanoencapsulation of zeaxanthin extracted from *Lycium barbarum* L. by complex coacervation with gelatin and CMC. *Food HydroColloids*, 112, 106280. <https://doi.org/10.1016/j.foodhyd.2020.106280>
- Zhang, L., Han, C., Liu, M., Yang, H., Zhang, F., & Meng, X. (2020). The formation, stability of DHA/EPA nanoemulsion prepared by emulsion phase inversion method and its application in apple juice. *Food Research International*, 133, 109132. <https://doi.org/10.1016/j.foodres.2020.109132>
- Zhang, X., Song, R., Liu, X., Xu, Y., & Wei, R. (2022). Fabrication of vitamin D3 nanoemulsions stabilized by Tween 80 and Span 80 as a composite surface-

active surfactant: Characterization and stability. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 645(February), 1–10.
<https://doi.org/10.1016/j.colsurfa.2022.128873>

Zhang, X., Wang, Y., & Wang, D. (2024). Synergistic stabilization of garlic essential oil nanoemulsions by carboxymethyl chitosan/Tween 80 and application for coating preservation of chilled fresh pork. *International Journal of Biological Macromolecules*, 131370.
<https://doi.org/10.1016/j.ijbiomac.2024.131370>

Zhao, C., Jiang, G., Zhao, J., & Tian, D. (2012). Extraction and fatty acid composition analysis of Antarctic krill oil]. *Food and Fermentation Industries*, 38(3), 182–185. <https://doi.org/10.13386/j.issn1002-0306.2012.03.028>

Zuta, C.P., Simpson, B.K., Chan, H.M., Phillips, L., 2003. Concentrating PUFA from Mackerel Processing Waste. *JAOCs, J. Am. Oil Chem. Soc.* 80, 933–936. <https://doi.org/10.1007/s11746-003-0799-5>.