

DAFTAR PUSTAKA

- Acton, R., Barton, J., Snively, B., McLaren, C., Adams, P., Harris, E., *et al.* (2006). Geographic and racial/ethnic differences in HFE mutation frequencies in the Hemochromatosis and Iron Overload Screening (HEIRS) Study. *Ethnicity & disease*, 16 : 815–21.
- Aghamohammadi, A., Maghsoodlu, M., Teimori Naghadeh, H. and Ferdowsi, S. (2017). Iron stores in blood donors: A literature mini review. *Journal of Basic Research in Medical Sciences*, 4(4) : 50–55.
- Akpotuzor, J.O., Isong, C., Okpokam, D.C. and Etukudo, M.H. (2008). Levels of serum iron, total iron binding capacity, transferrin saturation fraction and packed cell volume of blood donors in Calabar, Cross River State, Nigeria. *Pakistan Journal of Nutrition*, 7(3) : 500–502.
- Alageeli, A.A., Alqahtany, F.S. and Algahtani, F.H. (2021). The Role of Reticulocyte Hemoglobin Content for the Diagnosis of Functional Iron Deficiency in Hemodialyzed patients. *Saudi Journal of Biological Sciences*, 28(1) : 50–54.
- Alaqzam, T.S., Stanley, A.C., Simpson, P.M., Flood, V.H. and Menon, S. (2018). Treatment Modalities in Adolescents Who Present with Heavy Menstrual Bleeding. *Journal of Pediatric and Adolescent Gynecology*, 31(5) : 451–458.
- Almashjary, M. N., Barefah, A., Bahashwan, S., Ashankyty, I., El-Fayoumi, R. I., Alzahrani, M. M., Assaqaf, D. M., Aljabri, R. S., Aljohani, A. Y., Muslim, R., Baawad, S. A., Bawazir, W. M. and Alharthy, S. A. (2022) Reticulocyte Hemoglobin-Equivalent Potentially Detects, Diagnoses and Discriminates between Stages of Iron Deficiency with High Sensitivity and Specificity, *Stomatology*, 11(19), p. 5675, [online] Available at: <https://www.mdpi.com/2077-0383/11/19/5675/pdf?version=1664193225>.
- Di Angelantonio, E., Thompson, S.G., Kaptoge, S., Moore, C., Walker, M., Armitage, J., *et al.* (2017). Efficiency and safety of varying the frequency of whole blood donation (INTERVAL): a randomised trial of 45 000 donors. *The Lancet*, 390(10110) : 2360–2371.
- Atipo-Tsiba, F.O.G., Mouyabi, E.Q.G., Angounda, B.M., Mokono, S.O., Gokaba, L.T.O. and Dokekias, A.E. (2023). Prevalence of Iron Deficiency, Anemia, and Associated Factors in a Blood Donor Population in Brazzaville, Congo. *Anemia*, 2023.
- Awotiwon, O.F., Cois, A., Pacella, R., Turawa, E.B., Dhansay, M.A., Stuijvenberg, L., *et al.* (2022). Estimating the changing disease burden attributable to iron deficiency in South Africa, 2000, 2006 and 2012. *South African medical journal = Suid-Afrikaanse tydskrif vir geneeskunde*, 112(8) : 684–692.
- Aziz Ali, S., Abbasi, Z., Feroz, A., Hambidge, K.M., Krebs, N.F., Westcott, J.E., *et al.* (2019). Factors associated with anemia among women of the

reproductive age group in Thatta district: study protocol. *Reproductive Health*, 16(1) : 34.

- Badr, M., Elashkar, S., Ismail, W., Abozid, A. and Hanna, D. (2023) Role of reticulocyte haemoglobin content in diagnosis of iron deficiency anaemia, *Alexandria Journal of Pediatrics*, 36, pp. 105–112.
- Beck, K.L., Conlon, C.A., Kruger, R., Heath, A.-L.M., Matthys, C., Coad, J., *et al.* (2014). Blood Donation, Being Asian, and a History of Iron Deficiency Are Stronger Predictors of Iron Deficiency than Dietary Patterns in Premenopausal Women. *BioMed Research International*, 2014 : 1–7.
- Beguin, Y. (2003). Soluble transferrin receptor for the evaluation of erythropoiesis and iron status. *Clinica Chimica Acta*, 329(1) : 9–22.
- Bialkowski, W., Bryant, B.J., Schlumpf, K.S., Wright, D.J., Birch, R., Kiss, J.E., *et al.* (2015). The strategies to reduce iron deficiency in blood donors randomized trial: design, enrolment and early retention. *Vox sanguinis*, 108(2) : 178–185.
- Bouri, S. and Martin, J. (2018). Investigation of iron deficiency anaemia. *Clinical Medicine*, 18(3) : 242–244.
- Braga, F., Pasqualetti, S., Frusciante, E., Borrillo, F., Chibireva, M. and Panteghini, M. (2022). Harmonization Status of Serum Ferritin Measurements and Implications for Use as Marker of Iron-Related Disorders. *Clinical Chemistry*, 68(9) : 1202–1210.
- Brugnara, C. (2000). Reticulocyte cellular indices: A new approach in the diagnosis of anemias and monitoring of erythropoietic function. *Crit Rev Clin Lab Sci*, 37(2) : 93–130.
- Brugnara, C., Schiller, B. dan Moran, J. (2006). Reticulocyte hemoglobin equivalent (Ret He) and assessment of iron-deficient states. *Clin. Lab. Haem.*, 28 : 303–308.
- Cable, R.G., Birch, R.J., Spencer, B.R., Wright, D.J., Bialkowski, W., Kiss, J.E., *et al.* (2017). The operational implications of donor behaviors following enrollment in STRIDE (Strategies to Reduce Iron Deficiency in blood donors). *Transfusion*, 57(10) : 2440–2448.
- Cable, R.G., Brambilla, D., Glynn, S.A., Kleinman, S., Mast, A.E., Spencer, B.R., *et al.* (2016). Effect of iron supplementation on iron stores and total body iron after whole blood donation. *Transfusion*, 56(8) : 2005–2012.
- Cable, R.G., Glynn, S.A., Kiss, J.E., Mast, A.E., Steele, W.R., Murphy, E.L., *et al.* (2011). Iron deficiency in blood donors: analysis of enrollment data from the REDS-II Donor Iron Status Evaluation (RISE) study. *Transfusion*, 51(3) : 511–522.
- Cable, R.G., Glynn, S.A., Kiss, J.E., Mast, A.E., Steele, W.R., Murphy, E.L., *et al.* (2012). Iron deficiency in blood donors: the REDS-II Donor Iron Status Evaluation (RISE) study. *Transfusion*, 52(4) : 702–711.

- Cacoub, P., Vandewalle, C. and Peoc'h, K.(2019). Using transferrin saturation as a diagnostic criterion for iron deficiency: A systematic review. *Critical Reviews in Clinical Laboratory Sciences*, 56(8) : 526–532.
- Cai, J., Wu, M., Ren, J., Du, Y., Long, Z., Li, G., *et al.*(2017). Evaluation of the Efficiency of the Reticulocyte Hemoglobin Content on Diagnosis for Iron Deficiency Anemia in Chinese Adults. *Nutrients*, 9(5) : 450.
- Camaschella, C.(2019). Iron deficiency. *Blood*, 133(1) : 30–39.
- Chambers, J.C., Zhang, W., Li, Y., Sehmi, J., Wass, M.N., Zabaneh, D., *et al.*(2009). Genome-wide association study identifies variants in TMPRSS6 associated with hemoglobin levels. *Nature Genetics*, 41(11) : 1170–1172.
- Chaparro, C.M. and Suchdev, P.S.(2019). Anemia epidemiology, pathophysiology, and etiology in low- and middle-income countries. *Annals of the New York Academy of Sciences*, 1450(1) : 15–31.
- Charlebois, E. and Pantopoulos, K.(2021). Iron overload inhibits BMP/SMAD and IL-6/ STAT3 signaling to hepcidin in cultured hepatocytes. *PLoS ONE*, 16(6 June) : 1–12.
- Consort(2010). Correspondence CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. *Notes and Queries*, 8:18(274) : 1–9.
- Conway, D. and Henderson, M.A.(2019). Iron metabolism. *Anaesthesia and Intensive Care Medicine*, 20(3) : 175–177.
- Cook, J.D. and Skikne, B.S.(1982). Serum ferritin: a possible model for the assessment of nutrient stores. *The American Journal of Clinical Nutrition*, 35(5) : 1180–1185.
- Coskun, A.(2024). Diagnosis Based on Population Data versus Personalized Data: The Evolving Paradigm in Laboratory Medicine. *Diagnostics*, 14(19).
- Dara, R.C., Marwaha, N., Khetan, D. and Patidar, G.K.(2016a). A Randomized Control Study to Evaluate Effects of Short-term Oral Iron Supplementation in Regular Voluntary Blood Donors. *Indian Journal of Hematology and Blood Transfusion*, 32(3) : 299–306.
- De la Cruz-Góngora, V., Méndez-Gómez-Humarán, I., Gaona-Pineda, E.B., Shamah-Levy, T. and Dary, O.(2022). Drops of Capillary Blood Are Not Appropriate for Hemoglobin Measurement with Point-of-Care Devices: A Comparative Study Using Drop Capillary, Pooled Capillary, and Venous Blood Samples. *Nutrients*, 14(24).
- Dev, S. and Babitt, J.L.(2017). Overview of iron metabolism in health and disease: Iron metabolism in health and disease. *Hemodialysis International*, 21 : S6–S20.
- Diepeveen, L.E., Laarakkers, C.M.M., Martos, G., Pawlak, M.E., Uğuz, F.F., Verberne, K.E.S.A., *et al.*(2019). Provisional standardization of hepcidin assays: Creating a traceability chain with a primary reference material,

- candidate reference method and a commutable secondary reference material. *Clinical Chemistry and Laboratory Medicine*, 57(6) : 864–872.
- Dignass, A., Farrag, K. and Stein, J.(2018). Limitations of Serum Ferritin in Diagnosing Iron Deficiency in Inflammatory Conditions. *Int J Chronic Dis*, 1–11.
- El-Gendy, F.M., El-Hawy, M.A., Rizk, M.S., El-Hefnawy, S.M. and Mahmoud, M.Z.(2018). Value of Soluble Transferrin Receptors and sTfR/log Ferritin in the Diagnosis of Iron Deficiency Accompanied by Acute Infection. *Indian Journal of Hematology and Blood Transfusion*, 34(1) : 104–109.
- Elsayed, M.E., Sharif, M.U. and Stack, A.G.(2016). Transferrin Saturation. in *Advances in Clinical Chemistry*. Elsevier, 71–97.
- Fletcher, A., Forbes, A., Svenson, N. and Wayne Thomas, D.(2022). Guideline for the laboratory diagnosis of iron deficiency in adults (excluding pregnancy) and children. *British Journal of Haematology*, 196(3) : 523–529.
- Fung, Y.-L., Lee, C.-K., Kwon, S.-Y., Soedarmono, Y., Satake, M., Alcantara, R., et al.(2022). The Contribution and Potential of Older Blood Donors in Asia Pacific Regions. *Transfusion Medicine*, 71–76. Available at: <https://onlinelibrary.wiley.com/doi/epdf/10.1111/tme.12845>.
- Garcia-Casal, M.N., Pasricha, S.R., Martinez, R.X., Lopez-Perez, L. and Peña-Rosas, J.P.(2021). Serum or plasma ferritin concentration as an index of iron deficiency and overload. *Cochrane Database of Systematic Reviews*, 2021(5).
- Gattermann, N., Muckenthaler, M.U., Kulozik, A.E., Metzgeroth, G. and Hastka, J.(2021). The evaluation of iron deficiency and iron overload. *Deutsches Ärzteblatt international* [Preprint].
- Girelli, D., Nemeth, E. and Swinkels, D.W.(2016). Heparin in the diagnosis of iron disorders. *Blood*, 127(23) : 2809–2813.
- Goddard, A.F., James, M.W., McIntyre, A.S. and Scott, B.B.(2011). Guidelines for the management of iron deficiency anaemia. *Gut*, 60(10) : 1309–1316.
- Goldman, M., Uzicanin, S., Scalia, J., Scalia, V. and O'Brien, S.F.(2016). Impact of informing donors of low ferritin results. *Transfusion*, 56(9) : 2193–2198.
- Gonzo, M., Maramba, A. and Taylor, G.(2017). Heparin testing; establishing reference values for the Namibian blood donor population. *International Journal of Blood Transfusion and Immunohematology*, 7(1) : 1–6.
- Gupte, S., Mahida, V. and Bhatti, A.(2008). Iron status of regular voluntary blood donors. *Asian Journal of Transfusion Science*, 2(1) : 9.
- Harms, K. and Kaiser, T.(2015). Beyond soluble transferrin receptor: Old challenges and new horizons. *Best Practice and Research: Clinical Endocrinology and Metabolism*, 29(5) : 799–810.
- Hentze, M.W., Muckenthaler, M.U., Galy, B. and Camaschella, C.(2010). Two to

Tango: Regulation of Mammalian Iron Metabolism. *Cell*, 142(1) : 24–38.

Hoenemann, C., Ostendorf, N., Zarbock, A., Doll, D., Hagemann, O., Zimmermann, M., *et al.* (2021). Reticulocyte and Erythrocyte Hemoglobin Parameters for Iron Deficiency and Anemia Diagnostics in Patient Blood Management. A Narrative Review. *Journal of Clinical Medicine*, 10(18) : 4250.

Hu, W., Meng, H., Hu, Q., Feng, L. and Qu, X. (2019). Blood donation from 2006 to 2015 in Zhejiang Province, China: Annual consecutive cross-sectional studies. *BMJ Open*, 9(5) : 1–11.

Institute of Medicine (US) Panel on Micronutrients. (2001). *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*. Washington (DC): National Academies Press (US); 2001. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK222310/> doi: 10.17226/10026.

Jain, A., Chowdhury, N., Jain, S., Uttam, N. and Meinia, S.K. (2018). Altered Red Cell Indices in Repeat Blood Donors: Experience of a North Indian Blood Bank. *Indian Journal of Hematology and Blood Transfusion*, 34(4) : 666–670.

Karagülle, M., Gündüz, E., Mutlu, F.Ş. and Akay, M.O. (2013). Clinical Significance of Reticulocyte Hemoglobin Content in the Diagnosis of Iron Deficiency Anemia. *Turkish J Haematol*, 30(2) : 153–156.

Karregat, J.H.M., Meulenbeld, A., Quee, F.A., Novotny, V.M.J., de Haas, M., Swinkels, D., *et al.* (2025). Ferritin-guided iron supplementation in whole blood donors (FORTE): results of a double-blind randomized controlled trial. *medRxiv*, 2025.01.03.25319940. Available at: <http://medrxiv.org/content/early/2025/01/03/2025.01.03.25319940.abstract>

Kaur, Pandeep, Kaur, G., Kaur, Paramjit and Tahlan, A. (2021). Hcpidin as a diagnostic marker of iron deficiency in blood donors. *Transfusion and Apheresis Science*, 60(3) : 103121.

Keshelashvili, K., O'Meara, A., Stern, M., Jirout, Z., Pehlic, V., Holbro, A., *et al.* (2016). Adverse events and retention of donors of double red cell units by apheresis. *Blood Transfusion* [Preprint].

Kiss, J.E., Birch, R.J., Steele, W.R., Wright, D.J. and Cable, R.G. (2017). Quantification of body iron and iron absorption in the REDS-II Donor Iron Status Evaluation (RISE) study. *Transfusion*, 57(7) : 1656–1664.

Kiss, J.E., Brambilla, D., Glynn, S.A., Mast, A.E., Spencer, B.R., Stone, M., *et al.* (2015a). Oral iron supplementation after blood donation: A randomized clinical trial. *JAMA - Journal of the American Medical Association*, 313(6) : 575–583.

Kiss, J.E., Brambilla, D., Glynn, S.A., Mast, A.E., Spencer, B.R., Stone, M., *et*

- al.*(2015b). Oral Iron Supplementation After Blood Donation: A Randomized Clinical Trial. *JAMA*, 313(6) : 575.
- Kiss, J.E. and Vassallo, R.R.(2018). How do we manage iron deficiency after blood donation? *British Journal of Haematology*, 181(5) : 590–603.
- Lalosevic, M.S., Toncev, L., Stankovic, S., Dragasevic, S., Stojkovic, S., Jovicic, I., *et al.*(2020). Hepcidin is a reliable marker of iron deficiency anemia in newly diagnosed patients with inflammatory bowel disease. *Disease Markers*, 2020.
- Lam, J.R., Schneider, J.L., Quesenberry, C.P. and Corley, D.A.(2017). Proton Pump Inhibitor and Histamine-2 Receptor Antagonist Use and Iron Deficiency. *Gastroenterology*, 152(4) : 821-829.e1.
- Lankhorst, C.E. and Wish, J.B.(2010). Anemia in renal disease: Diagnosis and management. *Blood Reviews*, 24(1) : 39–47.
- Lee, S.J., Min, H.K., Jang, J.S., Lee, S., Chung, Y. and KIM, M.J.(2020). Donor protection: Iron supplementation for frequent blood donors in Korea. *Transfusion and Apheresis Science*, 59(1) : 1–6.
- Levi, M., Rosselli, M., Simonetti, M., Brignoli, O., Cancian, M., Masotti, A., *et al.*(2016). Epidemiology of iron deficiency anaemia in four European countries: a population-based study in primary care. *European Journal of Haematology*, 97(6) : 583–593.
- Li, Y., Huang, X., Wang, J., Huang, R., Wan, D. and Yang, G.(2020). Regulation of Iron Homeostasis and Related Diseases. *Mediators of Inflammation*, 2020.
- Lin, X.M., Tian, W., Ma, L., Long, Z., Zhang, J., Shen, X.Y., *et al.*(2008). The responses of serum transferrin receptors to iron supplements in subjects with iron-deficiency erythropoiesis and iron-deficiency anaemia. *British Journal of Nutrition*, 99(2) : 416–420.
- Loria, F., Maret, E., Schobinger, C., Kuuranne, T. and Grabherr, S., Leuenberger, N.(2023). The effects of iron injection on blood doping biomarkers in dried blood spots. *Drug testing and analysis*, 15 (4) : 444–448. Available at: <https://analyticalsciencejournals.onlinelibrary.wiley.com/doi/epdf/10.1002/dta.3407>.
- Maghsudlu, M., Nasizadeh, S., Toogeh, G.R., Zandieh, T., Parandoush, S. and Rezayani, M.(2008). Short-term ferrous sulfate supplementation in female blood donors. *Transfusion*, 48(6) : 1192–1197.
- Magnussen, K. and Ladelund, S.(2015). Handling low hemoglobin and iron deficiency in a blood donor population: 2 years' experience. *Transfusion*, 55(10) : 2473–2478.
- Mantilla-Gutiérrez, C.Y. dan Cardona-Arias, J.A.(2012). Meta-analysis: prevalence of iron deficiency in systematic blood donors and its association with gender, 14.

- Mast, A.E.(2020). Reducing iron deficiency in teen-aged blood donors. *Pediatrics*, 146(1) : 1–2.
- Mast, A.E., Bialkowski, W., Bryant, B.J., Wright, D.J., Birch, R., Kiss, J.E., *et al.*(2016). A randomized, blinded, placebo-controlled trial of education and iron supplementation for mitigation of iron deficiency in regular blood donors. *Transfusion*, 56(6) : 1588–1597.
- Mast, A.E., Blinder, M.A., Gronowski, A.M., Chumley, C. and Scott, M.G.(1998). Clinical utility of the soluble transferrin receptor and comparison with serum ferritin in several populations. *Clinical Chemistry*, 44(1) : 45–51.
- Mathew, B.M., Santhakumar, R., Innah, S.J., Cherian, V.J. and Rafi, A.M.(2024). Evaluating RET-Hb vs Hb in Predicting Early Response in Oral Iron Therapy: A Prospective Interventional Study. *Journal of Clinical and Diagnostic Research*, 1–5.
- Mehta, S., Goyal, L., Kaushik, D., Gulati, S., Sharma, N., Harshvardhan, L., *et al.*(2017). Reticulocyte hemoglobin vis-à-vis immature reticulocyte fraction, as the earliest indicator of response to therapy in iron deficiency anemia. *Journal of Association of Physicians of India*, 65(October) : 14–17.
- Mirrezaie, S.M., Parsi, R., Torabgahromi, S.A. and Askarian, M.(2008). Low Dose, Short-Term Iron Supplementation in Female Blood Donors of Childbearing Age: a Randomized, Double-Masked, Placebo-Controlled Study, 6.
- Misje, A.H., Bosnes, V. and Heier, H.E.(2010). Gender differences in presentation rates, deferrals and return behaviour among Norwegian blood donors. *Vox Sanguinis*, 98(3p1) : e241–e248.
- Moretti, D., Goede, J.S., Zeder, C., Jiskra, M., Chatzinakou, V., Tjalsma, H., *et al.*(2015). Oral iron supplements increase hepcidin and decrease iron absorption from daily or twice-daily doses in iron-depleted young women. *Blood*, 126(17) : 1981–1989.
- Muckenthaler, M.U., Rivella, S., Hentze, M.W. and Galy, B.(2017). A Red Carpet for Iron Metabolism. *Cell*, 168(3) : 344–361.
- Nadarajan, V.S. and Eow, G.I.(2002). Anaemia and iron status among blood donors in a blood transfusion unit in Malaysia. *The Malaysian journal of pathology*, 24(2) : 99–102.
- Namaste, S.M., Rohner, F., Huang, J., Bhushan, N.L., Flores-Ayala, R., Kupka, R., *et al.*(2017). Adjusting ferritin concentrations for inflammation: Biomarkers Reflecting Inflammation and Nutritional Determinants of Anemia (BRINDA) project, 13.
- Napolitano, M., Dolce, A., Celenza, G., Grandone, E., Perilli, M.G., Siragusa, S., *et al.*(2014). Iron-dependent erythropoiesis in women with excessive menstrual blood losses and women with normal menses. *Annals of Hematology*, 93(4) : 557–563.
- Newman, B.H.(2004). Adjusting our management of female blood donors: the key

to an adequate blood supply. *Transfusion*, 44(4) : 591–596.

- Norashikin, J., Roshan, T.M., Rosline, H., Zaidah, A.W., Suhair, A.A. and Rapiaah, M.(2006). A study of serum ferritin levels among male blood donors in Hospital Universiti sains Malaysia. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 37(2) : 370–373.
- Nwagha, T.U., Ugwu, A.O. and Nwaekpe, C.N.(2023). Iron supplementation and blood donation in Nigeria: Effect on Hemoglobin, red cell indices, and iron stores - The ranferonTM study. *Annals of African Medicine*, 22(1) : 70–76.
- Pasricha, S.R., Marks, D.C., Salvin, H., Brama, T., Keller, A.J., Pink, J., *et al.*(2017). Postdonation iron replacement for maintaining iron stores in female whole blood donors in routine donor practice: results of two feasibility studies in Australia. *Transfusion*, 57(8) : 1922–1929.
- Pasricha, S.R., Tye-Din, J., Muckenthaler, M.U. and Swinkels, D.W.(2021). Iron deficiency. *The Lancet*, 397(10270) : 233–248.
- Pehlic, V., Volken, T., Holbro, A., Jirout, Z., Drexler, B., Buser, A., *et al.*(2021). Long-Term Course of Haemoglobin and Ferritin Values in High-Frequency Donors of Whole Blood and Double Erythrocyte Apheresis. *Transfusion Medicine and Hemotherapy*, 48(2) : 71–78.
- Pei, L.X., Kroeun, H., Vercauteren, S.M., Barr, S.I., Green, T.J., Albert, A.Y., *et al.*(2021). Baseline Hemoglobin, Hpcidin, Ferritin, and Total Body Iron Stores are Equally Strong Diagnostic Predictors of a Hemoglobin Response to 12 Weeks of Daily Iron Supplementation in Cambodian Women. *Journal of Nutrition*, 151(8) : 2255–2263.
- Penkova, M. and Ivanova, N.(2019). Serum Iron metabolism variables in clinically healthy persons. *Open Access Macedonian Journal of Medical Sciences*, 7(3) : 318–321.
- Petersen, P., Hakimjavadi, H., Chamala, S. and Mathur, G.(2024). Evaluating utility of routine ferritin testing in blood donors: A hospital-based blood donor centre experience. *Transfusion Medicine*, 34(6) : 491–498.
- Piskin, E., Ciansiosi, D., Gulec, S., Tomas, M. and Capanoglu, E.(2022). Iron Absorption: Factors, Limitations, and Improvement Methods. *ACS Omega*, 7(24) : 20441–20456.
- Pittori, C., Buser, A., Gasser, U.E., Sigle, J., Job, S., Rüesch, M., *et al.*(2011). A pilot Iron Substitution Programme in female blood donors with iron deficiency without anaemia: Iron substitution in non-anaemic premenopausal female blood donors. *Vox Sanguinis*, 100(3) : 303–311.
- Piva, E., Brugnara, C., Spolaore, F. and Plebani, M.(2014). Clinical Utility of Reticulocyte Parameters. *Clin Lab Med*, 35(1) : 133–163.
- Poggiali, E., Andreozzi, F., Nava, I., Consonni, D., Graziadei, G. and Cappellini, M.D.(2015). The role of TMPRSS6 polymorphisms in iron deficiency anemia partially responsive to oral iron treatment. *American Journal of*

Hematology, 90(4) : 306–309.

- Polin, V., Coriat, R., Perkins, G., Dhooze, M., Abitbol, V., Leblanc, S., *et al.* (2013). Iron deficiency: From diagnosis to treatment. *Digestive and Liver Disease*, 45(10) : 803–809.
- Powers, J.M., Buchanan, G.R., Adix, L., Zhang, S., Gao, A. and McCavit, T.L. (2017). Effect of Low-Dose Ferrous Sulfate vs Iron Polysaccharide Complex on Hemoglobin Concentration in Young Children With Nutritional Iron-Deficiency Anemia: A Randomized Clinical Trial. *JAMA*, 317(22) : 2297–2304.
- Prabhakar, E. (2022). Iron in Cell Metabolism and Disease. in M. Zakaria and T. Hassan (eds) *Biochemistry*. IntechOpen.
- Prados Madrona, D. and Fernández Herrera, M.D. (2014). Women as whole blood donors: offers, donations and deferrals in the province of Huelva, south-western Spain. *Blood Transfusion* [Preprint].
- Punnonen, K., Irjala, K. and Rajamäki, A. (1997). Serum Transferrin Receptor and Its Ratio to Serum Ferritin in the Diagnosis of Iron Deficiency. *Blood*, 89(3) : 1052–1057.
- Radtke, H., Tegtmeier, J., Röcker, L., Salama, A. and Kiesewetter, H. (2004). Daily doses of 20 mg of elemental iron compensate for iron loss in regular blood donors: a randomized, double-blind, placebo-controlled study: IRON SUPPLEMENTATION IN BLOOD DONORS. *Transfusion*, 44(10) : 1427–1432.
- Rigas, A.S., Sørensen, C.J., Pedersen, O.B., Petersen, M.S., Thørner, L.W., Kotzé, S., *et al.* (2014). Predictors of iron levels in 14,737 Danish blood donors: results from the Danish Blood Donor Study. *Transfusion*, 54(3 Pt 2) : 789–796.
- Ringwald, J., Zimmermann, R. and Eckstein, R. (2010). Keys to Open the Door for Blood Donors to Return. *Transfusion Medicine Reviews*, 24(4) : 295–304.
- Riskika, F., Briawan, D. and Tanzihah, I. (2023). Factors related to the level of iron adequacy of adolescent girls in Indonesia. *AcTion: Aceh Nutrition Journal*, 8(4) : 624.
- Røsvik, A.S., Hervig, T., Wentzel-Larsen, T. and Ulvik, R.J. (2010). Effect of iron supplementation on iron status during the first week after blood donation. *Vox Sanguinis*, 98(3 Pt 1) : e249-256.
- Sanyoto, A., Suega, K., Adnyana, L. and Bakta, I.M. (2017). Diagnostic Test Equivalent Hemoglobin Reticulocyte in Iron Deficiency Anemia. *The Indonesian Biomedical Journal*, 9(3) : 143.
- Schotten, N., Pasker-de Jong, P.C.M., Moretti, D., Zimmermann, M.B., Geurts-Moespot, A.J., Swinkels, D.W., *et al.* (2016). The donation interval of 56 days requires extension to 180 days for whole blood donors to recover from changes in iron metabolism. *Blood*, 128(17) : 2185–2188.

- Sethi, V., Yadav S., Agrawal, S., Sareen, N., Kathuria, N., Mishra, P., Kapoor, J., Dureja, S. (2019). Incidence of Side-effects After Weekly Iron and Folic Acid Consumption Among School-going Indian Adolescents. *Indian Pediatrics*, 56: 33-36.
- Singh, A., Chaudhary, R., Sonker, A. and Pandey, H.C.(2016). Importance of donor history of restless leg syndrome and pica to asses iron deficiency. *Transfusion and Apheresis Science*, 54(2) : 259–261.
- Singh, A., Pandey, H.Ch. and Chaudhary, R.(2022). Establishment of normal reference range of serum hepcidin in Indian blood donors, 17(1) : 1–6.
- Singh, D.S.(2018). Pattern and Trend of Female Donors in Blood Bank and Outdoor Blood Donation Camps - A 7 year Comparative Study from Blood Bank, Chamba, (H.P). *Journal of Medical Science And clinical Research*, 6(7) : 897–903.
- Skikne, B., Flowers, C. and Cook, J.(no date). Serum transferrin receptor: a quantitative measure of tissue iron deficiency, 9.
- Skikne, B.S., Flowers, C.H. and Cook, J.D.(1990). Serum Transferrin Receptor: A Quantitative Measure of Tissue Iron Deficiency. *Blood*, 75(9) : 1870–1876.
- Smith, G.A., Fisher, S.A., Doree, C., Di Angelantonio, E. and Roberts, D.J.(2014). Oral or parenteral iron supplementation to reduce deferral, iron deficiency and/or anaemia in blood donors. *Cochrane Database of Systematic Reviews*, 2014(7).
- Speeckaert, M.M., Speeckaert, R. and Delanghe, J.R.(2010). Biological and clinical aspects of soluble transferrin receptor. *Critical Reviews in Clinical Laboratory Sciences*, 47(5–6) : 213–228.
- Spencer, B.R., Johnson, B., Wright, D.J., Kleinman, S., Glynn, S.A. and Cable, R.G.(2016). Potential impact on blood availability and donor iron status of changes to donor hemoglobin cutoff and interdonation intervals: Blood Availability and Hb/Interval Changes. *Transfusion*, 56(8) : 1994–2004.
- Stoffel, N.U., Cercamondi, C.I., Brittenham, G., Zeder, C., Geurts-Moespot, A.J., Swinkels, D.W., *et al.*(2017). Iron absorption from oral iron supplements given on consecutive versus alternate days and as single morning doses versus twice-daily split dosing in iron-depleted women: two open-label, randomised controlled trials. *The Lancet Haematology*, 4(11) : e524–e533.
- Suria, N., Kaur, R., Mittal, K., Palta, A., Sood, T., Kaur, P. & Kaur, G. (2021). Utility of reticulocyte haemoglobin content and immature reticulocyte fraction in early diagnosis of latent iron deficiency in whole blood donors. *Vox Sanguinis*. [Online]. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/vox.13216>.
- Szymczyk-Nuzka, M. and Wołowicz, D.(2003). Iron stores in regular blood donors]. *Polskie Archiwum Medycyny Wewnętrznej*, 110(6) : 1415–1421.
- Tiwari, A. K., Bhardwaj, G., Arora, D., Aggarwal, G., Pabbi, S., Dara, R. C.,

- Sachdev, R., Raizada, A. and Sethi, M. (2018) Applying newer parameter Ret-He (reticulocyte haemoglobin equivalent) to assess latent iron deficiency (LID) in blood donors—study at a tertiary care hospital in India, *Vox Sanguinis*, Vox Sang, 113(7), pp. 639–646, [online] Available at: <https://pubmed.ncbi.nlm.nih.gov/30101488/>.
- Tolkien, Z., Stecher, L., Mander, A.P., Pereira, D.I.A. and Powell, J.J.(2015). Ferrous Sulfate Supplementation Causes Significant Gastrointestinal Side-Effects in Adults: A Systematic Review and Meta-Analysis. *PLOS ONE*. Edited by P. Strnad, 10(2) : e0117383.
- Trisna, G.(2016)*Profil Status Besi dan Kejadian Defisiensi Besi pada Donor Darah di Unit Donor Darah PMI Kota Yogyakarta*. Universitas Gadjah Mada.
- Triyono, T., Taslima, M. and Sukorini, U.(2023). Profile of female blood donors in the blood transfusion units of Dr. Sardjito Central General Hospital Yogyakarta. *Bali Medical Journal*, 12(1) : 274–277.
- Tuan, V.P., Anh, S.D., Van, P.H. and Sy, T.A.B.(2024). Iron supplementation post-blood donation in Vietnam: Impact on hemoglobin, red cell indices, and iron reserves. *Romanian Medical Journal*, 71(3) : 225–229.
- Urrechaga, E., Boveda, O., Aguayo, F.J., de la Hera, P., Muñoz, R.I., Gallardo, I., *et al.*(2016). Percentage of hypochromic erythrocytes and reticulocyte hemoglobin equivalent predictors of response to intravenous iron in hemodialysis patients. *International Journal of Laboratory Hematology*, 38(4) : 360–365.
- Utkurovich, N.A.(2021). An Overview of Iron Metabolism in Health and Disease. *Journal of Advances in Medicine and Medical Research*, 33(8) : 70–78.
- Vassallo, R.R., Hilton, J.F., Bravo, M.D., Vittinghoff, E., Custer, B. and Kamel, H.(2020). Recovery of iron stores after adolescents donate blood. *Pediatrics*, 146(1).
- Verga Falzacappa, M.V., Vujic Spasic, M., Kessler, R., Stolte, J., Hentze, M.W. and Muckenthaler, M.U.(2007). STAT3 mediates hepatic hepcidin expression and its inflammatory stimulation. *Blood*, 109(1) : 353–358.
- Vhanda, D., Chinowaita, F., Nkomo, S., Timire, C. and Kouamou, V.(2021). Effects of repeated blood donation on iron status of blood donors in Zimbabwe: A cross-sectional study. *Health Science Reports*, 4(4).
- Vogt, A.C.S., Arsiwala, T., Mohsen, M., Vogel, M., Manolova, V. and Bachmann, M.F.(2021). On iron metabolism and its regulation. *International Journal of Molecular Sciences*, 22(9) : 1–17.
- Vos, T., Abajobir, A.A., Abbafati, C., Abbas, K.M., Abate, K.H., Abd-Allah, F., *et al.*(2017). Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: A systematic analysis for the Global Burden of Disease Study 2016. *The Lancet*, 390(10100) : 1211–1259.

- Waldvogel, S., Rochat, B., Peduzzi, D., Vaucher, P., Tissot, J.D. and Favrat, B.(2016). Effects of oral supplementation of iron on hepcidin blood concentrations among non-anaemic female blood donors: A randomized controlled trial. *Vox Sanguinis*, 110(2) : 166–171.
- Westgard QC(2014). Biologic Variation Database, the 2014 Update - Westgard. Available at: <https://www.westgard.com/biodatabase-2014-update.htm>.
- World Health Organization(2017)*The 2016 global status report on blood safety and availability*. Geneva: World Health Organization.
- World Health Organization(2021)*Global status report on blood safety and availability*. Available at: <https://www.who.int/publications/i/item/9789240051683>.
- Xiao, X., Yeoh, B.S. and Vijay-Kumar, M.(2017). Lipocalin 2: An Emerging Player in Iron Homeostasis and Inflammation. *Annual Review of Nutrition*, 37(1) : 103–130.
- Yaranal, P.J. and S G, H.(2013). Evaluation of Iron Stores in Blood Donors by Estimation of Serum Ferritin level. *International Journal of Medical Research & Health Sciences*, 1(7) : 275–281.
- Yaznil, M.R., Lubis, M.P., Lumbanraja, S.N., Barus, M.N.G., Sakina, A., Sarirah, M. (2019). Side Effects of Daily versus Weekly Iron Supplementation in Pregnant Women. *J Bahrain Med Soc*. 31(1):17-22