

DAFTAR PUSTAKA

- Agyei-Amponsah, J., Macakova, L., DeKock, H. L., Emmambux, M. N. (2019). Sensory, tribological, and rheological profiling of “clean label” starch–lipid complexes as fat replacers. *Starch - Stärke*, 71(9) 1-9. <https://doi.org/10.1002/star.201800340>
- Ai, Y., Jovin Hasjim, J., & Jane, J.-L. (2013) Effects of lipids on enzymatic hydrolysis and physical properties of starch. *Carbohydrate Polymers*, 92 (1), 120-127. <https://doi.org/10.1016/j.carbpol.2012.08.092>
- Akbari, S., & Nour, A. H. (2018). Emulsion types, stability mechanisms and rheology: A review. *International Journal of Innovative Research and Scientific Studies*, 1(1), 14-21. <https://doi.org/10.53894/ijirss.v1i1.4>
- Aktaş, N., & Gerçekaslan, K. E. (2019). Effect of pregelatinization and retrogradation on some physicochemical changes of wheat-potato starches. *Journal of Agricultural Sciences*, 25(3), 281–289. <https://doi.org/10.15832/ankutbd.426252>
- Alcázar-Alay, S. C., & Meireles, M. A. A. (2015). Physicochemical properties, modifications and applications of starches from different botanical sources. *Food Science and Technology (Brazil)*, 35(2), 215-236. <https://doi.org/10.1590/1678-457X.6749>
- Ariviani, S., Anggrahini, S., Naruki, S., & Raharjo, S. (2015). Characterization and chemical stability evaluation of β -karoten microemulsions prepared by spontaneous emulsification method using VCO and palm oil as oil phase. *International Food Research Journal*, 22(6), 2432-2439.
- Aserin, A. (2007). Multiple Emulsions Technology And Applications. Wiley-Interscience. <https://doi.org/10.1002/smll.200990090>
- BeMiller, J., & Whistler, R. (2009). *Starch Chemistry and Technology* (3 ed.). Academic Press. Burlington, USA.
- Bernardi, D. S., Pereira, T. A., Maciel, N. R., Bortoloto, J., Viera, G. S., Oliveira, G. C., & Rocha-Filho, P. A. (2011). Formation and stability of oil-in-water nanoemulsions containing rice bran oil: In vitro and in vivo assessments. *Journal of Nanobiotechnology*, 9(1), 44-44. <https://doi.org/10.1186/1477-3155-9-44>
- Bhopatkar, D., Feng, T., Chen, F., Zhang, G., Carignano, M., Park, S. H., Zhuang, H., Campanella, O. H., & Hamaker, B. R. (2015). Self-assembled nanoparticle of common food constituents that carries a sparingly soluble small molecule. *Journal of Agricultural and Food Chemistry*, 63(17), 4312-4319.

- Bhosale, R. G., & Ziegler, G. R. (2010). Preparation of spherulites from amylose-palmitic acid complexes. *Carbohydrate Polymers*, 80(1), 53-64. <https://doi.org/10.1016/j.carbpol.2009.10.069>
- Blazek, J., & Gilbert, E. P. (2011). Application of small-angle X-ray and neutron scattering techniques to the characterisation of starch structure: A review. *Carbohydrate Polymers*, 85, 281–293. <https://doi.org/10.1016/j.carbpol.2011.02.041>
- Borba, C. M., Tavares, M. N., Macedo, L. P., Araújo, G. S., Furlong, E. B., C. L., & Burkert J. F. M. (2019). Physical and chemical stability of β -Carotene nanoemulsions during storage and thermal process. *Food Research International* 121, 229–237. <https://doi.org/10.1016/j.foodres.2019.03.045>
- Cabezas-Terán, K., Charlotte Grootaert, John Van Camp, Johana Ortiz, Jenny Ruales, Silvana Donoso, Tom Van de Wiele. (2025). Bioaccessibility of β -carotene during in vitro co-digestion of encapsulated mango peel carotenoids with milk. *Food Research International*, 214, 116576.
- Chao, C., Yu, J., Wang, S., Copeland, L., & Wang, S. (2018). Mechanisms Underlying the Formation of Complexes between Maize Starch and Lipids. *Journal of Agricultural and Food Chemistry*, 66(1), 272-278. <https://doi.org/10.1021/acs.jafc.7b05025>
- Chao, C., Huang, S., Yu, J., Copeland, L., Yang, Y., & Wang, S. (2024). The influence of short-range molecular order in gelatinized starch on the formation of starch-lauric acid complexes. *International Journal of Biological Macromolecules*, 260, 129526-129526. <https://doi.org/10.1016/j.ijbiomac.2024.129526>
- Chaves, M. A., & de Pinho, S. C. (2018). Effect of production parameters and stress conditions on beta-carotene-loaded lipid particles produced with palm stearin and whey protein isolate. *Brazilian Journal of Food Technology*, 21, 1-10. <http://dx.doi.org/10.1590/1981-6723.03517>
- Chen, B., McClements, D. J., & Decker, E. A. (2011). Minor Components in Food Oils: A Critical Review of their Roles on Lipid Oxidation Chemistry in Bulk Oils and Emulsions. *Critical Reviews in Food Science and Nutrition*, 51(10), 901–916. <https://doi.org/10.1080/10408398.2011.606379>
- Chen, X., He, X., Fu, X., Zhang, B., & Huang, Q. (2017a). Complexation of rice starch/flour and maize oil through heat moisture treatment: Structural, in vitro digestion and physicochemical properties. *International Journal of Biological Macromolecules*, 98, 557-564. <https://doi.org/10.1016/j.ijbiomac.2017.01.105>
- Chen, B., Zeng, S., Zeng, H., Guo, Z., Zhang, Y., & Zheng, B. (2017b). Properties of lotus seed starch–glycerin monostearin complexes formed by high pressure homogenization. *Food Chemistry*, 226, 119-127. <https://doi.org/10.1016/j.foodchem.2017.01.018>

- Chen, J., He, J., Zhao, Z., Li, X., Tang, J., Liu, Q., & Wang, H. (2023). Effect of heat treatment on the physical stability, interfacial composition and protein-lipid co-oxidation of whey protein isolate-stabilised O/W emulsions. *Food Research International*, 172, 113126. <https://doi.org/10.1016/j.foodres.2023.113126>
- Chen, H., Li, H., Liu, S., Wang, Z., & Kan, J. (2024). Insight into esterified and granular esterified-pregelatinized starch formation during esterification modification : Key role of temperature. *Food Chemistry*, 460(2), 140809. <https://doi.org/10.1016/j.foodchem.2024.140809>
- Chevalier, Y., & Bolzinger, M.-a. (2013). Emulsions stabilized with solid nanoparticles : Pickering emulsions. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 439, 23-34. <http://dx.doi.org/10.1016/j.colsurfa.2013.02.054>
- Chuacharoen, T., & Sabliov, C. M. (2019). Comparative effects of curcumin when delivered in a nanoemulsion or nanoparticle form for food applications: Study on stability and lipid oxidation inhibition. *LWT*, 113(2019) 108319. <https://doi.org/10.1016/j.lwt.2019.108319>
- De Pilli, T., Legrand, J., Derossi, A., & Severini, C. (2014). Effect of proteins on the formation of starch-lipid complexes during extrusion cooking of wheat flour with the addition of oleic acid. *International Journal of Food Science and Technology*, 50(2), 515-521. <https://doi.org/10.1111/ijfs.12698>
- Derycke, V., Vandeputte, G. E., Vermeylen, R., De Man, W., Goderis, B., Koch, M. H. J., & Delcour, J. A. (2005). Starch gelatinization and amylose-lipid interactions during rice parboiling investigated by temperature resolved wide angle X-ray scattering and differential scanning calorimetry. *Journal of Cereal Science*, 42(3), 334-343. <https://doi.org/10.1016/j.jcs.2005.05.002>
- Destribats, M., Gineste, S., Laurichesse, E., Tanner, H., Leal-Calderon, F., Héroguez, V., & Schmitt, V. (2014). Pickering emulsions: What are the main parameters determining the emulsion type and interfacial properties? *Langmuir*, 30(31), 9313-9326. <https://doi.org/10.1021/la501299u>
- Dewi, A. M. P. (2023). Sifat Fisikokimia Active Biodegradable Film Pati Sagu (Metroxylon sp) Termodifikasi Heat Moisture Treatment-Oktenil Suksinilasi Dengan Penambahan Ekstrak Kulit Kayu Akway (Drymis piperita Hook f.). *Disertasi*. Universitas Gadjah Mada. Yogyakarta.
- Dewi, A. M. P., Santoso, U., Pranoto, Y., & Marseno, D. W. (2022). Dual Modification of Sago Starch via Heat Moisture Treatment and Octenyl Succinylation to Improve Starch Hydrophobicity. *Polymers* 14, 1-17. <https://doi.org/10.3390/polym14061086>
- Dewi, A. M. P., Santoso, U., Pranoto, Y., & Marseno, D. W. (2023). Optimizing reaction condition of octenyl succinic anhydride on heat-moisture-treated sago starch and its application for biodegradable film. *Food Science and Technology*, 43, 1-9. <https://doi.org/10.5327/fst.17523>

- Dokić, L., Krstonošić, V., Nikolić, I. (2012). Physicochemical characteristics and stability of oil-in-water emulsions stabilized by OSA starch. *Food Hydrocolloids*, 29(1), 185-192. <https://doi.org/10.1016/j.foodhyd.2012.02.008>.
- dos Santos, T. P. R., Franco, C. M. L., Demiate, I. M., Li, X. H., Garcia, E. L., Jane, J. I., & Leonel, M. (2018). Spray-drying and extrusion processes: Effects on morphology and physicochemical characteristics of starches isolated from Peruvian carrot and cassava. *International Journal of Biological Macromolecules*, 118, 1346-1353. <https://doi.org/10.1016/j.ijbiomac.2018.06.070>
- Drosou C., & Krokida M. A. (2024) Comparative Study of Encapsulation of β -Carotene via Spray-Drying and Freeze-Drying Techniques Using Pullulan and Whey Protein Isolate as Wall Material. *Foods*. 13(12):1933. doi: 10.3390/foods13121933.
- D'Silva, T. V., Taylor, J. R.N., & Emmambux, M. N. (2011). Enhancement of the pasting properties of teff and maize starches through weteheat processing with added stearic acid. *Journal of Cereal Science*, 53(2), 192-197. <https://doi.org/10.1016/j.jcs.2010.12.002>
- Ehara, H., Toyoda, Y., & Johnson, D. V. (2018). Sago palm: Multiple contributions to food security and sustainable livelihoods. Springer Nature. Singapore. <https://doi.org/10.1007/978-981-10-5269-9>
- Fanta, G. F., Kenar, J. A., & Felker, F. C. (2015). Nanoparticle formation from amylose-fatty acid inclusion complexes prepared by steam jet cooking. *Industrial Crops and Products*, 74, 36-44. <https://doi.org/10.1016/j.indcrop.2015.04.046>
- Feng, T., Zhuang, H., Chen, F., Campanella, O., Bhotpakar, D., Carignano, M. A., & Park, S. H. (2018). Starch-lipid and starch-protein complexes and their application. In Z. Jin (Ed.), (pp. 1-226). Springer Nature. Singapore. <https://doi.org/10.1007/978-981-13-1077-5>
- Flores-Miranda, G. A., del Toro, G. V., & Yanez-Fernandez, J. (2015). Stability evaluation of karoten nanoemulsions prepared by homogenization-emulsification process using stearic acid as oil phase. *Revista Mexicana de Ingeniería Química*, 14(3), 667-680.
- Fu, Z.-Q., Wang, L.-J., Li, D., & Adhikari, B. (2012). Effects of partial gelatinization on structure and thermal properties of corn starch after spray drying. *Carbohydrate Polymers*, 88(4), 1319-1325. <https://doi.org/10.1016/j.carbpol.2012.02.010>
- Garcia, M. C., Pereira-Da-Silva, M. A., Taboga, S., & Franco, C. M. L. (2016). Structural characterization of complexes prepared with glycerol monoestearate and maize starches with different amylose contents. *Carbohydrate Polymers*, 148, 371-379. <https://doi.org/10.1016/j.carbpol.2016.04.067>

- Gelders, G. G., Vanderstukken, T. C., Goesaert, H., & Delcour, J. A. (2004). Amylose-lipid complexation: A new fractionation method. *Carbohydrate Polymers*, 56(4), 447-458. <https://doi.org/10.1016/j.carbpol.2004.03.012>
- Goderis, B., Putseys, J. A., Gommers, C. J., Bosmans, G. M., & Delcour, J. A. (2014). The structure and thermal stability of amylose–lipid complexes: a case study on amylose–glycerol monostearate. *Crystal Growth & Design*, 14(7), 3221–3233. <https://doi.org/10.1021/cg4016355>
- Goodarzi, F., & Zendehboudi, S. (2019). A Comprehensive Review on Emulsions and Emulsion Stability in Chemical and Energy Industries. *Canadian Journal of Chemical Engineering*, 97(1), 281-309. <https://doi.org/10.1002/cjce.23336>
- Guida, C., Aguiar, A. C., Magalhaes, A. E. R., Soares, M. G., & Cunha, R. L. (2024). Impact of ultrasound process on cassava starch nanoparticles and Pickering emulsions stability. *Food Research International*, 192, 114810. <https://doi.org/10.1016/j.foodres.2024.114810>
- Gul, K., Tak, A., Singh, A. K., Preeti Singh, P., Yousuf, B., and Wani, A. A. (2015). Chemistry, encapsulation, and health benefits of β -carotene - A review. *Cogent Food & Agriculture* (2015), 1: 1018696. <http://dx.doi.org/10.1080/23311932.2015.1018696>
- Hasanvand, E., Fathi, M. & Bassiri, A. (2018). Production and characterization of vitamin D3 loaded starch nanoparticles: effect of amylose to amylopectin ratio and sonication parameters. *Journal of Food Science and Technology*, 55, 1314–1324. <https://doi.org/10.1007/s13197-018-3042-0>
- Hedayati, S., Shahidi, F., Koocheki, A., Farahnaky, A., & Majzoobi, M. (2020). Influence of pregelatinized and granular cold water swelling maize starches on stability and physicochemical properties of low fat oil-in-water emulsions. *Food Hydrocolloids*, 102(November 2019). <https://doi.org/10.1016/j.foodhyd.2019.105620>
- Hong, L. F., Cheng, L. H., Lee, C. Y., & Peh, K. K. (2015). Characterisation of physicochemical properties of propionylated corn starch and its application as stabiliser. *Food Technology and Biotechnology*, 53(3), 278-285. <https://doi.org/10.17113/ftb.53.03.15.3907>
- Hong, L. F., Cheng, L. H., Gan, C. Y., Lee, C. Y., & Peh, K. K. (2018). Evaluation of starch propionate as emulsion stabiliser in comparison with octenylsuccinate starch. *LWT - Food Science and Technology*, 91(August 2017), 526-531. <https://doi.org/10.1016/j.lwt.2018.01.076>
- Irani, F., Ranjbar, Z., Moradian, S., & Jannesari, A. (2017). Microencapsulation of n-heptadecane phase change material with starch shell. *Progress in Organic Coatings*, 113(May), 31-38. <https://doi.org/10.1016/j.porgcoat.2017.08.006>

- Jading, A., Tethool, E. F., Payung, P., & Gultom, O. (2011). Karakteristik fisikokimia pati sagu hasil pengeringan secara fluidisasi menggunakan alat pengering cross flow fluidized bed bertenaga surya dan biomassa. *Jurnal Reaktor*, 13(3), 155-164.
- Jian Jin, Hongyun Li, Kaiqi Zhao, Jun Su, Lu Xia, Xiaohu Yuan, Guangming Jiang, Bo Xu, and Lidong Sun. (2022). Dissecting the Chain Length Effect on Separation of Alkane-in-Water Emulsions with Superwetting Microchannels. *ACS Applied Materials & Interfaces*, 14 (4), 6157-6166.
- Kang, X., Liu, P., Gao, W., Wu, Z., Yu, B., Wang, R., Cui, B., Qiu, L., & Sun, C. (2020). Preparation of starch-lipid complex by ultrasonication and its film forming capacity. *Food Hydrocolloids*, 99, 105340-105340. <https://doi.org/10.1016/j.foodhyd.2019.105340>
- Kasemwong, K., & Itthisoponkul, T. (2013). Encapsulation of flavor compounds as helical inclusion complexes of starch. *Advances in Applied Nanotechnology for Agriculture, Chapter 14, ACS Symposium Series, Vol. 1143*. 235-245, doi: 10.1021/bk-2013-1143.ch014
- Kasprzak, M. M., Macnaughtan, W., Harding, S., Wilde, P., & Wolf, B. (2018). Stabilisation of oil-in-water emulsions with non-chemical modified gelatinised starch. *Food Hydrocolloids*, 81, 409-418. <https://doi.org/10.1016/j.foodhyd.2018.03.002>.
- Keatkrai, J., Lumdubwong, N., Chaiseri, S., & Jirapakkul, W. (2017). Characteristics of menthone encapsulated complex by mungbean, tapioca, and rice starches. *International Journal of Food Properties*, 20(4), 810-820. <https://doi.org/10.1080/10942912.2016.1183129>
- Kibar, E. A. A., Gönenç, İ., & Us, F. (2014). Effects of fatty acid addition on the physicochemical properties of corn starch. *International Journal of Food Properties*, 17(1), 204-218. <https://doi.org/10.1080/10942912.2011.619289>
- Kim, J.-H., Decker, E. A., & McClements D. J. (2006). Preparation of multiple emulsions based on thermodynamic incompatibility of heat-denatured whey protein and pectin solutions. *Food Hydrocolloids*, 20, 586–595. <https://doi.org/10.1016/j.foodhyd.2005.06.007>
- Kim, J.-Y., & Huber, K. C. (2016). Preparation and characterization of corn starch- β -karoten composites. *Carbohydrate Polymers*, 136, 394-401.
- Kong, L., Bhosale, R., & Ziegler, G. R. (2018). Encapsulation and stabilization of β -karoten by amylose inclusion complexes. *Food Research International*, 105, 446-452.
- Królikowska, K., Fortuna, T., Pietrzyk, S., Gryszkin, A. (2017). Effect of modification of octenyl succinate starch with mineral elements on the stability and rheological properties of oil-in-water emulsions. *Food Hydrocolloids*, 66(2017), 118-127. <https://doi.org/10.1016/j.foodhyd.2016.12.012>

- Le-Bail, P., Lorentz, C., Pencreac'h, G., Saultani-Vigneron, S., Pontoire, B., Giraldo, L. J. L., Villeneuve, P., J. Hendrickx, J. & Tran, V. (2015). Trapping by amylose of the aliphatic chain grafted onto chlorogenic acid: Importance of the graft position. *Carbohydrate Polymers*, 117, 910–916. <http://dx.doi.org/10.1016/j.carbpol.2014.10.029>
- Lee, H. S., Kim, K. H., Park, S. H., Hur, S. W., & Auh, J. H. (2020). Amylose-lipid complex as a fat replacement in the preparation of low-fat white pan bread. *Foods*, 9(2), 1-12. <https://doi.org/10.3390/foods9020194>
- Lesmes, U., Cohen, S. H., Shener, Y., & Shimoni, E. (2009). Effects of long chain fatty acid unsaturation on the structure and controlled release properties of amylose complexes. *Food Hydrocolloids*, 23(3), 667-675. <https://doi.org/10.1016/j.foodhyd.2008.04.003>
- Li, C., Li, Y., Sun, P., & Yang, C. (2013). Pickering emulsions stabilized by native starch granules. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 431, 142-149. <https://doi.org/10.1016/j.colsurfa.2013.04.025>
- Li, P., Hao Huang, Yuan Fang, Yu Wang, Da Som No, Rohil S. Bhatnagar, Alireza Abbaspourrad (2023). Interfacial engineering of clear emulsions: Surfactant hydrophobicity and the hidden role of chain structure. *Colloids and Surfaces A: Physicochemical and Engineering Aspects* . 676, 132242.
- Liang, R., Shoemaker, C. F., Yang, X., Zhong, F., Huang, Q. (2013). Stability and Bioaccessibility of β -Carotene in Nanoemulsions Stabilized by Modified Starches. *Journal of Agricultural and Food Chemistry*, 61(6) 1249–1257. <https://doi.org/10.1021/jf303967f>
- Lin, X., Zhang, X., Du, B., & Xu, B. (2023). Morphological, Structural, Thermal, Pasting, and Digestive Properties of Starches Isolated from Different Varieties of Rice: A Systematic Comparative Study. *Foods*, 12(24). <https://doi.org/10.3390/foods12244492>
- Lin, Q., Jiang, H., Li, X., McClements, D. J., Sang, S., Wang, J., Jiao, A., Jin, Z., Qiu, C. (2024). Encapsulation and protection of β -carotene in Pickering emulsions stabilized by chitosan-phytic acid-cyclodextrin nanoparticles. *Food Bioscience*, 59,(2024), 103845, <https://doi.org/10.1016/j.fbio.2024.103845>.
- Liu, Y., Chen, J., Luo, S., Li, C., Ye, J., Liu, C., & Gilbert, R. G. (2017). Physicochemical and structural properties of pregelatinized starch prepared by improved extrusion cooking technology. *Carbohydrate Polymers*, 175, 265-272. <https://doi.org/10.1016/j.carbpol.2017.07.084>
- Liu, P., Wang, R., Kang, X., Cui, B., & Yu, B. (2018). Effects of ultrasonic treatment on amylose-lipid complex formation and properties of sweet potato starch-based films. *Ultrasoniks - Sonochemistry*, 44, 215-222. <https://doi.org/10.1016/j.ultsonch.2018.02.029>
- Liu, P., Kang, X., Cui, B., Gao, W., Wu, Z., & Yu, B. (2019). Effects of amylose content and enzymatic debranching on the properties of maize starch-

- glycerol monolaurate complexes. *Carbohydrate Polymers*, 222, 1-8. <https://doi.org/10.1016/j.carbpol.2019.115000>
- Liu, Z., Wang, C., Liao, X., & Shen, Q. (2020). Measurement and comparison of multi-scale structure in heat and pressure treated corn starch granule under the same degree of gelatinization. *Food Hydrocolloids*, 108, 1-9. <https://doi.org/10.1016/j.foodhyd.2020.106081>
- Liu, Q., Guan, H., Guo, Y., Wang, D., Yang, Y., Ji, H., Jiao, A., & Jin, Z. (2024). Structure and in vitro digestibility of amylose–lipid complexes formed by an extrusion-debranching-complexing strategy. *Food Chemistry*, 437, 137950. <https://doi.org/10.1016/j.foodchem.2023.137950>
- Lopez-Silva, M., Bello-Pereza, L. A., Agama-Acevedoa, E., & Alvarez-Ramirez, J. (2019). Effect of amylose content in morphological, functional and emulsification properties of OSA modified corn starch. *Food Hydrocolloids*, 97, 1-8. <https://doi.org/10.1016/j.foodhyd.2019.105212>
- Lu, X., Liu, H., & Huang, Q. (2020). Fabrication and characterization of resistant starch stabilized Pickering emulsions. *Food Hydrocolloids*, 103, 105703-105703. <https://doi.org/10.1016/j.foodhyd.2020.105703>
- Majzoobi, M., Kaveh, Z., & Farahnaky, A. (2016). Effect of acetic acid on physical properties of pregelatinized wheat and corn starch gels. *Food Chemistry*, 196, 720-725. <https://doi.org/10.1016/j.foodchem.2015.09.096>
- Mao, L., Wang, D., Liu, F., & Gao, Y. (2018). Emulsion design for the delivery of β -karoten in complex food systems. *Critical Reviews in Food Science and Nutrition*, 58(5), 770-784.
- Marefati, A. (2018). Starch Pickering Emulsions: Process and Encapsulation Stability. [Doctoral Thesis (compilation), Lund University]. Department of Food Technology, Lund University.
- Marinopoulou, A., Papastergiadis, E., Raphaelides, S. N. & Kontominas, M. G. (2016). Structural characterization and thermal properties of amylose-fatty acid complexes prepared at different temperatures. *Food Hydrocolloids* 58 224-234. <http://dx.doi.org/10.1016/j.foodhyd.2016.02.034>
- McClements, D. J. (2007). Critical review of techniques and methodologies for characterization of emulsion stability. *Critical Reviews in Food Science and Nutrition*, 47(7), 611-649. <https://doi.org/10.1080/10408390701289292>
- Montgomery, D.C. 2020. Introduction to Statistical Quality Control, 8th ed. John Wiley & Sons, Inc., New York.
- Pawlik, A. K., Fryer, P. J., & Norton, I. T. (2013). Formulation Engineering of Food Emulsions. In J. E. Norton, P. J. Fryer, & I. T. Norton (Eds.). Wiley - Blackwell.
- Putseys, J. A., Derde, L. J., Lamberts, L., Östman, E., Bjorck, I. M., & Delcour, J. A. (2010). Functionality of short chain amylose-Lipid complexes in starch-Water systems and their impact on in vitro starch degradation. *Journal of*

- Agricultural and Food Chemistry*, 58(3), 1939-1945.
<https://doi.org/10.1021/jf903523h>
- Qi, W., Zhanga, Z., & Wu, T. (2020). Encapsulation of β -karoten in oleogel-in-water Pickering emulsion with improved stability and bioaccessibility. *International Journal of Biological Macromolecules*, 164, 1432-1442.
- Qin, R., Yu, J., Li, Y., Copeland, L., Wang, S., & Wang, S. (2019). Structural changes of starch-lipid complexes during postprocessing and their effect on in vitro enzymatic digestibility. *Journal of Agricultural & Food Chemistry*, 67, 1530-1536. <https://doi.org/10.1021/acs.jafc.8b06371>
- Reddy, C. K., Choi, S. M., Lee, D.-J., & Lim, S.-T. (2018). Complex formation between starch and stearic acid: Effect of enzymatic debranching for starch. *Food Chemistry*, 244, 136-142. <https://doi.org/10.1016/j.foodchem.2017.10.040>
- Salaün, F., Butstraen, C., & Devaux, E. (2016). Sol-Gel Microencapsulation Based on Pickering Emulsion. In Selcan Karakuş (ed). *Science and Technology Behind Nanoemulsions*. Intechopen (pp. 43-63).
- Sang, S. S., Xu, X., Zhu, X., & Narsimhan, G. (2021). Complexation of 26-Mer Amylose with Egg Yolk Lipids with Different Numbers of Tails Using a Molecular Dynamics Simulation. *Foods*, 10(10). <https://doi.org/10.3390/foods10102355>
- Santos, J., Trujillo-Cayado, L. A., Barquero, M., Calero, N. (2022) Influence of Type and Concentration of Biopolymer on β -Carotene Encapsulation Efficiency in Nanoemulsions Based on Linseed Oil. *Polymers (Basel)*, 14(21):4640. doi: 10.3390/polym14214640.
- Santoso, B., Sakakura, K., Naito, H., Ohmi, M., Nishimura, Y., Uchiyama, T., Itaya, A., Hisamatsu, M., Ehara, H., & Mishima, T. (2015). Effects of Micro Powder Milling on Physicochemical Properties of Sago Starch. *Journal of Applied Glycoscience*, 62(2), 73-80. https://doi.org/10.5458/jag.jag.jag-2015_008
- Santoso, B., Sakakura, K., Naito, H., Ohmi, M., Nishimura, Y., Uchiyama, T., Itaya, A., Hisamatsu, M., & Ehara, H. (2017). Effects of Wet Milling Sago Pith Waste on Yield and Physical Properties of Sago Starch. *International Journal of Engineering & Technology (IJET-IJENS)*, 17(02), 1-6.
- Santoso, B., Sarungallo, Z. L., & Dewi, A. M. P. (2021). Physicochemical and functional properties of spineless, short-spines, and long-spines sago starch. *Biodiversitas*, 22(1), 137-143. <https://doi.org/10.13057/biodiv/d220119>
- Saw, S. S., Shariffa, N. Y., Ruri, A. S., & Uthumporn, U. (2020). Physicochemical and emulsifying properties of pre-treated octenyl succinic anhydride (Osa) sago starch in simple emulsion system. *Food Research*, 4(4), 1326-1332. [https://doi.org/10.26656/fr.2017.4\(4\).426](https://doi.org/10.26656/fr.2017.4(4).426)

- Seo, T. R., Kim, J. Y., & Lim, S. T. (2015). Preparation and characterization of crystalline complexes between amylose and C18 fatty acids. *LWT - Food Science and Technology*, 64(2), 889-897. <https://doi.org/10.1016/j.lwt.2015.06.021>
- Shi, S., Dong, Y., Li, Q., Liu, T., & Yu, X. (2020). Morphology, structural, thermal and rheological properties of wheat starch-palmitic acid complexes prepared during steam cooking. *RSC Advances*, 10(50), 30087–30093. <https://doi.org/10.1039/d0ra05954d>
- Subagio, A., & Morita, N. (2001). Instability of carotenoids is a reason for their promotion on lipid oxidation. *Food Research International*, 34, (2–3), 183-188. [https://doi.org/10.1016/S0963-9969\(00\)00150-2](https://doi.org/10.1016/S0963-9969(00)00150-2).
- Sun, X., Jin, R., Ma, F., Ma, W., Pan, Y., Liu, J., Liu, X., Zhu, J., & Zhang, J. (2025). Effects of different fatty acids on the structure, physicochemical properties, and in vitro digestibility of Chinese yam resistant starch-lipid complexes. *Food Chemistry*, 465(2), 142159. <https://doi.org/10.1016/j.foodchem.2024.142159>
- Sweedman, M. C., Tizzotti, M. J., Schäfer, C., Gilbert, R. G. (2013). Structure and physicochemical properties of octenyl succinic anhydride modified starches: A review. *Carbohydrate Polymers*, 92(1) 905-920. <https://doi.org/10.1016/j.carbpol.2012.09.040>.
- Tang, M. C., & Copeland, L. (2007). Analysis of complexes between lipids and wheat starch. *Carbohydrate Polymers*, 67(1), 80-85. <https://doi.org/10.1016/j.carbpol.2006.04.016>
- Tethool, E. F., Jading, A., & Santoso, B. (2012). Characterization of Physicochemical and Baking Expansion Properties of Oxidized Sago Starch Using Hydrogen Peroxide and Sodium Hypochlorite Catalyzed By UV Irradiation. *Food Science and Quality Management*, 10, 1-11.
- Tethool, E. F., Santoso, B., & Dewi, A. M. P. (2019). *Teknologi Pengolahan Ubi-ubian dan Sagu* (1 ed.). Deepublish. Yogyakarta
- Timgren, A., Rayner, M., Dejmek, P., Marku, D., & Sjöo, M. (2013). Emulsion stabilizing capacity of intact starch granules modified by heat treatment or octenyl succinic anhydride. *Food Science & Nutrition*, 1(2), 157-171. <https://doi.org/10.1002/fsn3.17>
- Torrejon, M. V., Song, H., Wu, B., Luo, G., & Song, J. (2023). Effect of Starch Type and Pre-Treatment on the Properties of Gelatin–Starch Foams Produced by Mechanical Foaming. *Polymers*, 15(7), 1-18. <https://doi.org/10.3390/polym15071775>
- Tufvesson, F., Wahlgren, M., & Eliasson, A-C. (2003a). Formation of amylose-lipid complexes and effects of temperature treatment. Part 1. Monoglycerides. *Starch - Stärke*, 55(2), 61-71. <https://doi.org/10.1002/star.200390018>

- Tufvesson, F., Wahlgren, M., & Eliasson, A-C. (2003b). Formation of amylose-lipid complexes and effects of temperature treatment. Part 2. Fatty Acids. *Starch - Stärke*, 55(34), 138-149. <https://doi.org/10.1002/star.200390028>
- Vanmarcke, A., Leroy, L., Stoclet, G., Duchatel-Crépy, L., Lefebvre, J-M., Joly, N., & Gaucher, V. (2017) Influence of fatty chain length and starch composition on structure and properties of fully substituted fatty acid starch esters. *Carbohydrate Polymers* 164, 249–257. <http://dx.doi.org/10.1016/j.carbpol.2017.02.013>
- Vasiliadou, E., Raphaelides, S. N., & Papastergiadis, E. (2015). Effect of heating time and temperature on partially gelatinized starch fatty acid interactions. *LWT - Food Science and Technology*, 60, 698-707. <http://dx.doi.org/10.1016/j.lwt.2014.10.026>
- Wang, S., Wang, J., Yu, J., & Wang, S. (2016). Effect of fatty acids on functional properties of normal wheat and waxy wheat starches: A structural basis. *Food Chemistry*, 190, 285-292. <https://doi.org/10.1016/j.foodchem.2015.05.086>
- Wang, L., Wang, W., Wang, Y., Xiong, G., Mei, X., Wu, W., Ding, A., Li, X., Qiao, Y., & Liao, L. (2018). Effects of fatty acid chain length on properties of potato starch–fatty acid complexes under partially gelatinization. *International Journal of Food Properties*, 21(1), 2121-2134. <https://doi.org/10.1080/10942912.2018.1489842>
- Wang, R., Liu, P., Cui, B., Kang, X., & Yu, B. (2019a). Effects of different treatment methods on properties of potato starch-lauric acid complex and potato starch-based films. *International Journal of Biological Macromolecules*, 124, 34-40. <https://doi.org/10.1016/j.ijbiomac.2018.11.207>
- Wang, H., Wu, Y., Wang, N., Yang, L., & Zhou, Y. (2019b). Effect of water content of high-amylose corn starch and glutinous rice starch combined with lipids on formation of starch–lipid complexes during deep-fat frying. *Food Chemistry*, 278, 515-522. <https://doi.org/10.1016/j.foodchem.2018.11.092>
- Wang, S., Wang, S., Chao, C., Cai, J., Niu, B., & Copeland, L. (2020). Starch – lipid and starch – lipid – protein complexes : A review. *Comprehensive Reviews in Food Science and Food Safety*. Wiley. 1-24. <https://doi.org/10.1111/1541-4337.12550>
- Wulff, G., Avgenaki, G., & Guzman, M. S. P. (2005). Molecular encapsulation of flavours as helical inclusion complexes of amylose. *Journal of Cereal Science*, 41, 239–249. <https://doi.org/10.1016/j.jcs.2004.06.002>
- Xu, T., Yang, J., Hua, S., Hong, Y., Gu, Z., Cheng, L., Li, Z., & Li, C. (2020). Characteristics of starch-based Pickering emulsions from the interface perspective. *Trends in Food Science and Technology*, 105(September), 334-346. <https://doi.org/10.1016/j.tifs.2020.09.026>

- Yan, C., Wu, X., Wang, Y., Peng, S., Chen, J., Zou, L., McClements, D. J., & Liu, W. (2021). Utilization of polysaccharide-based high internal phase emulsion for nutraceutical encapsulation: Enhancement of carotenoid loading capacity and stability. *Journal of Functional Foods*, *84*, (2021), 104601. <https://doi.org/10.1016/j.jff.2021.104601>.
- Yang, Y., Fang, Z., Chen, X., Zhang, W., Xie, Y., Chen, Y., Liu, Z., & Yuan, W. (2017). An overview of pickering emulsions: Solid-particle materials, classification, morphology, and applications. *Frontiers in Pharmacology*, *8*(MAY), 1-20. <https://doi.org/10.3389/fphar.2017.00287>
- Yotsawimonwat, S., Sriroth, K., Kaewvichit, S., Piyachomkwan, P., Jane, J.-L., & Sirithunyalug, J. (2008). Effect of pH on complex formation between debranched waxy rice starch and fatty acids. *International Journal of Biological Macromolecules*, *43*(2), 94–99. <https://doi.org/10.1016/j.ijbiomac.2008.03.006>.
- Yulianingsih, R., & Gohtani, S. (2019). Dispersion characteristics of pregelatinized waxy rice starch and its performance as an emulsifier for oil-in-water emulsions: Effect of gelatinization temperature and starch concentration. *Food Hydrocolloids*, *95*(December 2018), 476-486. <https://doi.org/10.1016/j.foodhyd.2018.12.013>
- Zheng, M., Chao, C., Yu, J., Copeland, L., Wang, S., & Wang, S. (2018). Effects of Chain Length and Degree of Unsaturation of Fatty Acids on Structure and in Vitro Digestibility of Starch-Protein-Fatty Acid Complexes. *Journal of Agricultural and Food Chemistry*, *66*(8), 1872-1880. <https://doi.org/10.1021/acs.jafc.7b04779>
- Zheng, Y., Wang, B., Guo, Z., Zhang, Y., Zheng, B., Zeng, S., & Zeng, H. (2020). Properties of lotus seed starch-glycerin monostearin V-complexes after long-term retrogradation. *Food Chemistry*, *311*(September 2019), 125887-125887. <https://doi.org/10.1016/j.foodchem.2019.125887>
- Zhu, F. (2019). Starch based Pickering emulsions: Fabrication, properties, and applications. *Trends in Food Science and Technology*, *85*(January), 129-137. <https://doi.org/10.1016/j.tifs.2019.01.012>