

DAFTAR PUSTAKA

- Acharya, P., Chouhan, K., Weiskirchen, S., & Weiskirchen, R. (2021). Cellular mechanisms of liver fibrosis. *Frontiers in Pharmacology*, 12, 671640.
- Aricandana, P.A.E., Putra, I.G.M., Megadhana, I.W., Anantasika, A.A.N., Manuaba, I.B.G.F. and Surya, I.G.N.H.W., 2022. Polimorfisme gen COL1A1 sebagai faktor risiko terjadinya prolaps organ panggul pada perempuan Bali, Indonesia. *Intisari Sains Medis*, 13(1), pp.191-196.
- Asai, A., Miethke, A. and Bezerra, J.A., 2015. Pathogenesis of biliary atresia: defining biology to understand clinical phenotypes. *Nature reviews Gastroenterology & hepatology*, 12(6), pp.342-352.
- Brahee, D. D., & Lampl, B. S. (2022). Neonatal diagnosis of biliary atresia: a practical review and update. *Pediatric radiology*, 52(4), 685-692.
- Chardot, C., Carton, M., Spire-Bendelac, N., Le Pommelet, C., Golmard, J. L., Reding, R., & Auvert, B. (2001). Is the Kasai operation still indicated in children older than 3 months diagnosed with biliary atresia?. *The Journal of pediatrics*, 138(2), 224-228.
- Chen, Y., Gilbert, M. A., Grochowski, C. M., McEldrew, D., Llewellyn, J., Waisbourd-Zinman, O., ... & Devoto, M. (2018). A genome-wide association study identifies a susceptibility locus for biliary atresia on 2p16. 1 within the gene EFEMP1. *PLoS genetics*, 14(8), e1007532.

Hopkins, P.C., Yazigi, N. and Nylund, C.M., 2017. Incidence of biliary atresia and timing of hepatoportoenterostomy in the United States. *The Journal of pediatrics*, 187, pp.253-257.

Hsiao, C.H., Chang, M.H., Chen, H.L., Lee, H.C., Wu, T.C., Lin, C.C., Yang, Y.J., Chen, A.C., Tiao, M.M., Lau, B.H. and Chu, C.H., 2008. Universal screening for biliary atresia using an infant stool color card in Taiwan. *Hepatology*, 47(4), pp.1233-1240.

Itoh S, Imagawa M, Kokubo T, *et al.* Potential Role of Leukocyte Cell-Derived Chemotaxin 2 as a Fibrosis Prognostic Marker in Nonalcoholic Fatty Liver Disease. *Biomed Res Int.* 2019;2019:7514262. doi:10.1155/2019/7514262

Kakos, C. D., Ziogas, I. A., Alexopoulos, S. P., & Tsoulfas, G. (2021). Management of biliary atresia: To transplant or not to transplant. *World Journal of Transplantation*, 11(9), 400.

Kong, M., & Xiang, B. (2021). Identifying Biomarkers to Predict the Prognosis of Biliary Atresia by Weighted Gene Co-Expression Network Analysis. *Frontiers in genet*, 12, 760182. doi: 10.3389/fgene.2021.760182

Kitagawa, H. (2021). Definition and classifications of biliary atresia. *Introduction to Biliary Atresia*, 11-17.

Lee, K.J., Kim, J.W., Moon, J.S. and Ko, J.S., 2017. Epidemiology of biliary atresia in Korea. *Journal of Korean Medical Science*, 32(4), p.656.

Lenhardt, R., Hopf, H.W., Marker, E., Akça, O., Kurz, A., Scheuenstuhl, H. and Sessler, D.I., 2000. Perioperative collagen deposition in elderly and young men and women. *Archives of surgery*, 135(1), pp.71-74.

Livesey, E., Borja, M.C., Sharif, K., Alizai, N., McClean, P., Kelly, D., Hadzic, N. and Davenport, M., 2009. Epidemiology of biliary atresia in England and Wales (1999–2006). *Archives of Disease in Childhood-Fetal and Neonatal Edition*, 94(6), pp.F451-F455.

Lorent K, Gong W, Koo KA *et al.* Identification of a plant isoflavonoid that causes biliary atresia. *Sci. Transl. Med.* 7(286), 286ra67 (2015).

Li, Jingxiong & Cen, Bohong & Chen, Siping & He, Yuanli. 2016. MicroRNA-29b inhibits TGF- β 1-induced fibrosis via regulation of the TGF- β 1/Smad pathway in primary human endometrial stromal cells. *Molecular Medicine Reports*. 13. doi: 10.3892/mmr.2016.5062.

Ma, H. P., Chang, H. L., Bamodu, O. A., Yadav, V. K., Huang, T. Y., Wu, A. T., ... & Lee, W. H. (2019). Collagen 1A1 (*COL1A1*) is a reliable biomarker and putative therapeutic target for hepatocellular carcinogenesis and metastasis. *Cancers*, 11(6), 786.

Makhmudi, A., Kalim, A.S., Gunadi, 2019. microRNA-21 expressions impact on liver fibrosis in biliary atresia patients. *BMC Notes* 12. <https://doi.org/10.1186/S13104-019-4227-Y>

Makhmudi, A., Supanji, R., Putra, B. P., & Gunadi. (2020). The effect of APTR, Fn14 and CD133 expressions on liver fibrosis in biliary atresia patients. *Pediatric Surgery International*, 36, 75-79.

Nio, M., Ohi, R., Miyano, T., Saeki, M., Shiraki, K. and Tanaka, K., 2003. Five- and 10-year survival rates after surgery for biliary atresia: a report from the

Japanese Biliary Atresia Registry. *Journal of pediatric surgery*, 38(7), pp.997-1000.

Pradana D.A., 2022. Hubungan Ekspresi *LECT2* dengan Liver Fibrosis pada Pasien Atresia Bilier. Tesis. Perpustakaan Universitas Gadjah Mada.

Sanchez-Valle, A., Kassira, N., Varela, V. C., Radu, S. C., Paidas, C., & Kirby, R. S. (2017). Biliary atresia: epidemiology, genetics, clinical update, and public health perspective. *Advances in Pediatrics*, 64(1), 285-305.

Sangkhatat S, Laochareonsuk W, Maneechay W, Kayasut K, Chiengkriwate P. Variants associated with infantile cholestatic syndromes detected in extrahepatic biliary atresia by whole exome studies: a 20-case series from Thailand. *J. Pediatr. Genet.* 7(2), 67–73 (2018).

Schreiber, R.A., Harpavat, S., Hulscher, J.B.F., Wildhaber, B.E. (2022). Biliary Atresia in 2021: Epidemiology, Screening, and Public Policy, *J Clin Med*, 11 (4):999.

Shinkai, M., Ohhama, Y., Take, H., Kitagawa, N., Kudo, H., Mochizuki, K., & Hatata, T. (2009). Long-term outcome of children with biliary atresia who were not transplanted after the Kasai operation:> 20-year experience at a children's hospital. *Journal of pediatric gastroenterology and nutrition*, 48(4), 443-450.

Wang H, Zhang Y, Zhang Z, *et al.* *LECT2*, a Ligand for Tie1, Plays a Crucial Role in Liver Fibrogenesis. *Cell.* 2019;178(6):1478-1492.e20. doi:10.1016/j.cell.2019.08.022

Waisbourd-Zinman O, Koh H, Tsai S *et al.* The toxin biliatresone causes mouse extrahepatic cholangiocyte damage and fibrosis through decreased glutathione and SOX17. *Hepatology* 64(3), 880–893 (2016).

Wildhaber, B.E., Majno, P., Mayr, J., Zachariou, Z., Hohlfeld, J., Schwoebel, M., Kistler, W., Meuli, M., Le Coultre, C., Mentha, G. and Belli, D., 2008. Biliary atresia: Swiss national study, 1994–2004. *Journal of pediatric gastroenterology and nutrition*, 46(3), pp.299-307.

Wildhaber, B.E., Coran, A.G., Drongowski, R.A., Hirschl, R.B., Geiger, J.D., Lelli, J.L. and Teitelbaum, D.H., 2003. The Kasai portoenterostomy for biliary atresia: a review of a 27-year experience with 81 patients. *Journal of pediatric surgery*, 38(10), pp.1480-1485.

Vij, M. and Rela, M., 2020. Biliary atresia: pathology, etiology and pathogenesis. *Future science OA*, 6(5), p.FSO466.

Xu Y, Yu J, Zhang R *et al.* The perinatal infection of cytomegalovirus is an important etiology for biliary atresia in China. *Clin. Pediatr. (Phila)*. 51(2), 109–113 (2012).

Xu, M., Xu, H. H., Lin, Y., Sun, X., Wang, L. J., Fang, Z. P., ... & Zhou, W. J. (2019). *LECT2*, a ligand for Tie1, plays a crucial role in liver fibrogenesis. *Cell*, 178(6), 1478-1492.

Xu, H., Li, X., Wu, Z., Zhao, L., Shen, J., Liu, J., ... & Gao, Y. (2021). *LECT2*, a novel and direct biomarker of liver fibrosis in patients with CHB. *Frontiers in Molecular Biosciences*, 8, 749648.

Yang L, Inokuchi S, Roh YS, *et al.* Transforming growth factor- β signaling in hepatocytes promotes hepatic fibrosis and carcinogenesis in mice with hepatocyte-specific deletion of TAK1. *Gastroenterology*. 2013;144(5):1042-1054.e4.

Zhang S, Goswami S, Ma J *et al.* CD4(+)T cell subset profiling in biliary atresia reveals ICOS(-) regulatory T cells as a favorable prognostic factor. *Front. Pediatr.* 7, 279 (2019).